



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 15, 2017 – 07:08 pm GMT

PDB ID : 2YBV  
Title : STRUCTURE OF RUBISCO FROM THERMOSYNECHOCOCCUS ELONGATUS  
Authors : Terlecka, B.; Wilhelmi, V.; Bialek, W.; Gubernator, B.; Szczepaniak, A.; Hofmann, E.  
Deposited on : 2011-03-10  
Resolution : 2.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.9-1692
EDS	:	trunk28620
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	trunk28620


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	4130 (2.30-2.30)
Clashscore	112137	4751 (2.30-2.30)
Ramachandran outliers	110173	4705 (2.30-2.30)
Sidechain outliers	110143	4704 (2.30-2.30)
RSRZ outliers	101464	4156 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	475	<div> <div>%</div> <div> <div></div> <div>80%</div> <div>10%</div> <div>10%</div> </div> </div>
1	C	475	<div> <div>%</div> <div> <div></div> <div>79%</div> <div>10%</div> <div>10%</div> </div> </div>
1	E	475	<div> <div>%</div> <div> <div></div> <div>77%</div> <div>12%</div> <div>• 10%</div> </div> </div>
1	G	475	<div> <div></div> <div> <div>79%</div> <div>10%</div> <div>10%</div> </div> </div>
1	I	475	<div> <div>2%</div> <div> <div></div> <div>78%</div> <div>12%</div> <div>• 10%</div> </div> </div>
1	K	475	<div> <div>%</div> <div> <div></div> <div>78%</div> <div>12%</div> <div>10%</div> </div> </div>

Continued on next page...

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
1	M	475	<p>77% 12% • 10%</p>
1	O	475	<p>78% 11% • 10%</p>
2	B	118	<p>75% 11% • 12%</p>
2	D	118	<p>65% 21% • 12%</p>
2	F	118	<p>65% 20% • 12%</p>
2	H	118	<p>72% 16% 12%</p>
2	J	118	<p>65% 21% • 12%</p>
2	L	118	<p>69% 19% • 12%</p>
2	N	118	<p>77% 10% • 12%</p>
2	P	118	<p>67% 19% • 12%</p>

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 34583 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called RIBULOSE BISPHOSPHATE CARBOXYLASE LARGE CHAIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	428	Total	C	N	O	S	0	0	0
			3348	2123	588	616	21			
1	C	428	Total	C	N	O	S	0	0	0
			3348	2123	588	616	21			
1	E	428	Total	C	N	O	S	0	0	0
			3348	2123	588	616	21			
1	G	428	Total	C	N	O	S	0	0	0
			3348	2123	588	616	21			
1	I	428	Total	C	N	O	S	0	0	0
			3348	2123	588	616	21			
1	K	428	Total	C	N	O	S	0	0	0
			3348	2123	588	616	21			
1	M	428	Total	C	N	O	S	0	0	0
			3348	2123	588	616	21			
1	O	428	Total	C	N	O	S	0	0	0
			3348	2123	588	616	21			

- Molecule 2 is a protein called RIBULOSE BISPHOSPHATE CARBOXYLASE SMALL SUB-UNIT.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	104	Total	C	N	O	S	0	0	0
			869	557	146	159	7			
2	D	104	Total	C	N	O	S	0	0	0
			869	557	146	159	7			
2	F	104	Total	C	N	O	S	0	0	0
			869	557	146	159	7			
2	H	104	Total	C	N	O	S	0	0	0
			869	557	146	159	7			
2	J	104	Total	C	N	O	S	0	0	0
			869	557	146	159	7			

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	L	104	Total 869	C 557	N 146	O 159	S 7	0	0	0
2	N	104	Total 869	C 557	N 146	O 159	S 7	0	0	0
2	P	104	Total 869	C 557	N 146	O 159	S 7	0	0	0

- Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	G	2	Total 2	Cl 2	0	0
3	K	2	Total 2	Cl 2	0	0
3	E	1	Total 1	Cl 1	0	0
3	I	2	Total 2	Cl 2	0	0
3	C	2	Total 2	Cl 2	0	0
3	A	2	Total 2	Cl 2	0	0
3	O	2	Total 2	Cl 2	0	0
3	M	2	Total 2	Cl 2	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	106	Total 106	O 106	0	0
4	B	3	Total 3	O 3	0	0
4	C	119	Total 119	O 119	0	0
4	D	7	Total 7	O 7	0	0
4	E	118	Total 118	O 118	0	0
4	F	8	Total 8	O 8	0	0

*Continued on next page...*


*Continued from previous page...*

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	G	112	Total 112	O 112	0	0
4	H	4	Total 4	O 4	0	0
4	I	87	Total 87	O 87	0	0
4	J	16	Total 16	O 16	0	0
4	K	73	Total 73	O 73	0	0
4	L	7	Total 7	O 7	0	0
4	M	77	Total 77	O 77	0	0
4	N	9	Total 9	O 9	0	0
4	O	80	Total 80	O 80	0	0
4	P	6	Total 6	O 6	0	0




- Molecule 1: RIBULOSE BISPHOSPHATE CARBOXYLASE LARGE CHAIN






• Molecule 1: RIBULOSE BISPHOSPHATE CARBOXYLASE LARGE CHAIN

Chain G: 79% 10% 10%




• Molecule 1: RIBULOSE BISPHOSPHATE CARBOXYLASE LARGE CHAIN

Chain I: 2% 78% 12% 10%



• Molecule 1: RIBULOSE BISPHOSPHATE CARBOXYLASE LARGE CHAIN

Chain K: % 78% 12% 10%




• Molecule 1: RIBULOSE BISPHOSPHATE CARBOXYLASE LARGE CHAIN


Chain M: % 77% 12% 10%









• Molecule 1: RIBULOSE BISPHOSPHATE CARBOXYLASE LARGE CHAIN



• Molecule 2: RIBULOSE BISPHOSPHATE CARBOXYLASE SMALL SUBUNIT



• Molecule 2: RIBULOSE BISPHOSPHATE CARBOXYLASE SMALL SUBUNIT




• Molecule 2: RIBULOSE BISPHOSPHATE CARBOXYLASE SMALL SUBUNIT








• Molecule 2: RIBULOSE BISPHOSPHATE CARBOXYLASE SMALL SUBUNIT




• Molecule 2: RIBULOSE BISPHOSPHATE CARBOXYLASE SMALL SUBUNIT



• Molecule 2: RIBULOSE BISPHOSPHATE CARBOXYLASE SMALL SUBUNIT



• Molecule 2: RIBULOSE BISPHOSPHATE CARBOXYLASE SMALL SUBUNIT



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	115.48Å 163.47Å 163.54Å 90.00° 108.96° 90.00°	Depositor
Resolution (Å)	39.84 – 2.30 39.83 – 2.30	Depositor EDS
% Data completeness (in resolution range)	100.0 (39.84-2.30) 100.0 (39.83-2.30)	Depositor EDS
$R_{merge}$	0.17	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.08 (at 2.29Å)	Xtriage
Refinement program	REFMAC 5.5.0088	Depositor
R, $R_{free}$	0.196 , 0.232 0.196 , 0.233	Depositor DCC
$R_{free}$ test set	12993 reflections (5.39%)	DCC
Wilson B-factor (Å <sup>2</sup> )	11.8	Xtriage
Anisotropy	0.532	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 14.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.40$ , $\langle L^2 \rangle = 0.22$	Xtriage
Estimated twinning fraction	0.057 for h,-k,-h-l	Xtriage
$F_o, F_c$ correlation	0.91	EDS
Total number of atoms	34583	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	18.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.15% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section:  
CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	1.12	9/3427 (0.3%)	0.99	10/4642 (0.2%)
1	C	1.11	8/3427 (0.2%)	0.93	7/4642 (0.2%)
1	E	1.11	11/3427 (0.3%)	0.97	9/4642 (0.2%)
1	G	1.11	7/3427 (0.2%)	0.99	9/4642 (0.2%)
1	I	1.07	6/3427 (0.2%)	0.96	6/4642 (0.1%)
1	K	1.09	5/3427 (0.1%)	0.93	7/4642 (0.2%)
1	M	1.09	9/3427 (0.3%)	0.94	10/4642 (0.2%)
1	O	1.08	3/3427 (0.1%)	0.92	7/4642 (0.2%)
2	B	0.90	0/893	0.85	0/1215
2	D	0.94	0/893	0.83	0/1215
2	F	0.99	2/893 (0.2%)	0.86	0/1215
2	H	0.98	1/893 (0.1%)	0.85	0/1215
2	J	1.02	0/893	0.84	0/1215
2	L	0.95	0/893	0.86	0/1215
2	N	0.91	0/893	0.83	0/1215
2	P	0.94	0/893	0.81	0/1215
All	All	1.07	61/34560 (0.2%)	0.93	65/46856 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	G	0	1

The worst 5 of 61 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	K	172	CYS	CB-SG	10.14	1.99	1.82
1	M	218	PHE	CE2-CZ	-9.31	1.19	1.37

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	139	ARG	CZ-NH1	-8.98	1.21	1.33
1	M	218	PHE	CG-CD2	-8.66	1.25	1.38
1	A	139	ARG	CZ-NH2	-8.63	1.21	1.33

The worst 5 of 65 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	G	215	ARG	NE-CZ-NH1	19.10	129.85	120.30
1	I	421	ARG	NE-CZ-NH1	15.58	128.09	120.30
1	E	139	ARG	NE-CZ-NH2	14.95	127.77	120.30
1	M	421	ARG	NE-CZ-NH1	13.97	127.28	120.30
1	A	139	ARG	NE-CZ-NH2	13.84	127.22	120.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	G	215	ARG	Sidechain

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3348	0	3273	35	2
1	C	3348	0	3273	41	2
1	E	3348	0	3273	44	0
1	G	3348	0	3273	37	0
1	I	3348	0	3273	45	0
1	K	3348	0	3273	45	0
1	M	3348	0	3273	47	0
1	O	3348	0	3273	44	0
2	B	869	0	836	12	0
2	D	869	0	836	22	0
2	F	869	0	836	18	0
2	H	869	0	836	10	0
2	J	869	0	836	22	0
2	L	869	0	836	19	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	N	869	0	836	16	0
2	P	869	0	836	21	0
3	A	2	0	0	0	0
3	C	2	0	0	0	0
3	E	1	0	0	0	0
3	G	2	0	0	0	0
3	I	2	0	0	0	0
3	K	2	0	0	0	0
3	M	2	0	0	0	0
3	O	2	0	0	0	0
4	A	106	0	0	2	0
4	B	3	0	0	0	0
4	C	119	0	0	3	0
4	D	7	0	0	1	0
4	E	118	0	0	4	0
4	F	8	0	0	0	0
4	G	112	0	0	3	0
4	H	4	0	0	0	0
4	I	87	0	0	2	0
4	J	16	0	0	0	0
4	K	73	0	0	4	0
4	L	7	0	0	0	0
4	M	77	0	0	0	0
4	N	9	0	0	0	0
4	O	80	0	0	5	0
4	P	6	0	0	0	0
All	All	34583	0	32872	412	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 412 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:49:PRO:HD2	1:C:52:GLU:HG3	1.37	1.06
1:O:177:LYS:O	1:O:206:ILE:HD11	1.63	0.99
4:C:2011:HOH:O	1:K:178:LEU:HD12	1.63	0.98
1:I:49:PRO:HD2	1:I:52:GLU:HG3	1.45	0.98
2:D:51:ARG:HD2	1:E:223:ASP:OD2	1.63	0.98

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the sym-

metry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:439:ARG:NH2	1:C:52:GLU:OE2[2_656]	2.07	0.13
1:A:52:GLU:OE2	1:C:439:ARG:NH2[2_656]	2.18	0.02

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	422/475 (89%)	412 (98%)	10 (2%)	0	100	100
1	C	422/475 (89%)	410 (97%)	12 (3%)	0	100	100
1	E	422/475 (89%)	411 (97%)	11 (3%)	0	100	100
1	G	422/475 (89%)	411 (97%)	11 (3%)	0	100	100
1	I	422/475 (89%)	414 (98%)	8 (2%)	0	100	100
1	K	422/475 (89%)	412 (98%)	10 (2%)	0	100	100
1	M	422/475 (89%)	411 (97%)	10 (2%)	1 (0%)	51	63
1	O	422/475 (89%)	411 (97%)	11 (3%)	0	100	100
2	B	102/118 (86%)	95 (93%)	4 (4%)	3 (3%)	5	3
2	D	102/118 (86%)	95 (93%)	4 (4%)	3 (3%)	5	3
2	F	102/118 (86%)	92 (90%)	8 (8%)	2 (2%)	9	7
2	H	102/118 (86%)	96 (94%)	5 (5%)	1 (1%)	18	20
2	J	102/118 (86%)	94 (92%)	7 (7%)	1 (1%)	18	20
2	L	102/118 (86%)	92 (90%)	8 (8%)	2 (2%)	9	7
2	N	102/118 (86%)	92 (90%)	9 (9%)	1 (1%)	18	20
2	P	102/118 (86%)	92 (90%)	10 (10%)	0	100	100
All	All	4192/4744 (88%)	4040 (96%)	138 (3%)	14 (0%)	44	55

5 of 14 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	62	ASN
2	D	62	ASN
2	F	93	ILE
2	J	62	ASN
2	L	93	ILE

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	346/386 (90%)	341 (99%)	5 (1%)	71	85
1	C	346/386 (90%)	341 (99%)	5 (1%)	71	85
1	E	346/386 (90%)	339 (98%)	7 (2%)	60	77
1	G	346/386 (90%)	340 (98%)	6 (2%)	66	81
1	I	346/386 (90%)	342 (99%)	4 (1%)	75	87
1	K	346/386 (90%)	341 (99%)	5 (1%)	71	85
1	M	346/386 (90%)	341 (99%)	5 (1%)	71	85
1	O	346/386 (90%)	341 (99%)	5 (1%)	71	85
2	B	97/108 (90%)	95 (98%)	2 (2%)	59	76
2	D	97/108 (90%)	92 (95%)	5 (5%)	27	36
2	F	97/108 (90%)	93 (96%)	4 (4%)	35	48
2	H	97/108 (90%)	96 (99%)	1 (1%)	80	90
2	J	97/108 (90%)	94 (97%)	3 (3%)	45	61
2	L	97/108 (90%)	95 (98%)	2 (2%)	59	76
2	N	97/108 (90%)	95 (98%)	2 (2%)	59	76
2	P	97/108 (90%)	95 (98%)	2 (2%)	59	76
All	All	3544/3952 (90%)	3481 (98%)	63 (2%)	64	79

5 of 63 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	G	52	GLU

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type
1	I	52	GLU
1	O	163	ASN
1	G	163	ASN
1	G	215	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 12 such sidechains are listed below:

Mol	Chain	Res	Type
1	K	163	ASN
1	K	226	HIS
2	N	97	GLN
2	J	65	ASN
2	N	65	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 15 ligands modelled in this entry, 15 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	428/475 (90%)	-0.28	4 (0%) 84 87	2, 11, 39, 53	0
1	C	428/475 (90%)	-0.30	4 (0%) 84 87	2, 11, 39, 53	0
1	E	428/475 (90%)	-0.30	3 (0%) 87 90	2, 11, 39, 53	0
1	G	428/475 (90%)	-0.27	2 (0%) 90 93	2, 11, 38, 53	0
1	I	428/475 (90%)	-0.31	9 (2%) 64 70	2, 11, 39, 53	0
1	K	428/475 (90%)	-0.30	7 (1%) 72 77	2, 11, 39, 53	0
1	M	428/475 (90%)	-0.32	6 (1%) 75 80	2, 11, 39, 53	0
1	O	428/475 (90%)	-0.29	10 (2%) 61 67	2, 11, 39, 53	0
2	B	104/118 (88%)	0.13	5 (4%) 31 38	19, 33, 52, 64	0
2	D	104/118 (88%)	0.17	3 (2%) 52 59	17, 32, 48, 59	0
2	F	104/118 (88%)	-0.01	3 (2%) 52 59	15, 27, 44, 57	0
2	H	104/118 (88%)	0.08	3 (2%) 52 59	16, 28, 51, 62	0
2	J	104/118 (88%)	0.08	3 (2%) 52 59	15, 30, 47, 59	0
2	L	104/118 (88%)	0.20	3 (2%) 52 59	17, 32, 52, 63	0
2	N	104/118 (88%)	0.13	1 (0%) 82 86	18, 33, 53, 62	0
2	P	104/118 (88%)	0.08	3 (2%) 52 59	15, 31, 52, 63	0
All	All	4256/4744 (89%)	-0.22	69 (1%) 72 77	2, 14, 42, 64	0

The worst 5 of 69 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	I	332	VAL	4.1
2	L	60	LEU	3.9
1	C	451	TRP	3.8
1	O	72	ASP	3.8
1	I	74	LEU	3.8

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
3	CL	A	1460	1/1	0.95	0.10	-	36,36,36,36	0
3	CL	I	1461	1/1	0.94	0.11	-	34,34,34,34	0
3	CL	K	1461	1/1	0.96	0.18	-	31,31,31,31	0
3	CL	M	1461	1/1	0.95	0.14	-	34,34,34,34	0
3	CL	O	1461	1/1	0.94	0.12	-	35,35,35,35	0
3	CL	A	1461	1/1	0.98	0.07	-	33,33,33,33	0
3	CL	K	1460	1/1	0.98	0.14	-	27,27,27,27	0
3	CL	C	1461	1/1	0.97	0.08	-	36,36,36,36	0
3	CL	M	1460	1/1	0.99	0.13	-	30,30,30,30	0
3	CL	O	1460	1/1	0.96	0.14	-	32,32,32,32	0
3	CL	G	1461	1/1	0.98	0.08	-	30,30,30,30	0
3	CL	C	1460	1/1	0.98	0.15	-	38,38,38,38	0
3	CL	E	1460	1/1	0.98	0.11	-	34,34,34,34	0
3	CL	G	1460	1/1	0.98	0.11	-	33,33,33,33	0
3	CL	I	1460	1/1	0.97	0.13	-	28,28,28,28	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.