



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 15, 2017 – 12:57 am GMT

PDB ID : 4YBQ
Title : Rat GLUT5 with Fv in the outward-open form
Authors : Nomura, N.; Shimamura, T.; Iwata, S.
Deposited on : 2015-02-19
Resolution : 3.27 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.9-1692
EDS	:	trunk28620
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.27 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	1006 (3.34-3.22)
Clashscore	112137	1070 (3.34-3.22)
Ramachandran outliers	110173	1050 (3.34-3.22)
Sidechain outliers	110143	1049 (3.34-3.22)
RSRZ outliers	101464	1011 (3.34-3.22)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	508	<div> <div>9%</div> <div> <div></div> <div>62%</div> <div>26%</div> <div>•</div> <div>11%</div> </div> </div>
1	B	508	<div> <div>9%</div> <div> <div></div> <div>65%</div> <div>25%</div> <div>•</div> <div>9%</div> </div> </div>
2	C	122	<div> <div>3%</div> <div> <div></div> <div>83%</div> <div>11%</div> <div>6%</div> </div> </div>
2	E	122	<div> <div>7%</div> <div> <div></div> <div>78%</div> <div>16%</div> <div>6%</div> </div> </div>
3	D	136	<div> <div>14%</div> <div> <div></div> <div>74%</div> <div>15%</div> <div>12%</div> </div> </div>
3	F	136	<div> <div>13%</div> <div> <div></div> <div>68%</div> <div>19%</div> <div>•</div> <div>13%</div> </div> </div>

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 10657 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Solute carrier family 2, facilitated glucose transporter member 5.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	452	Total	C	N	O	S	0	0	0
			3486	2313	553	601	19			
1	B	462	Total	C	N	O	S	0	0	0
			3567	2361	567	620	19			

There are 14 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	50	TYR	ASN	engineered mutation	UNP P43427
A	503	GLU	-	expression tag	UNP P43427
A	504	ASN	-	expression tag	UNP P43427
A	505	LEU	-	expression tag	UNP P43427
A	506	TYR	-	expression tag	UNP P43427
A	507	PHE	-	expression tag	UNP P43427
A	508	GLN	-	expression tag	UNP P43427
B	50	TYR	ASN	engineered mutation	UNP P43427
B	503	GLU	-	expression tag	UNP P43427
B	504	ASN	-	expression tag	UNP P43427
B	505	LEU	-	expression tag	UNP P43427
B	506	TYR	-	expression tag	UNP P43427
B	507	PHE	-	expression tag	UNP P43427
B	508	GLN	-	expression tag	UNP P43427

- Molecule 2 is a protein called antibody Fv fragment light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	115	Total	C	N	O	S	0	0	0
			885	562	147	174	2			
2	E	115	Total	C	N	O	S	0	0	0
			885	562	147	174	2			


- Molecule 3 is a protein called antibody Fv fragment heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	D	120	Total	C	N	O	S	0	0	0
			921	577	160	180	4			
3	F	119	Total	C	N	O	S	0	0	0
			913	571	159	179	4			


3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Solute carrier family 2, facilitated glucose transporter member 5





- Molecule 1: Solute carrier family 2, facilitated glucose transporter member 5








- Molecule 2: antibody Fv fragment light chain




- Molecule 2: antibody Fv fragment light chain



- Molecule 3: antibody Fv fragment heavy chain



- Molecule 3: antibody Fv fragment heavy chain



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	76.77Å 151.54Å 106.40Å 90.00° 97.25° 90.00°	Depositor
Resolution (Å)	36.49 – 3.27 38.08 – 3.24	Depositor EDS
% Data completeness (in resolution range)	99.8 (36.49-3.27) 90.2 (38.08-3.24)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.84 (at 3.25Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.9_1692)	Depositor
R, R_{free}	0.242 , 0.288 0.244 , 0.290	Depositor DCC
R_{free} test set	1842 reflections (4.94%)	DCC
Wilson B-factor (Å ²)	108.3	Xtriage
Anisotropy	0.347	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.26 , 78.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.89	EDS
Total number of atoms	10657	wwPDB-VP
Average B, all atoms (Å ²)	157.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.06% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.27	0/3564	0.48	0/4845
1	B	0.27	0/3647	0.47	0/4957
2	C	0.27	0/905	0.50	0/1226
2	E	0.29	0/905	0.53	0/1226
3	D	0.24	0/942	0.52	0/1275
3	F	0.26	0/934	0.56	0/1264
All	All	0.27	0/10897	0.49	0/14793

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3486	0	3646	91	0
1	B	3567	0	3715	88	0
2	C	885	0	871	10	0
2	E	885	0	871	12	0
3	D	921	0	880	14	0
3	F	913	0	866	20	0
All	All	10657	0	10849	216	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

The worst 5 of 216 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:364:LEU:HB3	1:A:371:MET:HG3	1.70	0.74
1:B:38:VAL:O	1:B:131:ARG:NH2	2.22	0.72
1:A:25:PHE:O	1:A:29:PHE:HB3	1.90	0.71
2:E:14:PRO:HB2	2:E:114:LYS:HG2	1.72	0.71
1:B:458:VAL:O	1:B:477:LYS:NZ	2.23	0.71

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	448/508 (88%)	436 (97%)	12 (3%)	0	100	100
1	B	458/508 (90%)	447 (98%)	11 (2%)	0	100	100
2	C	113/122 (93%)	110 (97%)	3 (3%)	0	100	100
2	E	113/122 (93%)	110 (97%)	3 (3%)	0	100	100
3	D	118/136 (87%)	117 (99%)	1 (1%)	0	100	100
3	F	117/136 (86%)	116 (99%)	1 (1%)	0	100	100
All	All	1367/1532 (89%)	1336 (98%)	31 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	378/433 (87%)	367 (97%)	11 (3%)	48	75
1	B	388/433 (90%)	379 (98%)	9 (2%)	56	79
2	C	101/108 (94%)	101 (100%)	0	100	100
2	E	101/108 (94%)	100 (99%)	1 (1%)	80	88
3	D	98/113 (87%)	96 (98%)	2 (2%)	60	81
3	F	97/113 (86%)	94 (97%)	3 (3%)	45	74
All	All	1163/1308 (89%)	1137 (98%)	26 (2%)	57	80

5 of 26 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	156	ASN
1	B	267	LEU
3	F	97	CYS
1	B	174	LEU
1	B	241	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
2	E	33	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry

There are no ligands in this entry.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	452/508 (88%)	0.40	46 (10%) 7 6	77, 135, 291, 452	0
1	B	462/508 (90%)	0.49	48 (10%) 7 6	78, 137, 297, 455	0
2	C	115/122 (94%)	0.20	4 (3%) 44 40	91, 127, 188, 232	0
2	E	115/122 (94%)	0.41	8 (6%) 17 16	100, 135, 193, 233	0
3	D	120/136 (88%)	0.64	19 (15%) 2 2	94, 156, 261, 285	0
3	F	119/136 (87%)	0.61	17 (14%) 3 3	94, 159, 264, 286	0
All	All	1383/1532 (90%)	0.45	142 (10%) 7 6	77, 139, 281, 455	0

The worst 5 of 142 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	188	GLU	15.5
1	B	190	GLY	14.0
1	B	259	ALA	13.7
1	B	258	ALA	12.7
1	B	260	GLY	11.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.