



# Full wwPDB X-ray Structure Validation Report ⓘ

Sep 18, 2017 – 09:33 AM EDT

PDB ID : 4YEC  
Title : Crystal structure of a clostripain (PARMER\_00083) from Parabacteroides merdae ATCC 43184 in complex with peptide inhibitor Ac-VLTK-AOMK  
Authors : Wolan, D.W.; Gonzalez-Paez, G.E.; Chatterjee, S.  
Deposited on : unknown  
Resolution : 1.12 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20029824  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20029824


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.12 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	1062 (1.16-1.08)
Clashscore	112137	1107 (1.16-1.08)
Ramachandran outliers	110173	1065 (1.16-1.08)
Sidechain outliers	110143	1062 (1.16-1.08)
RSRZ outliers	101464	1067 (1.16-1.08)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	126	<div> <div>6%</div> <div>91%</div> <div>9%</div> </div>
2	B	236	<div> <div>3%</div> <div>89%</div> <div>7%</div> <div>.</div> </div>
3	C	5	<div> <div>40%</div> <div>60%</div> <div>40%</div> </div>

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 3208 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Clostripain family.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	126	Total	C	N	O	S	0	7	0
			1008	638	166	198	6			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	22	MET	-	expression tag	UNP A7A9N3

- Molecule 2 is a protein called Clostripain family.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	228	Total	C	N	O	S	0	6	0
			1881	1215	287	366	13			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	376	LEU	-	expression tag	UNP A7A9N3
B	377	GLU	-	expression tag	UNP A7A9N3
B	378	HIS	-	expression tag	UNP A7A9N3
B	379	HIS	-	expression tag	UNP A7A9N3
B	380	HIS	-	expression tag	UNP A7A9N3
B	381	HIS	-	expression tag	UNP A7A9N3
B	382	HIS	-	expression tag	UNP A7A9N3
B	383	HIS	-	expression tag	UNP A7A9N3

- Molecule 3 is a protein called Peptide inhibitor Ac-VLTK-AOMK.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	C	5	Total	C	N	O	0	0	0
			35	24	5	6			


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	105	Total 105	O 105	0	0
4	B	176	Total 176	O 176	0	0
4	C	3	Total 3	O 3	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Clostripain family



- Molecule 2: Clostripain family



- Molecule 3: Peptide inhibitor Ac-VLTK-AOMK



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	38.89Å 107.34Å 40.94Å 90.00° 116.59° 90.00°	Depositor
Resolution (Å)	36.61 – 1.12 36.61 – 1.12	Depositor EDS
% Data completeness (in resolution range)	78.9 (36.61-1.12) 76.1 (36.61-1.12)	Depositor EDS
$R_{merge}$	0.11	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.69 (at 1.12Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.9_1692)	Depositor
R, $R_{free}$	0.158 , 0.191 0.158 , 0.190	Depositor DCC
$R_{free}$ test set	1923 reflections (2.20%)	DCC
Wilson B-factor (Å <sup>2</sup> )	14.7	Xtriage
Anisotropy	0.515	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.30 , 42.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.031 for h,-k,-h-l	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	3208	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	24.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.94% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CKC, ACE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.55	0/1050	0.64	0/1425
2	B	0.59	0/1947	0.71	0/2648
3	C	0.37	0/22	0.70	0/30
All	All	0.57	0/3019	0.68	0/4103

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	B	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	B	237	LEU	Peptide

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1008	0	1005	8	0
2	B	1881	0	1789	14	0
3	C	35	0	43	2	0
4	A	105	0	0	0	0
4	B	176	0	0	1	0
4	C	3	0	0	0	0
All	All	3208	0	2837	19	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (19) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:46:PHE:O	1:A:50[B]:ASN:ND2	2.06	0.88
2:B:238:ASN:N	2:B:238:ASN:HD22	1.73	0.86
2:B:238:ASN:H	2:B:238:ASN:HD22	1.24	0.85
2:B:365:GLN:HG3	2:B:366:ARG:HG2	1.66	0.76
1:A:39:ALA:O	1:A:44:ASN:ND2	2.24	0.70
2:B:238:ASN:H	2:B:238:ASN:ND2	1.91	0.68
1:A:135:THR:HG1	2:B:148:ALA:N	1.98	0.61
2:B:330:SER:HB2	2:B:341:ILE:HD11	1.91	0.52
2:B:177:ASP:OD1	3:C:405:CKC:NAM	2.42	0.52
2:B:209:TRP:HE3	2:B:214[B]:MET:SD	2.33	0.52
2:B:238:ASN:N	2:B:238:ASN:ND2	2.46	0.49
2:B:230:GLU:OE1	4:B:401:HOH:O	2.18	0.49
2:B:206:ALA:HB2	3:C:403:LEU:HB3	1.99	0.44
1:A:49[A]:LYS:HB2	1:A:49[A]:LYS:HE3	1.77	0.44
2:B:236:TYR:O	2:B:242:PRO:HA	2.18	0.43
1:A:47:ALA:HB2	1:A:131:TRP:CH2	2.54	0.42
1:A:130:LEU:HB3	2:B:149:PHE:CZ	2.54	0.42
1:A:131:TRP:O	1:A:132:SER:HB2	2.19	0.42
1:A:111:ARG:NH2	2:B:167:PRO:HA	2.37	0.40

There are no symmetry-related clashes.



## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	131/126 (104%)	125 (95%)	6 (5%)	0	100	100
2	B	232/236 (98%)	226 (97%)	5 (2%)	1 (0%)	38	11
3	C	2/5 (40%)	2 (100%)	0	0	100	100
All	All	365/367 (100%)	353 (97%)	11 (3%)	1 (0%)	44	15

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	238	ASN

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	115/110 (104%)	115 (100%)	0	100	100
2	B	203/208 (98%)	201 (99%)	2 (1%)	80	45
3	C	3/3 (100%)	3 (100%)	0	100	100
All	All	321/321 (100%)	319 (99%)	2 (1%)	89	63

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	183	SER
2	B	238	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
2	B	238	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	126/126 (100%)	0.36	8 (6%) 21 21	13, 21, 36, 55	0
2	B	228/236 (96%)	0.27	6 (2%) 56 53	12, 21, 35, 47	0
3	C	3/5 (60%)	2.65	2 (66%) 0 0	34, 34, 39, 45	0
All	All	357/367 (97%)	0.32	16 (4%) 34 32	12, 21, 37, 55	0

All (16) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	238	ASN	9.0
2	B	288	TYR	5.0
1	A	22	MET	4.0
1	A	23	GLU	4.0
3	C	402	VAL	3.9
2	B	154	ASN	3.7
1	A	44	ASN	2.8
1	A	147	LYS	2.7
2	B	207	ASP	2.4
1	A	24	THR	2.3
1	A	45	SER	2.3
1	A	42	SER	2.2
3	C	403	LEU	2.2
1	A	76	GLY	2.1
2	B	308	GLU	2.1
2	B	306	THR	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

### 6.5 Other polymers [i](#)

There are no such residues in this entry.