



# Full wwPDB X-ray Structure Validation Report ⓘ

Oct 16, 2017 – 11:20 AM EDT

PDB ID : 5YFM  
Title : Human isocitrate dehydrogenase 1 bound with NADP  
Authors : Nordlund, P.; Chen, D.; Jansson, A.; Larsson, A.  
Deposited on : unknown  
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20030345  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20030345


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	3166 (2.40-2.40)
Clashscore	112137	3674 (2.40-2.40)
Ramachandran outliers	110173	3616 (2.40-2.40)
Sidechain outliers	110143	3617 (2.40-2.40)
RSRZ outliers	101464	3195 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	414	
1	B	414	
1	C	414	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	MG	A	502	-	-	-	X

2 Entry composition ⓘ


There are 4 unique types of molecules in this entry. The entry contains 10322 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Isocitrate dehydrogenase [NADP] cytoplasmic.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	409	Total	C	N	O	S	0	1	0
			3244	2062	549	615	18			
1	B	409	Total	C	N	O	S	0	1	0
			3248	2065	550	615	18			
1	C	408	Total	C	N	O	S	0	0	0
			3223	2049	547	609	18			

- Molecule 2 is NADP NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (three-letter code: NAP) (formula: C<sub>21</sub>H<sub>28</sub>N<sub>7</sub>O<sub>17</sub>P<sub>3</sub>).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			48	21	7	17	3		
2	B	1	Total	C	N	O	P	0	0
			48	21	7	17	3		
2	C	1	Total	C	N	O	P	0	0
			48	21	7	17	3		

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total 1	Mg 1	0	0
3	A	1	Total 1	Mg 1	0	0
3	C	1	Total 1	Mg 1	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	194	Total 194	O 194	0	0
4	B	155	Total 155	O 155	0	0
4	C	111	Total 111	O 111	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Isocitrate dehydrogenase [NADP] cytoplasmic



- Molecule 1: Isocitrate dehydrogenase [NADP] cytoplasmic



- Molecule 1: Isocitrate dehydrogenase [NADP] cytoplasmic



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	203.02Å 85.58Å 86.07Å 90.00° 98.08° 90.00°	Depositor
Resolution (Å)	100.50 – 2.40 41.37 – 2.40	Depositor EDS
% Data completeness (in resolution range)	99.0 (100.50-2.40) 99.0 (41.37-2.40)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.11	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.73 (at 2.39Å)	Xtriage
Refinement program	REFMAC 5.8.0158	Depositor
R, $R_{free}$	0.173 , 0.239 0.181 , 0.241	Depositor DCC
$R_{free}$ test set	2875 reflections (5.34%)	DCC
Wilson B-factor (Å <sup>2</sup> )	23.7	Xtriage
Anisotropy	0.471	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 38.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	10322	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	30.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.65% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAP, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	1.01	4/3312 (0.1%)	0.90	3/4469 (0.1%)
1	B	0.95	0/3316	0.91	0/4473
1	C	0.86	1/3291 (0.0%)	0.84	1/4441 (0.0%)
All	All	0.94	5/9919 (0.1%)	0.88	4/13383 (0.0%)

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	247	GLU	CD-OE2	-5.52	1.19	1.25
1	C	274	GLY	C-O	-5.24	1.15	1.23
1	A	391	TYR	CE1-CZ	-5.08	1.31	1.38
1	A	332	SER	CB-OG	-5.07	1.35	1.42
1	A	326	SER	CB-OG	-5.02	1.35	1.42

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	132	ARG	NE-CZ-NH1	5.81	123.20	120.30
1	A	317	ARG	NE-CZ-NH1	5.70	123.15	120.30
1	A	249	ARG	NE-CZ-NH1	5.38	122.99	120.30
1	A	20	ARG	NE-CZ-NH1	5.08	122.84	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen



atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3244	0	3198	18	0
1	B	3248	0	3209	20	0
1	C	3223	0	3178	15	0
2	A	48	0	25	0	0
2	B	48	0	25	0	0
2	C	48	0	25	2	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
3	C	1	0	0	0	0
4	A	194	0	0	2	2
4	B	155	0	0	0	0
4	C	111	0	0	0	1
All	All	10322	0	9660	47	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (47) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:211:THR:HG23	1:C:220:ASP:HB3	1.56	0.88
1:A:160:ASP:OD1	1:A:162:THR:HG22	1.96	0.66
1:A:100:ARG:NH2	4:A:601:HOH:O	2.34	0.60
1:C:314:ARG:NH2	2:C:501:NAP:O3X	2.35	0.60
1:A:239:PHE:CD2	1:A:244[A]:ILE:HG22	2.36	0.59
1:A:219:TYR:CD2	1:B:178:VAL:HG13	2.38	0.59
1:C:13:MET:HB3	1:C:44:LEU:HD12	1.85	0.58
1:A:18:MET:HB2	1:A:316:TYR:HB2	1.85	0.57
1:A:15:GLY:O	1:A:20:ARG:HD2	2.07	0.55
1:B:142:THR:HG23	1:B:142:THR:O	2.07	0.55
1:A:139:TYR:CE2	1:B:215:ILE:HD13	2.44	0.53
1:B:362:VAL:HG23	1:B:408:LYS:HD2	1.92	0.52
1:C:211:THR:CG2	1:C:220:ASP:HB3	2.35	0.52
1:C:68:ASN:HA	1:C:302:THR:HG23	1.92	0.51
1:C:75:THR:O	2:C:501:NAP:H2N	2.10	0.51
1:A:362:VAL:HG23	1:A:408:LYS:HD2	1.93	0.50
1:A:215:ILE:HD13	1:B:139:TYR:CE2	2.47	0.50
1:A:25:LEU:HD11	1:A:395:PHE:CE1	2.46	0.50
1:B:121:VAL:HG22	1:B:259:MET:HE1	1.94	0.50

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:191:ASP:HB3	1:B:298:PRO:HG3	1.95	0.48
1:C:207:LEU:HD13	1:C:246:TYR:CD1	2.49	0.48
1:C:133:HIS:HB2	1:C:192:PHE:CE2	2.48	0.48
1:A:178:VAL:HG13	1:B:219:TYR:CD2	2.50	0.47
1:B:371:PHE:CZ	1:B:400:LYS:HE3	2.50	0.47
1:B:408:LYS:HG3	1:B:408:LYS:O	2.13	0.47
1:B:362:VAL:HG21	1:B:405:LEU:HA	1.96	0.46
1:A:361:GLU:OE2	1:A:408:LYS:NZ	2.49	0.46
1:C:314:ARG:HB3	1:C:314:ARG:NH2	2.31	0.45
1:A:117:ILE:HD12	1:A:376:LEU:HB3	1.98	0.45
1:C:112:ILE:HG12	1:C:334:PHE:CE2	2.51	0.45
1:A:219:TYR:CE2	1:B:178:VAL:HG13	2.52	0.44
1:B:18:MET:HB2	1:B:316:TYR:HB2	1.99	0.44
1:C:13:MET:HB3	1:C:44:LEU:CD1	2.46	0.44
1:A:334:PHE:HA	1:A:337:THR:OG1	2.17	0.44
1:B:392:LEU:HB3	1:B:396:GLU:HB2	1.99	0.44
1:C:324:GLU:OE2	1:C:388:ARG:NH1	2.51	0.44
1:C:334:PHE:HA	1:C:337:THR:OG1	2.17	0.43
1:A:155:THR:HG22	4:A:666:HOH:O	2.17	0.43
1:B:212:LYS:HE3	1:B:215:ILE:HD12	2.01	0.43
1:B:397:PHE:CE2	1:B:401:LEU:HD11	2.55	0.42
1:B:336:TRP:O	1:B:340:LEU:HG	2.20	0.41
1:A:178:VAL:CG2	1:B:219:TYR:HA	2.50	0.41
1:C:211:THR:CG2	1:C:213:ASN:HB3	2.51	0.41
1:A:25:LEU:HD11	1:A:395:PHE:HE1	1.85	0.41
1:B:400:LYS:HD3	1:B:400:LYS:HA	1.89	0.40
1:C:126:LYS:HB3	1:C:127:PRO:HD2	2.03	0.40
1:B:108:PHE:HA	1:B:130:ILE:O	2.21	0.40

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:C:687:HOH:O	4:C:704:HOH:O[2_456]	1.56	0.64
4:A:652:HOH:O	4:A:772:HOH:O[4_455]	1.81	0.39
4:A:765:HOH:O	4:A:785:HOH:O[4_455]	1.93	0.27

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	408/414 (99%)	395 (97%)	12 (3%)	1 (0%)	51	67
1	B	408/414 (99%)	387 (95%)	21 (5%)	0	100	100
1	C	406/414 (98%)	393 (97%)	13 (3%)	0	100	100
All	All	1222/1242 (98%)	1175 (96%)	46 (4%)	1 (0%)	55	72

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	122	SER

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	345/350 (99%)	321 (93%)	24 (7%)	18	28
1	B	346/350 (99%)	328 (95%)	18 (5%)	27	43
1	C	342/350 (98%)	313 (92%)	29 (8%)	12	19
All	All	1033/1050 (98%)	962 (93%)	71 (7%)	18	28

All (71) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	39	LEU
1	A	65	LYS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	82	ARG
1	A	89	LYS
1	A	115	LYS
1	A	119	ARG
1	A	122	SER
1	A	125	VAL
1	A	155	THR
1	A	162	THR
1	A	168	LEU
1	A	169	VAL
1	A	174	GLU
1	A	178	VAL
1	A	184	ASN
1	A	207	LEU
1	A	216	LEU
1	A	222	ARG
1	A	249	ARG
1	A	381	LYS
1	A	383	LEU
1	A	403	GLU
1	A	409	LEU
1	A	411	GLN
1	B	4	LYS
1	B	39	LEU
1	B	82	ARG
1	B	115	LYS
1	B	119	ARG
1	B	125	VAL
1	B	169	VAL
1	B	174	GLU
1	B	178	VAL
1	B	184	ASN
1	B	207	LEU
1	B	216	LEU
1	B	222	ARG
1	B	240	GLU
1	B	249	ARG
1	B	403	GLU
1	B	409	LEU
1	B	411	GLN
1	C	12	GLU
1	C	19	THR

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	C	22	ILE
1	C	58	LYS
1	C	62	GLU
1	C	82	ARG
1	C	89	LYS
1	C	103	LEU
1	C	109	ARG
1	C	168	LEU
1	C	180	MET
1	C	184	ASN
1	C	198	GLN
1	C	203	LYS
1	C	207	LEU
1	C	211	THR
1	C	216	LEU
1	C	260	LYS
1	C	280	SER
1	C	288	LEU
1	C	304	GLU
1	C	312	VAL
1	C	313	THR
1	C	314	ARG
1	C	338	ARG
1	C	383	LEU
1	C	388	ARG
1	C	393	ASN
1	C	409	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	323	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 3 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	NAP	A	501	-	44,52,52	1.97	14 (31%)	51,80,80	1.65	8 (15%)
2	NAP	B	501	-	44,52,52	1.88	15 (34%)	51,80,80	1.77	12 (23%)
2	NAP	C	501	-	44,52,52	1.33	5 (11%)	51,80,80	1.42	5 (9%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAP	A	501	-	-	0/27/67/67	0/5/5/5
2	NAP	B	501	-	-	0/27/67/67	0/5/5/5
2	NAP	C	501	-	-	0/27/67/67	0/5/5/5

All (34) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	501	NAP	P2B-O2B	-4.28	1.52	1.59
2	A	501	NAP	P2B-O2B	-4.26	1.52	1.59
2	B	501	NAP	P2B-O3X	-4.01	1.38	1.54
2	A	501	NAP	P2B-O3X	-3.87	1.38	1.54
2	A	501	NAP	P2B-O2X	-3.82	1.39	1.54
2	B	501	NAP	O7N-C7N	-3.36	1.17	1.24
2	A	501	NAP	P2B-O1X	-3.17	1.39	1.50
2	A	501	NAP	PN-O2N	-3.03	1.39	1.55
2	B	501	NAP	PN-O2N	-3.01	1.40	1.55

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	501	NAP	PA-O2A	-2.90	1.40	1.55
2	A	501	NAP	O7N-C7N	-2.89	1.18	1.24
2	A	501	NAP	C4A-N3A	-2.82	1.31	1.35
2	B	501	NAP	C5A-N7A	-2.76	1.30	1.39
2	B	501	NAP	P2B-O1X	-2.75	1.41	1.50
2	B	501	NAP	PA-O2A	-2.73	1.41	1.55
2	B	501	NAP	P2B-O2X	-2.71	1.43	1.54
2	A	501	NAP	PN-O1N	-2.70	1.40	1.50
2	A	501	NAP	PA-O1A	-2.69	1.40	1.50
2	B	501	NAP	PN-O1N	-2.65	1.41	1.50
2	A	501	NAP	C5A-N7A	-2.63	1.30	1.39
2	B	501	NAP	PA-O1A	-2.60	1.41	1.50
2	C	501	NAP	C2D-C1D	-2.56	1.49	1.53
2	B	501	NAP	C8A-N7A	-2.35	1.30	1.34
2	A	501	NAP	O4D-C4D	-2.32	1.39	1.45
2	A	501	NAP	O4B-C4B	-2.27	1.39	1.45
2	A	501	NAP	C8A-N7A	-2.27	1.30	1.34
2	B	501	NAP	C2D-C1D	-2.25	1.50	1.53
2	B	501	NAP	O4D-C4D	-2.18	1.40	1.45
2	C	501	NAP	O7N-C7N	-2.10	1.19	1.24
2	C	501	NAP	PN-O2N	-2.03	1.44	1.55
2	C	501	NAP	C5A-N7A	-2.01	1.32	1.39
2	B	501	NAP	PN-O5D	-2.01	1.50	1.59
2	B	501	NAP	C5A-C4A	2.08	1.45	1.40
2	C	501	NAP	C5A-C4A	2.55	1.46	1.40

All (25) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	501	NAP	N3A-C2A-N1A	-6.65	123.06	128.86
2	C	501	NAP	N3A-C2A-N1A	-6.06	123.58	128.86
2	B	501	NAP	N3A-C2A-N1A	-5.42	124.14	128.86
2	B	501	NAP	C4A-C5A-N7A	-3.84	105.70	109.41
2	A	501	NAP	C4A-C5A-N7A	-3.62	105.91	109.41
2	B	501	NAP	C5N-C4N-C3N	-3.58	116.14	120.35
2	A	501	NAP	C5N-C4N-C3N	-3.27	116.50	120.35
2	C	501	NAP	C4A-C5A-N7A	-3.06	106.46	109.41
2	C	501	NAP	O2B-P2B-O1X	-2.94	97.72	109.26
2	B	501	NAP	O3D-C3D-C4D	-2.94	102.49	111.09
2	B	501	NAP	O7N-C7N-N7N	-2.49	119.04	122.58
2	A	501	NAP	O7N-C7N-C3N	2.04	122.01	119.62
2	C	501	NAP	C4D-O4D-C1D	2.04	111.94	109.77

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
2	B	501	NAP	O2X-P2B-O1X	2.13	118.82	110.50
2	A	501	NAP	O3X-P2B-O2X	2.15	116.28	107.61
2	B	501	NAP	O2B-C2B-C3B	2.27	120.05	111.63
2	B	501	NAP	O3B-C3B-C4B	2.33	117.88	111.09
2	B	501	NAP	C2A-N1A-C6A	2.36	122.90	118.77
2	C	501	NAP	C2A-N1A-C6A	2.39	122.95	118.77
2	A	501	NAP	C3N-C2N-N1N	2.43	122.87	120.43
2	B	501	NAP	C3N-C7N-N7N	2.82	121.00	117.77
2	A	501	NAP	C4D-O4D-C1D	2.85	112.80	109.77
2	A	501	NAP	C2A-N1A-C6A	3.08	124.16	118.77
2	B	501	NAP	C4D-O4D-C1D	3.28	113.26	109.77
2	B	501	NAP	C3N-C2N-N1N	3.58	124.03	120.43

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	501	NAP	2	0

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.



## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	409/414 (98%)	-0.57	0 <span>100</span> <span>100</span>	11, 22, 45, 64	0
1	B	409/414 (98%)	-0.58	2 (0%) <span>90</span> <span>89</span>	12, 21, 51, 69	0
1	C	408/414 (98%)	-0.11	16 (3%) <span>40</span> <span>39</span>	13, 40, 73, 99	0
All	All	1226/1242 (98%)	-0.42	18 (1%) <span>74</span> <span>72</span>	11, 24, 62, 99	0

All (18) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	88	LEU	4.2
1	C	322	GLY	3.6
1	C	90	GLN	3.6
1	C	83	VAL	3.5
1	B	161	GLY	3.3
1	C	389	SER	3.2
1	C	84	GLU	3.0
1	C	87	LYS	3.0
1	C	6	SER	2.8
1	C	407	ILE	2.6
1	C	321	LYS	2.6
1	C	92	TRP	2.5
1	C	409	LEU	2.5
1	C	317	ARG	2.3
1	C	320	GLN	2.3
1	C	91	MET	2.2
1	C	354	PHE	2.0
1	B	88	LEU	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
3	MG	A	502	1/1	0.91	0.19	2.83	24,24,24,24	0
2	NAP	C	501	48/48	0.93	0.14	-0.24	47,54,65,69	0
2	NAP	A	501	48/48	0.97	0.09	-0.96	14,20,32,35	0
2	NAP	B	501	48/48	0.98	0.08	-1.29	14,18,25,29	0
3	MG	C	502	1/1	0.92	0.24	-	32,32,32,32	0
3	MG	B	502	1/1	0.84	0.27	-	33,33,33,33	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.