



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 14, 2017 – 05:52 pm GMT

PDB ID : 2YMU  
Title : Structure of a highly repetitive propeller structure  
Authors : Zeth, K.; Chaudhuri, I.; Martin, J.; Lupas, A.N.  
Deposited on : 2012-10-10  
Resolution : 1.79 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.9-1692  
EDS : trunk28620  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.79 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	4827 (1.80-1.80)
Clashscore	112137	5742 (1.80-1.80)
Ramachandran outliers	110173	5676 (1.80-1.80)
Sidechain outliers	110143	5675 (1.80-1.80)
RSRZ outliers	101464	4906 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	577	<div> <div>3%</div> <div> <div></div> <div>93%</div> <div>6% •</div> </div> </div>
1	B	577	<div> <div>4%</div> <div> <div></div> <div>91%</div> <div>8%</div> </div> </div>

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 9719 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called WD-40 REPEAT PROTEIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	577	Total	C	N	O	S	0	24	0
			4419	2722	801	895	1			
1	B	577	Total	C	N	O	S	0	22	0
			4413	2718	801	893	1			

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	GLY	-	EXPRESSION TAG	UNP B2J0I0
A	2	SER	-	EXPRESSION TAG	UNP B2J0I0
A	3	HIS	-	EXPRESSION TAG	UNP B2J0I0
A	4	MET	-	EXPRESSION TAG	UNP B2J0I0
A	336	HIS	LEU	CONFLICT	UNP B2J0I0
B	1	GLY	-	EXPRESSION TAG	UNP B2J0I0
B	2	SER	-	EXPRESSION TAG	UNP B2J0I0
B	3	HIS	-	EXPRESSION TAG	UNP B2J0I0
B	4	MET	-	EXPRESSION TAG	UNP B2J0I0
B	336	HIS	LEU	CONFLICT	UNP B2J0I0


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	469	Total	O	0	0
			469	469		
2	B	418	Total	O	0	0
			418	418		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: WD-40 REPEAT PROTEIN



#### • Molecule 1: WD-40 REPEAT PROTEIN



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	88.47Å 107.73Å 136.26Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.90 – 1.79 29.90 – 1.79	Depositor EDS
% Data completeness (in resolution range)	99.5 (29.90-1.79) 99.5 (29.90-1.79)	Depositor EDS
$R_{merge}$	0.13	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.54 (at 1.79Å)	Xtriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
R, $R_{free}$	0.174 , 0.207 0.174 , 0.207	Depositor DCC
$R_{free}$ test set	6077 reflections (5.00%)	DCC
Wilson B-factor (Å <sup>2</sup> )	19.3	Xtriage
Anisotropy	0.529	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.40 , 49.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	9719	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	21.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.80% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.52	0/4576	0.71	3/6224 (0.0%)
1	B	0.51	0/4564	0.72	1/6207 (0.0%)
All	All	0.51	0/9140	0.72	4/12431 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	1
All	All	0	2

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	128	ASN	N-CA-C	5.76	126.54	111.00
1	B	128	ASN	N-CA-C	5.74	126.49	111.00
1	A	83	LEU	CA-CB-CG	5.27	127.41	115.30
1	A	324	ASP	CB-CG-OD1	5.17	122.95	118.30

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	137	GLY	Peptide
1	B	137	GLY	Peptide

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4419	0	4354	25	0
1	B	4413	0	4344	35	0
2	A	469	0	0	7	5
2	B	418	0	0	9	3
All	All	9719	0	8698	58	5

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (58) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:526:ALA:HB2	1:B:555[A]:VAL:HG23	1.44	0.96
1:A:526:ALA:HB2	1:A:555[A]:VAL:HG23	1.52	0.91
1:A:439:GLN:OE1	2:A:2388:HOH:O	1.94	0.85
1:B:290:ASN:HD21	1:B:294:GLN:HE21	1.26	0.81
1:B:22:VAL:HG13	1:B:280:ALA:HB2	1.70	0.74
1:A:136:THR:O	2:A:2164:HOH:O	2.06	0.74
1:A:417:GLN:OE1	2:A:2370:HOH:O	2.06	0.72
1:A:128:ASN:ND2	2:A:2151:HOH:O	2.22	0.71
1:A:435[A]:SER:OG	1:A:437:ASP:OD1	2.11	0.68
1:B:323:ASP:HA	1:B:346[B]:SER:HB3	1.76	0.67
1:B:127:ARG:NH2	2:B:2094:HOH:O	2.20	0.67
1:B:323:ASP:HA	1:B:346[A]:SER:HB3	1.77	0.66
1:B:436:PRO:O	2:B:2320:HOH:O	2.13	0.65
1:B:526:ALA:HB2	1:B:555[B]:VAL:HG13	1.77	0.65
1:B:133:GLN:OE1	2:B:2127:HOH:O	2.16	0.63
1:B:127:ARG:NE	2:B:2094:HOH:O	2.21	0.62
1:B:526:ALA:HB2	1:B:555[A]:VAL:CG2	2.25	0.60
1:A:1:GLY:N	1:B:95:THR:O	2.34	0.59
1:B:434:PHE:HE1	1:B:455:ARG:HE	1.50	0.59
1:B:45:ARG:HG2	2:B:2053:HOH:O	2.02	0.59
1:B:439:GLN:CD	2:B:2319:HOH:O	2.42	0.57
1:A:53:LEU:HD22	1:A:88:GLY:HA2	1.88	0.55
1:B:444:ALA:HB1	1:B:470:VAL:HG12	1.90	0.54

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:110:GLY:O	1:A:127:ARG:NH1	2.40	0.53
1:A:9:ARG:HG3	1:A:577:ASN:HB3	1.90	0.53
1:B:434:PHE:HE1	1:B:455:ARG:NE	2.07	0.53
1:B:176:LEU:CD2	1:B:211:GLY:HA2	2.40	0.52
1:B:53:LEU:HD22	1:B:88:GLY:HA2	1.92	0.52
1:B:454[A]:ASN:HD21	1:B:458:GLN:HB2	1.74	0.51
1:A:30:THR:OG1	1:A:86:ARG:NH2	2.38	0.51
1:B:30:THR:OG1	1:B:86:ARG:NH2	2.32	0.51
1:B:110:GLY:O	1:B:127:ARG:NH1	2.36	0.50
1:B:422:LEU:CD2	1:B:457:GLY:HA2	2.41	0.50
1:B:176:LEU:HD22	1:B:211:GLY:HA2	1.93	0.49
1:B:530:LYS:HB2	1:B:530:LYS:HE2	1.63	0.49
1:A:340:LEU:CD2	1:A:375:GLY:HA2	2.43	0.48
1:B:202:LYS:HE3	1:B:222:SER:HA	1.95	0.48
1:B:290:ASN:OD1	1:B:294:GLN:HG2	2.14	0.47
1:A:526:ALA:HB2	1:A:555[A]:VAL:CG2	2.35	0.47
1:B:454[A]:ASN:ND2	1:B:458:GLN:HB2	2.30	0.46
1:A:374:ASN:OD1	2:A:2335:HOH:O	2.21	0.46
1:A:202:LYS:HE3	1:A:222:SER:HA	1.98	0.46
1:A:496:ARG:HD2	2:A:2426:HOH:O	2.16	0.45
1:B:297:GLN:OE1	2:B:2228:HOH:O	2.21	0.44
1:B:437:ASP:OD1	2:B:2318:HOH:O	2.21	0.43
1:A:118:ASP:HA	1:A:141:SER:HB2	1.99	0.43
1:A:2:SER:OG	1:B:92:GLN:OE1	2.36	0.43
1:A:126:ASN:O	1:A:129:GLY:HA2	2.19	0.43
1:A:128:ASN:N	1:A:129:GLY:HA2	2.34	0.43
1:A:290:ASN:HD21	1:A:294:GLN:HG3	1.84	0.42
1:B:340:LEU:HD13	1:B:371:TRP:CG	2.54	0.42
1:A:441:ILE:HD11	1:A:455:ARG:HG2	2.01	0.42
1:B:128:ASN:N	1:B:129:GLY:HA2	2.35	0.41
1:B:441:ILE:HD11	1:B:455:ARG:HD3	2.03	0.41
1:B:475:PHE:O	2:B:2348:HOH:O	2.22	0.41
1:A:11:ARG:HD3	2:A:2024:HOH:O	2.20	0.40
1:A:113[B]:ILE:HD11	1:A:127:ARG:HD2	2.03	0.40
1:A:526:ALA:HB2	1:A:555[B]:VAL:HG13	2.02	0.40

All (5) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:2147:HOH:O	2:A:2357:HOH:O[4_545]	1.91	0.29

Continued on next page...



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:2065:HOH:O	2:A:2366:HOH:O[4_545]	1.97	0.23
2:A:2269:HOH:O	2:B:2342:HOH:O[4_445]	2.00	0.20
2:A:2284:HOH:O	2:B:2048:HOH:O[3_554]	2.12	0.08
2:A:2209:HOH:O	2:B:2186:HOH:O[1_565]	2.14	0.06

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	599/577 (104%)	571 (95%)	28 (5%)	0	100	100
1	B	597/577 (104%)	567 (95%)	30 (5%)	0	100	100
All	All	1196/1154 (104%)	1138 (95%)	58 (5%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	503/479 (105%)	495 (98%)	8 (2%)	68	58
1	B	501/479 (105%)	489 (98%)	12 (2%)	54	40
All	All	1004/958 (105%)	984 (98%)	20 (2%)	66	48

All (20) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	22	VAL
1	A	83	LEU
1	A	113[A]	ILE
1	A	113[B]	ILE
1	A	176	LEU
1	A	294	GLN
1	A	311	PHE
1	A	332	ARG
1	B	42	LEU
1	B	99[A]	SER
1	B	99[B]	SER
1	B	113[A]	ILE
1	B	113[B]	ILE
1	B	288	LEU
1	B	292	ASN
1	B	311	PHE
1	B	332	ARG
1	B	346[A]	SER
1	B	346[B]	SER
1	B	419	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	294	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry

There are no ligands in this entry.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å <sup>2</sup> )	Q<0.9
1	A	577/577 (100%)	0.06	18 (3%)	49	44	9, 17, 33, 52	1 (0%)
1	B	577/577 (100%)	0.12	25 (4%)	36	30	11, 19, 36, 67	1 (0%)
All	All	1154/1154 (100%)	0.09	43 (3%)	42	37	9, 18, 34, 67	2 (0%)

All (43) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	438	ASP	5.6
1	B	437	ASP	4.7
1	B	439	GLN	4.2
1	B	455	ARG	4.1
1	A	555[A]	VAL	3.8
1	A	128	ASN	3.7
1	B	391	VAL	3.6
1	A	292	ASN	3.3
1	B	555[A]	VAL	3.2
1	A	514	VAL	3.1
1	A	473	VAL	3.1
1	B	128	ASN	3.0
1	B	22	VAL	3.0
1	B	45	ARG	2.9
1	B	292	ASN	2.9
1	A	432	VAL	2.8
1	B	458	GLN	2.8
1	A	22	VAL	2.7
1	B	415	ASN	2.6
1	B	432	VAL	2.6
1	A	130[A]	GLN	2.6
1	B	63	VAL	2.5
1	B	473	VAL	2.4
1	A	499	GLN	2.4

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	139	SER	2.4
1	B	145	VAL	2.3
1	B	227	VAL	2.3
1	A	316	GLN	2.3
1	B	310	ALA	2.3
1	A	309	VAL	2.2
1	A	104	VAL	2.2
1	A	169	ASN	2.2
1	A	439	GLN	2.2
1	B	456	ASN	2.2
1	B	186	VAL	2.2
1	B	350	VAL	2.2
1	B	417	GLN	2.2
1	A	350	VAL	2.1
1	A	431	GLY	2.1
1	B	436	PRO	2.1
1	A	145	VAL	2.1
1	A	497	ASN	2.1
1	B	169	ASN	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.