



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 13, 2017 – 04:13 am GMT

PDB ID : 4YNL
Title : Crystal structure of the hood domain of Anabaena HetR in complex with the hexapeptide ERGSGR derived from PatS
Authors : Hu, H.X.; Jiang, Y.L.; Zhao, M.X.; Zhang, C.C.; Chen, Y.; Zhou, C.Z.
Deposited on : 2015-03-10
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.9-1692
EDS	:	trunk28620
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	4243 (2.10-2.10)
Clashscore	112137	4788 (2.10-2.10)
Ramachandran outliers	110173	4740 (2.10-2.10)
Sidechain outliers	110143	4741 (2.10-2.10)
RSRZ outliers	101464	4275 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	90	<div> <div>67%</div> <div>19%</div> <div>14%</div> </div>
1	B	90	<div> <div>%</div> <div>60%</div> <div>24%</div> <div>16%</div> </div>
1	M	90	<div> <div>54%</div> <div>78%</div> <div>8%</div> <div>14%</div> </div>
1	N	90	<div> <div>44%</div> <div>71%</div> <div>13%</div> <div>16%</div> </div>
2	C	6	<div> <div>83%</div> <div>17%</div> </div>
2	D	6	<div> <div>67%</div> <div>33%</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	P	6	<div><div></div><div>50%</div><div>83%</div><div>17%</div></div>
2	R	6	<div><div></div><div>50%</div><div>83%</div><div>17%</div></div>

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 2958 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Heterocyst differentiation control protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	B	76	Total	C	N	O	S	0	1	0
			649	413	113	116	7			
1	A	77	Total	C	N	O	S	0	1	0
			658	418	114	119	7			
1	N	76	Total	C	N	O	S	0	0	0
			645	409	113	116	7			
1	M	77	Total	C	N	O	S	0	0	0
			654	414	114	119	7			

There are 36 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	210	MET	-	expression tag	UNP P27709
B	211	GLY	-	expression tag	UNP P27709
B	212	HIS	-	expression tag	UNP P27709
B	213	HIS	-	expression tag	UNP P27709
B	214	HIS	-	expression tag	UNP P27709
B	215	HIS	-	expression tag	UNP P27709
B	216	HIS	-	expression tag	UNP P27709
B	217	HIS	-	expression tag	UNP P27709
B	218	MET	-	expression tag	UNP P27709
A	210	MET	-	expression tag	UNP P27709
A	211	GLY	-	expression tag	UNP P27709
A	212	HIS	-	expression tag	UNP P27709
A	213	HIS	-	expression tag	UNP P27709
A	214	HIS	-	expression tag	UNP P27709
A	215	HIS	-	expression tag	UNP P27709
A	216	HIS	-	expression tag	UNP P27709
A	217	HIS	-	expression tag	UNP P27709
A	218	MET	-	expression tag	UNP P27709
N	210	MET	-	expression tag	UNP P27709
N	211	GLY	-	expression tag	UNP P27709
N	212	HIS	-	expression tag	UNP P27709

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
N	213	HIS	-	expression tag	UNP P27709
N	214	HIS	-	expression tag	UNP P27709
N	215	HIS	-	expression tag	UNP P27709
N	216	HIS	-	expression tag	UNP P27709
N	217	HIS	-	expression tag	UNP P27709
N	218	MET	-	expression tag	UNP P27709
M	210	MET	-	expression tag	UNP P27709
M	211	GLY	-	expression tag	UNP P27709
M	212	HIS	-	expression tag	UNP P27709
M	213	HIS	-	expression tag	UNP P27709
M	214	HIS	-	expression tag	UNP P27709
M	215	HIS	-	expression tag	UNP P27709
M	216	HIS	-	expression tag	UNP P27709
M	217	HIS	-	expression tag	UNP P27709
M	218	MET	-	expression tag	UNP P27709

- Molecule 2 is a protein called Heterocyst inhibition-signaling peptide.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	D	6	Total	C	N	O	0	0	0
			45	24	12	9			
2	C	6	Total	C	N	O	0	0	0
			45	24	12	9			
2	P	6	Total	C	N	O	0	0	0
			45	24	12	9			
2	R	6	Total	C	N	O	0	0	0
			45	24	12	9			

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	68	Total	O	0	0
			68	68		
3	A	82	Total	O	0	0
			82	82		
3	D	10	Total	O	0	0
			10	10		
3	C	11	Total	O	0	0
			11	11		
3	M	1	Total	O	0	0
			1	1		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: Heterocyst differentiation control protein





- Molecule 1: Heterocyst differentiation control protein



- Molecule 1: Heterocyst differentiation control protein



- Molecule 1: Heterocyst differentiation control protein




- Molecule 2: Heterocyst inhibition-signaling peptide






- Molecule 2: Heterocyst inhibition-signaling peptide

Chain C: 




- Molecule 2: Heterocyst inhibition-signaling peptide

Chain P: 



- Molecule 2: Heterocyst inhibition-signaling peptide

Chain R: 



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	218.22Å 43.46Å 55.11Å 90.00° 97.54° 90.00°	Depositor
Resolution (Å)	42.65 – 2.10 42.61 – 2.09	Depositor EDS
% Data completeness (in resolution range)	96.3 (42.65-2.10) 96.4 (42.61-2.09)	Depositor EDS
R_{merge}	0.14	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.34 (at 2.10Å)	Xtriage
Refinement program	REFMAC 5.7.0032	Depositor
R, R_{free}	0.212 , 0.256 0.213 , 0.252	Depositor DCC
R_{free} test set	1504 reflections (5.37%)	DCC
Wilson B-factor (Å ²)	26.2	Xtriage
Anisotropy	0.773	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 71.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	2958	wwPDB-VP
Average B, all atoms (Å ²)	72.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.40% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.81	0/676	0.87	0/911
1	B	0.79	0/667	0.94	2/899 (0.2%)
1	M	0.31	0/669	0.52	0/900
1	N	0.31	0/660	0.52	0/888
2	C	1.14	0/44	1.43	2/55 (3.6%)
2	D	1.03	0/44	1.35	1/55 (1.8%)
2	P	0.38	0/44	0.63	0/55
2	R	0.41	0/44	0.78	0/55
All	All	0.63	0/2848	0.77	5/3818 (0.1%)

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	2	ARG	NE-CZ-NH1	6.11	123.35	120.30
1	B	270	ASP	CB-CG-OD2	-6.09	112.82	118.30
2	D	2	ARG	NE-CZ-NH2	-5.82	117.39	120.30
1	B	278	ASP	CB-CG-OD2	-5.51	113.34	118.30
2	C	2	ARG	NE-CZ-NH2	-5.18	117.71	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	658	0	646	30	0
1	B	649	0	640	31	0
1	M	654	0	637	10	0
1	N	645	0	631	13	0
2	C	45	0	45	0	0
2	D	45	0	45	1	0
2	P	45	0	45	1	0
2	R	45	0	45	1	0
3	A	82	0	0	2	0
3	B	68	0	0	1	2
3	C	11	0	0	0	2
3	D	10	0	0	0	0
3	M	1	0	0	0	0
All	All	2958	0	2734	53	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (53) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:N:268:GLU:O	1:N:272:ILE:HG13	1.77	0.84
1:B:288:MET:HG3	1:A:257[A]:VAL:HG12	1.71	0.73
1:B:272:ILE:HG21	1:A:272:ILE:CG2	2.21	0.71
1:B:292:MET:CE	1:A:292:MET:HE2	2.20	0.70
1:B:292:MET:HE2	1:A:292:MET:CE	2.22	0.69
1:B:272:ILE:CG2	1:A:272:ILE:CG2	2.71	0.69
1:B:257[A]:VAL:HG12	1:A:288:MET:HG3	1.75	0.68
1:B:288:MET:HG3	1:A:257[A]:VAL:CG1	2.23	0.68
1:N:272:ILE:HG21	1:M:272:ILE:HG21	1.74	0.68
1:B:281:HIS:O	3:B:301:HOH:O	2.12	0.68
1:B:272:ILE:CG2	1:A:272:ILE:HG21	2.24	0.67
1:A:271:GLU:HG2	3:A:369:HOH:O	1.97	0.64
1:N:263:ASP:O	1:N:267:GLN:OE1	2.15	0.63
1:A:291:GLN:HG2	3:A:362:HOH:O	1.97	0.63
1:B:292:MET:CE	1:A:292:MET:CE	2.75	0.63
1:B:257[A]:VAL:CG1	1:A:288:MET:HG3	2.29	0.62
1:B:257[A]:VAL:HG12	1:A:288:MET:CG	2.30	0.61
1:B:288:MET:CG	1:A:257[A]:VAL:HG12	2.32	0.60
1:B:241:TRP:O	1:A:223:ARG:NH1	2.28	0.58
1:B:292:MET:HE2	1:A:292:MET:HE2	1.83	0.58
1:N:282:GLN:HE21	1:M:258:PRO:HG3	1.69	0.58

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:N:272:ILE:CG2	1:M:272:ILE:HG21	2.35	0.57
1:B:295:GLY:HA2	1:A:247:ASN:O	2.07	0.54
1:A:224:THR:HG22	1:N:227:MET:CE	2.38	0.54
1:B:232:ALA:HB1	1:B:236:ARG:HH12	1.72	0.54
1:A:261:ARG:NH1	1:A:264:GLU:OE2	2.42	0.53
1:A:257[A]:VAL:HG13	1:A:262:TRP:HE1	1.74	0.53
1:B:292:MET:HE2	1:A:292:MET:HE3	1.91	0.53
1:N:282:GLN:NE2	1:M:258:PRO:HG3	2.24	0.53
1:B:257[A]:VAL:HG13	1:B:262:TRP:CD1	2.45	0.51
1:B:257[A]:VAL:HG13	1:B:262:TRP:NE1	2.27	0.50
1:A:257[A]:VAL:HG13	1:A:262:TRP:NE1	2.27	0.49
1:B:261:ARG:NH1	1:B:264:GLU:OE1	2.48	0.46
1:B:230:ASP:OD1	1:B:233:ARG:NH2	2.48	0.46
1:N:265:ALA:HA	1:M:280:TYR:CE1	2.51	0.46
1:B:254:GLU:HG2	2:D:1:GLU:O	2.17	0.45
1:B:257[A]:VAL:HG13	1:B:262:TRP:HE1	1.81	0.45
1:N:255:LEU:O	1:M:288:MET:N	2.47	0.45
1:B:251:ALA:HB1	1:A:273:ILE:HD11	1.98	0.45
1:N:272:ILE:HG21	1:M:272:ILE:CG2	2.45	0.44
1:B:242:ALA:HA	1:A:223:ARG:HB3	2.00	0.44
1:N:273:ILE:HD11	2:R:6:ARG:NH2	2.33	0.44
1:A:235:PHE:CE2	1:M:228:VAL:HA	2.53	0.43
1:B:257[B]:VAL:HG22	1:A:280:TYR:O	2.19	0.43
1:M:281:HIS:CE1	2:P:2:ARG:HB2	2.54	0.42
1:B:250:ARG:HG2	1:B:251:ALA:N	2.33	0.42
1:N:296:ARG:HG2	1:N:297:LYS:N	2.35	0.41
1:A:228:VAL:HG13	1:M:235:PHE:HD1	1.86	0.41
1:B:286:ILE:O	1:B:288:MET:HG2	2.21	0.41
1:N:247:ASN:O	1:N:247:ASN:OD1	2.39	0.41
1:A:257[A]:VAL:HG13	1:A:262:TRP:CD1	2.56	0.40
1:B:290:LEU:HD11	1:A:292:MET:HG3	2.03	0.40
1:B:286:ILE:HD12	1:A:259:PRO:HD3	2.03	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:B:322:HOH:O	3:C:107:HOH:O[4_545]	2.06	0.14
3:B:322:HOH:O	3:C:105:HOH:O[4_545]	2.14	0.06

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	76/90 (84%)	76 (100%)	0	0	100	100
1	B	75/90 (83%)	74 (99%)	1 (1%)	0	100	100
1	M	75/90 (83%)	74 (99%)	1 (1%)	0	100	100
1	N	74/90 (82%)	74 (100%)	0	0	100	100
2	C	4/6 (67%)	4 (100%)	0	0	100	100
2	D	4/6 (67%)	4 (100%)	0	0	100	100
2	P	4/6 (67%)	4 (100%)	0	0	100	100
2	R	4/6 (67%)	4 (100%)	0	0	100	100
All	All	316/384 (82%)	314 (99%)	2 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	69/80 (86%)	69 (100%)	0	100	100
1	B	68/80 (85%)	68 (100%)	0	100	100
1	M	68/80 (85%)	68 (100%)	0	100	100
1	N	67/80 (84%)	67 (100%)	0	100	100
2	C	4/4 (100%)	4 (100%)	0	100	100
2	D	4/4 (100%)	4 (100%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	P	4/4 (100%)	4 (100%)	0	100	100
2	R	4/4 (100%)	4 (100%)	0	100	100
All	All	288/336 (86%)	288 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	N	282	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	77/90 (85%)	0.20	0 100 100	18, 27, 52, 69	0
1	B	76/90 (84%)	0.19	1 (1%) 77 81	17, 28, 58, 72	0
1	M	77/90 (85%)	3.04	49 (63%) 0 0	76, 119, 154, 162	0
1	N	76/90 (84%)	2.84	40 (52%) 0 0	71, 113, 157, 168	0
2	C	6/6 (100%)	0.08	0 100 100	20, 21, 25, 35	0
2	D	6/6 (100%)	0.06	0 100 100	21, 22, 25, 38	0
2	P	6/6 (100%)	2.51	3 (50%) 0 0	95, 105, 121, 126	0
2	R	6/6 (100%)	2.05	3 (50%) 0 0	73, 87, 108, 126	0
All	All	330/384 (85%)	1.54	96 (29%) 1 1	17, 71, 147, 168	0

All (96) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	M	285	GLY	14.9
1	M	225	TYR	12.0
1	M	284	GLY	10.1
1	N	261	ARG	9.1
1	M	294	PHE	8.7
1	M	283	VAL	8.3
1	M	286	ILE	8.1
1	N	286	ILE	7.8
1	N	255	LEU	7.8
1	N	259	PRO	7.6
1	M	282	GLN	7.3
1	N	262	TRP	6.9
1	N	260	GLU	6.7
1	N	284	GLY	6.5
1	N	287	PRO	6.0
1	N	252	LEU	6.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	N	226	ILE	5.9
1	N	282	GLN	5.8
1	M	281	HIS	5.7
1	N	257	VAL	5.5
1	M	276	TRP	5.5
1	M	245	ARG	5.2
1	N	273	ILE	5.1
1	M	235	PHE	5.0
2	P	2	ARG	5.0
1	M	290	LEU	4.8
1	M	287	PRO	4.8
1	M	255	LEU	4.7
1	M	262	TRP	4.6
1	N	258	PRO	4.6
2	R	2	ARG	4.5
1	M	226	ILE	4.4
1	N	283	VAL	4.3
2	P	1	GLU	4.2
1	M	228	VAL	4.2
1	N	269	LEU	4.1
1	N	275	THR	4.1
1	M	292	MET	4.1
1	N	237	MET	3.9
1	M	296	ARG	3.7
1	M	224	THR	3.7
1	N	281	HIS	3.7
1	N	296	ARG	3.6
2	R	1	GLU	3.6
1	N	249	MET	3.6
1	N	276	TRP	3.5
1	M	223	ARG	3.4
1	N	292	MET	3.4
1	M	244	LYS	3.4
1	M	297	LYS	3.4
1	M	230	ASP	3.4
1	M	252	LEU	3.4
1	N	280	TYR	3.3
1	N	256	ASP	3.2
1	M	232	ALA	3.2
1	M	288	MET	3.2
1	N	241	TRP	3.2
1	N	235	PHE	3.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	P	6	ARG	3.0
1	M	249	MET	3.0
1	N	290	LEU	3.0
1	M	233	ARG	3.0
1	N	294	PHE	2.9
1	M	280	TYR	2.9
1	N	293	VAL	2.8
1	N	267	GLN	2.8
1	N	245	ARG	2.8
1	M	229	GLU	2.8
1	N	247	ASN	2.8
1	N	225	TYR	2.8
1	N	254	GLU	2.8
1	M	254	GLU	2.8
1	N	277	ALA	2.7
1	M	258	PRO	2.7
1	N	250	ARG	2.7
1	M	269	LEU	2.7
1	M	289	ILE	2.6
1	M	293	VAL	2.6
1	N	285	GLY	2.5
1	M	257	VAL	2.5
1	M	264	GLU	2.4
1	M	266	MET	2.4
1	M	274	ARG	2.4
1	M	259	PRO	2.4
1	N	228	VAL	2.4
1	M	263	ASP	2.4
1	M	271	GLU	2.2
1	N	291	GLN	2.2
1	M	260	GLU	2.2
1	M	246	PRO	2.2
1	M	256	ASP	2.2
1	M	273	ILE	2.1
1	B	225	TYR	2.1
1	M	298	GLU	2.1
1	M	261	ARG	2.1
2	R	3	GLY	2.1

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.