



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 15, 2017 – 05:25 am GMT

PDB ID : 4YNN
Title : Structure of Legionella pneumophila DegQ (S190A variant)
Authors : Wrase, R.; Hilgenfeld, R.; Hansen, G.
Deposited on : 2015-03-10
Resolution : 3.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.9-1692
EDS	:	trunk28620
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	recalc28949


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R _{free}	100719	1015 (3.22-3.18)
Clashscore	112137	1009 (3.20-3.20)
Ramachandran outliers	110173	1118 (3.22-3.18)
Sidechain outliers	110143	1117 (3.22-3.18)
RSRZ outliers	101464	1020 (3.22-3.18)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	448	<div><div>%</div><div><div></div><div>72%</div><div>17%</div><div>•</div><div>11%</div></div></div>
1	B	448	<div><div>%</div><div><div></div><div>74%</div><div>15%</div><div></div><div>11%</div></div></div>
1	C	448	<div><div>%</div><div><div></div><div>73%</div><div>16%</div><div></div><div>11%</div></div></div>
1	D	448	<div><div>%</div><div><div></div><div>72%</div><div>17%</div><div></div><div>11%</div></div></div>
1	E	448	<div><div>%</div><div><div></div><div>74%</div><div>15%</div><div></div><div>11%</div></div></div>
1	F	448	<div><div>%</div><div><div></div><div>77%</div><div>13%</div><div></div><div>11%</div></div></div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	G	448	 74% 15% 11%
1	H	448	 74% 15% 11%
2	I	8	 88% 13%
3	J	6	 100%
3	L	6	 67% 33%
4	K	7	 86% 14%
4	M	7	 86% 14%
4	N	7	 100%
4	O	7	 100%
4	P	7	 100%
5	Q	3	 67% 33%
5	R	3	 100%
5	S	3	 67% 33%
5	T	3	 100%
5	U	3	 100%
5	V	3	 100%

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 24195 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Protease DO.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	400	Total	C	N	O	S	0	0	0
			2979	1885	523	566	5			
1	B	399	Total	C	N	O	S	0	0	0
			2971	1880	522	565	4			
1	C	400	Total	C	N	O	S	0	0	0
			2979	1885	523	566	5			
1	D	400	Total	C	N	O	S	0	0	0
			2979	1885	523	566	5			
1	E	400	Total	C	N	O	S	0	0	0
			2979	1885	523	566	5			
1	F	400	Total	C	N	O	S	0	0	0
			2979	1885	523	566	5			
1	G	400	Total	C	N	O	S	0	0	0
			2979	1885	523	566	5			
1	H	400	Total	C	N	O	S	0	0	0
			2979	1885	523	566	5			

There are 104 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-11	MET	-	initiating methionine	UNP Q5ZVV9
A	-10	ARG	-	expression tag	UNP Q5ZVV9
A	-9	GLY	-	expression tag	UNP Q5ZVV9
A	-8	SER	-	expression tag	UNP Q5ZVV9
A	-7	HIS	-	expression tag	UNP Q5ZVV9
A	-6	HIS	-	expression tag	UNP Q5ZVV9
A	-5	HIS	-	expression tag	UNP Q5ZVV9
A	-4	HIS	-	expression tag	UNP Q5ZVV9
A	-3	HIS	-	expression tag	UNP Q5ZVV9
A	-2	HIS	-	expression tag	UNP Q5ZVV9
A	-1	GLY	-	expression tag	UNP Q5ZVV9
A	0	SER	-	expression tag	UNP Q5ZVV9
A	190	ALA	SER	engineered mutation	UNP Q5ZVV9

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-11	MET	-	initiating methionine	UNP Q5ZVV9
B	-10	ARG	-	expression tag	UNP Q5ZVV9
B	-9	GLY	-	expression tag	UNP Q5ZVV9
B	-8	SER	-	expression tag	UNP Q5ZVV9
B	-7	HIS	-	expression tag	UNP Q5ZVV9
B	-6	HIS	-	expression tag	UNP Q5ZVV9
B	-5	HIS	-	expression tag	UNP Q5ZVV9
B	-4	HIS	-	expression tag	UNP Q5ZVV9
B	-3	HIS	-	expression tag	UNP Q5ZVV9
B	-2	HIS	-	expression tag	UNP Q5ZVV9
B	-1	GLY	-	expression tag	UNP Q5ZVV9
B	0	SER	-	expression tag	UNP Q5ZVV9
B	190	ALA	SER	engineered mutation	UNP Q5ZVV9
C	-11	MET	-	initiating methionine	UNP Q5ZVV9
C	-10	ARG	-	expression tag	UNP Q5ZVV9
C	-9	GLY	-	expression tag	UNP Q5ZVV9
C	-8	SER	-	expression tag	UNP Q5ZVV9
C	-7	HIS	-	expression tag	UNP Q5ZVV9
C	-6	HIS	-	expression tag	UNP Q5ZVV9
C	-5	HIS	-	expression tag	UNP Q5ZVV9
C	-4	HIS	-	expression tag	UNP Q5ZVV9
C	-3	HIS	-	expression tag	UNP Q5ZVV9
C	-2	HIS	-	expression tag	UNP Q5ZVV9
C	-1	GLY	-	expression tag	UNP Q5ZVV9
C	0	SER	-	expression tag	UNP Q5ZVV9
C	190	ALA	SER	engineered mutation	UNP Q5ZVV9
D	-11	MET	-	initiating methionine	UNP Q5ZVV9
D	-10	ARG	-	expression tag	UNP Q5ZVV9
D	-9	GLY	-	expression tag	UNP Q5ZVV9
D	-8	SER	-	expression tag	UNP Q5ZVV9
D	-7	HIS	-	expression tag	UNP Q5ZVV9
D	-6	HIS	-	expression tag	UNP Q5ZVV9
D	-5	HIS	-	expression tag	UNP Q5ZVV9
D	-4	HIS	-	expression tag	UNP Q5ZVV9
D	-3	HIS	-	expression tag	UNP Q5ZVV9
D	-2	HIS	-	expression tag	UNP Q5ZVV9
D	-1	GLY	-	expression tag	UNP Q5ZVV9
D	0	SER	-	expression tag	UNP Q5ZVV9
D	190	ALA	SER	engineered mutation	UNP Q5ZVV9
E	-11	MET	-	initiating methionine	UNP Q5ZVV9
E	-10	ARG	-	expression tag	UNP Q5ZVV9
E	-9	GLY	-	expression tag	UNP Q5ZVV9

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
E	-8	SER	-	expression tag	UNP Q5ZVV9
E	-7	HIS	-	expression tag	UNP Q5ZVV9
E	-6	HIS	-	expression tag	UNP Q5ZVV9
E	-5	HIS	-	expression tag	UNP Q5ZVV9
E	-4	HIS	-	expression tag	UNP Q5ZVV9
E	-3	HIS	-	expression tag	UNP Q5ZVV9
E	-2	HIS	-	expression tag	UNP Q5ZVV9
E	-1	GLY	-	expression tag	UNP Q5ZVV9
E	0	SER	-	expression tag	UNP Q5ZVV9
E	190	ALA	SER	engineered mutation	UNP Q5ZVV9
F	-11	MET	-	initiating methionine	UNP Q5ZVV9
F	-10	ARG	-	expression tag	UNP Q5ZVV9
F	-9	GLY	-	expression tag	UNP Q5ZVV9
F	-8	SER	-	expression tag	UNP Q5ZVV9
F	-7	HIS	-	expression tag	UNP Q5ZVV9
F	-6	HIS	-	expression tag	UNP Q5ZVV9
F	-5	HIS	-	expression tag	UNP Q5ZVV9
F	-4	HIS	-	expression tag	UNP Q5ZVV9
F	-3	HIS	-	expression tag	UNP Q5ZVV9
F	-2	HIS	-	expression tag	UNP Q5ZVV9
F	-1	GLY	-	expression tag	UNP Q5ZVV9
F	0	SER	-	expression tag	UNP Q5ZVV9
F	190	ALA	SER	engineered mutation	UNP Q5ZVV9
G	-11	MET	-	initiating methionine	UNP Q5ZVV9
G	-10	ARG	-	expression tag	UNP Q5ZVV9
G	-9	GLY	-	expression tag	UNP Q5ZVV9
G	-8	SER	-	expression tag	UNP Q5ZVV9
G	-7	HIS	-	expression tag	UNP Q5ZVV9
G	-6	HIS	-	expression tag	UNP Q5ZVV9
G	-5	HIS	-	expression tag	UNP Q5ZVV9
G	-4	HIS	-	expression tag	UNP Q5ZVV9
G	-3	HIS	-	expression tag	UNP Q5ZVV9
G	-2	HIS	-	expression tag	UNP Q5ZVV9
G	-1	GLY	-	expression tag	UNP Q5ZVV9
G	0	SER	-	expression tag	UNP Q5ZVV9
G	190	ALA	SER	engineered mutation	UNP Q5ZVV9
H	-11	MET	-	initiating methionine	UNP Q5ZVV9
H	-10	ARG	-	expression tag	UNP Q5ZVV9
H	-9	GLY	-	expression tag	UNP Q5ZVV9
H	-8	SER	-	expression tag	UNP Q5ZVV9
H	-7	HIS	-	expression tag	UNP Q5ZVV9
H	-6	HIS	-	expression tag	UNP Q5ZVV9

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
H	-5	HIS	-	expression tag	UNP Q5ZVV9
H	-4	HIS	-	expression tag	UNP Q5ZVV9
H	-3	HIS	-	expression tag	UNP Q5ZVV9
H	-2	HIS	-	expression tag	UNP Q5ZVV9
H	-1	GLY	-	expression tag	UNP Q5ZVV9
H	0	SER	-	expression tag	UNP Q5ZVV9
H	190	ALA	SER	engineered mutation	UNP Q5ZVV9

- Molecule 2 is a protein called Octapeptide.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	I	8	Total	C	N	O	0	0	0
			40	24	8	8			

- Molecule 3 is a protein called Hexa-peptide.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	J	6	Total	C	N	O	0	0	0
			30	18	6	6			
3	L	6	Total	C	N	O	0	0	0
			30	18	6	6			

- Molecule 4 is a protein called Hepta-peptide.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	K	7	Total	C	N	O	0	0	0
			35	21	7	7			
4	M	7	Total	C	N	O	0	0	0
			35	21	7	7			
4	N	7	Total	C	N	O	0	0	0
			35	21	7	7			
4	O	7	Total	C	N	O	0	0	0
			35	21	7	7			
4	P	7	Total	C	N	O	0	0	0
			35	21	7	7			

- Molecule 5 is a protein called UNK-UNK-UNK.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
5	Q	3	Total	C	N	O	0	0	0
			16	9	3	4			



Continued on next page...

Continued from previous page...




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
5	R	3	Total	C	N	O	0	0	0
			16	9	3	4			
5	S	3	Total	C	N	O	0	0	0
			16	9	3	4			
5	T	3	Total	C	N	O	0	0	0
			16	9	3	4			
5	U	3	Total	C	N	O	0	0	0
			16	9	3	4			
5	V	3	Total	C	N	O	0	0	0
			16	9	3	4			




- Molecule 1: Protease DO



- Molecule 1: Protease DO



- Molecule 2: Octapeptide



- Molecule 3: Hexa-peptide




There are no outlier residues recorded for this chain.

- Molecule 3: Hexa-peptide




- Molecule 4: Hepta-peptide

Chain K:  86% 14%



- Molecule 4: Hepta-peptide

Chain M:  86% 14%



- Molecule 4: Hepta-peptide

Chain N:  100%

There are no outlier residues recorded for this chain.

- Molecule 4: Hepta-peptide

Chain O:  100%

There are no outlier residues recorded for this chain.

- Molecule 4: Hepta-peptide

Chain P:  100%

There are no outlier residues recorded for this chain.

- Molecule 5: UNK-UNK-UNK

Chain Q:  67% 33%



- Molecule 5: UNK-UNK-UNK

Chain R:  100%

There are no outlier residues recorded for this chain.

- Molecule 5: UNK-UNK-UNK

Chain S:  67% 33%



- Molecule 5: UNK-UNK-UNK

Chain T:  100%

There are no outlier residues recorded for this chain.

- Molecule 5: UNK-UNK-UNK

Chain U:  100%

There are no outlier residues recorded for this chain.

- Molecule 5: UNK-UNK-UNK

Chain V:  100%

There are no outlier residues recorded for this chain.

4 Data and refinement statistics

Property	Value	Source
Space group	P 63	Depositor
Cell constants a, b, c, α , β , γ	147.91Å 147.91Å 383.51Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	28.54 – 3.20 28.54 – 3.20	Depositor EDS
% Data completeness (in resolution range)	99.0 (28.54-3.20) 99.0 (28.54-3.20)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.18 (at 3.17Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8.1_1168)	Depositor
R, R_{free}	0.181 , 0.219 0.180 , 0.219	Depositor DCC
R_{free} test set	3874 reflections (5.03%)	DCC
Wilson B-factor (Å ²)	72.1	Xtriage
Anisotropy	0.002	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 55.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.28$	Xtriage
Estimated twinning fraction	0.066 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	24195	wwPDB-VP
Average B, all atoms (Å ²)	60.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 1.53% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.24	0/3018	0.46	0/4089
1	B	0.25	0/3010	0.47	0/4078
1	C	0.25	0/3018	0.47	0/4089
1	D	0.24	0/3018	0.45	0/4089
1	E	0.25	0/3018	0.47	0/4089
1	F	0.25	0/3018	0.46	0/4089
1	G	0.24	0/3018	0.46	0/4089
1	H	0.25	0/3018	0.47	0/4089
All	All	0.25	0/24136	0.47	0/32701

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2979	0	3088	45	0
1	B	2971	0	3080	40	0
1	C	2979	0	3088	45	0
1	D	2979	0	3088	46	0
1	E	2979	0	3088	45	0
1	F	2979	0	3088	33	0
1	G	2979	0	3088	41	0
1	H	2979	0	3088	40	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	I	40	0	11	1	0
3	J	30	0	8	0	0
3	L	30	0	8	2	0
4	K	35	0	9	1	0
4	M	35	0	9	1	0
4	N	35	0	9	0	0
4	O	35	0	9	0	0
4	P	35	0	9	0	0
5	Q	16	0	5	1	0
5	R	16	0	5	0	0
5	S	16	0	6	1	0
5	T	16	0	5	0	0
5	U	16	0	5	0	0
5	V	16	0	5	0	0
All	All	24195	0	24799	329	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 329 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:125:LEU:HD11	1:E:201:LEU:HB2	1.58	0.85
1:D:23:ASN:HD21	1:D:144:ASN:HD22	1.25	0.84
1:D:230:VAL:HG13	1:D:240:ILE:HD11	1.67	0.77
1:D:125:LEU:HD11	1:D:201:LEU:HB2	1.66	0.75
1:B:230:VAL:HG13	1:B:240:ILE:HD11	1.69	0.72

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	396/448 (88%)	388 (98%)	8 (2%)	0	100	100
1	B	395/448 (88%)	387 (98%)	8 (2%)	0	100	100
1	C	396/448 (88%)	387 (98%)	9 (2%)	0	100	100
1	D	396/448 (88%)	387 (98%)	9 (2%)	0	100	100
1	E	396/448 (88%)	388 (98%)	8 (2%)	0	100	100
1	F	396/448 (88%)	388 (98%)	8 (2%)	0	100	100
1	G	396/448 (88%)	389 (98%)	7 (2%)	0	100	100
1	H	396/448 (88%)	387 (98%)	9 (2%)	0	100	100
All	All	3167/3584 (88%)	3101 (98%)	66 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	322/362 (89%)	315 (98%)	7 (2%)	57	85
1	B	321/362 (89%)	319 (99%)	2 (1%)	89	96
1	C	322/362 (89%)	316 (98%)	6 (2%)	62	86
1	D	322/362 (89%)	316 (98%)	6 (2%)	62	86
1	E	322/362 (89%)	318 (99%)	4 (1%)	75	91
1	F	322/362 (89%)	319 (99%)	3 (1%)	82	94
1	G	322/362 (89%)	319 (99%)	3 (1%)	82	94
1	H	322/362 (89%)	316 (98%)	6 (2%)	62	86
All	All	2575/2896 (89%)	2538 (99%)	37 (1%)	71	90

5 of 37 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	76	ILE
1	D	416	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	H	345	LEU
1	D	92	THR
1	D	189	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 18 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	23	ASN
1	D	155	GLN
1	F	233	GLN
1	B	233	GLN
1	C	233	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	400/448 (89%)	-0.37	4 (1%) 82 72	23, 58, 90, 112	0
1	B	399/448 (89%)	-0.35	3 (0%) 86 77	23, 51, 85, 112	0
1	C	400/448 (89%)	-0.40	5 (1%) 77 65	22, 49, 81, 117	0
1	D	400/448 (89%)	-0.23	6 (1%) 74 61	25, 62, 99, 127	0
1	E	400/448 (89%)	-0.35	4 (1%) 82 72	25, 53, 93, 129	0
1	F	400/448 (89%)	-0.33	5 (1%) 77 65	31, 60, 95, 119	0
1	G	400/448 (89%)	-0.32	4 (1%) 82 72	27, 59, 92, 120	0
1	H	400/448 (89%)	-0.32	6 (1%) 74 61	27, 58, 91, 125	0
2	I	0/8	-	-	-	-
3	J	0/6	-	-	-	-
3	L	0/6	-	-	-	-
4	K	0/7	-	-	-	-
4	M	0/7	-	-	-	-
4	N	0/7	-	-	-	-
4	O	0/7	-	-	-	-
4	P	0/7	-	-	-	-
5	Q	0/3	-	-	-	-
5	R	0/3	-	-	-	-
5	S	0/3	-	-	-	-
5	T	0/3	-	-	-	-
5	U	0/3	-	-	-	-
5	V	0/3	-	-	-	-
All	All	3199/3657 (87%)	-0.34	37 (1%) 79 67	22, 56, 92, 129	0

The worst 5 of 37 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	212	TYR	4.1
1	H	8	PRO	3.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	8	PRO	3.8
1	G	29	TYR	3.7
1	H	414	HIS	3.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.