



wwPDB X-ray Structure Validation Summary Report

Feb 15, 2017 – 03:45 am GMT

PDB ID : 2YZC
Title : Crystal structure of uricase from *Arthrobacter globiformis* in complex with allantoinate
Authors : Juan, E.C.M.; Hossain, M.T.; Hoque, M.M.; Suzuki, K.; Sekiguchi, T.; Takemura, A.
Deposited on : 2007-05-05
Resolution : 1.88 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the  symbol.

The following versions of software and data (see [references](#) ) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : trunk28620
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.88 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	7505 (1.90-1.86)
Clashscore	112137	8369 (1.90-1.86)
Ramachandran outliers	110173	8279 (1.90-1.86)
Sidechain outliers	110143	8280 (1.90-1.86)
RSRZ outliers	101464	7571 (1.90-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	302	
1	B	302	
1	C	302	
1	D	302	
1	E	302	
1	F	302	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	G	302	<p>2% 82% 12% • 5%</p>
1	H	302	<p>% 83% 11% 5%</p>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	1AL	A	303	-	-	X	X
2	1AL	B	303	-	-	-	X
2	1AL	B	304	-	-	-	X
2	1AL	D	304	-	-	-	X
2	1AL	E	308	-	-	X	X
2	1AL	F	306	-	-	-	X
2	1AL	G	307	-	-	X	X
2	1AL	H	305	-	-	-	X

2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 20334 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Uricase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	287	2281	1435	408	435	3	0	0	0
1	B	287	2281	1435	408	435	3	0	0	0
1	C	287	2281	1435	408	435	3	0	0	0
1	D	287	2281	1435	408	435	3	0	0	0
1	E	287	2281	1435	408	435	3	0	0	0
1	F	287	2281	1435	408	435	3	0	0	0
1	G	287	2281	1435	408	435	3	0	0	0
1	H	287	2281	1435	408	435	3	0	0	0

- Molecule 2 is ALLANTOATE ION (three-letter code: 1AL) (formula: C₄H₇N₄O₄).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
2	A	1	Total 12	C 4	N 4	O 4	0	0
2	B	1	Total 12	C 4	N 4	O 4	0	0
2	B	1	Total 12	C 4	N 4	O 4	0	0
2	D	1	Total 12	C 4	N 4	O 4	0	0
2	E	1	Total 12	C 4	N 4	O 4	0	0
2	F	1	Total 12	C 4	N 4	O 4	0	0
2	G	1	Total 12	C 4	N 4	O 4	0	0
2	H	1	Total 12	C 4	N 4	O 4	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	253	Total 253	O 253	0	0
3	B	251	Total 251	O 251	0	0
3	C	248	Total 248	O 248	0	0
3	D	254	Total 254	O 254	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	E	230	Total 230	O 230	0	0
3	F	268	Total 268	O 268	0	0
3	G	245	Total 245	O 245	0	0
3	H	241	Total 241	O 241	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Uricase




- Molecule 1: Uricase



- Molecule 1: Uricase




- Molecule 1: Uricase








• Molecule 1: Uricase




• Molecule 1: Uricase



• Molecule 1: Uricase



• Molecule 1: Uricase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	85.50Å 122.94Å 284.78Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 1.88 49.98 – 1.88	Depositor EDS
% Data completeness (in resolution range)	95.6 (20.00-1.88) 95.6 (49.98-1.88)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.48 (at 1.88Å)	Xtrriage
Refinement program	REFMAC 5.1.24	Depositor
R, R_{free}	0.159 , 0.195 0.182 , 0.210	Depositor DCC
R_{free} test set	22669 reflections (9.78%)	DCC
Wilson B-factor (Å ²)	23.1	Xtrriage
Anisotropy	0.574	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 48.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	20334	wwPDB-VP
Average B, all atoms (Å ²)	26.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.73% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 1AL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.32	0/2333	0.62	0/3167
1	B	0.31	0/2333	0.60	0/3167
1	C	0.32	0/2333	0.61	0/3167
1	D	0.31	0/2333	0.60	0/3167
1	E	0.31	0/2333	0.59	0/3167
1	F	0.32	0/2333	0.62	0/3167
1	G	0.32	0/2333	0.61	0/3167
1	H	0.32	0/2333	0.61	0/3167
All	All	0.32	0/18664	0.61	0/25336

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2281	0	2177	26	0
1	B	2281	0	2177	37	0
1	C	2281	0	2177	35	0
1	D	2281	0	2177	31	0
1	E	2281	0	2177	50	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	F	2281	0	2177	40	0
1	G	2281	0	2177	41	0
1	H	2281	0	2177	31	0
2	A	12	0	6	4	0
2	B	24	0	12	5	0
2	D	12	0	6	2	0
2	E	12	0	6	4	0
2	F	12	0	6	2	0
2	G	12	0	6	4	0
2	H	12	0	6	2	0
3	A	253	0	0	0	0
3	B	251	0	0	1	0
3	C	248	0	0	1	0
3	D	254	0	0	0	0
3	E	230	0	0	1	0
3	F	268	0	0	1	0
3	G	245	0	0	3	0
3	H	241	0	0	2	0
All	All	20334	0	17464	262	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 262 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:67:THR:H	2:E:308:1AL:H111	1.04	1.03
2:B:303:1AL:H111	1:C:67:THR:H	1.04	1.00
1:H:67:THR:H	2:H:305:1AL:H111	1.07	0.98
1:F:131:GLU:HB2	1:F:150:ILE:HD11	1.47	0.96
2:A:303:1AL:H111	1:D:67:THR:H	1.07	0.96

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	285/302 (94%)	279 (98%)	6 (2%)	0	100	100
1	B	285/302 (94%)	281 (99%)	4 (1%)	0	100	100
1	C	285/302 (94%)	279 (98%)	6 (2%)	0	100	100
1	D	285/302 (94%)	282 (99%)	3 (1%)	0	100	100
1	E	285/302 (94%)	281 (99%)	4 (1%)	0	100	100
1	F	285/302 (94%)	282 (99%)	3 (1%)	0	100	100
1	G	285/302 (94%)	281 (99%)	4 (1%)	0	100	100
1	H	285/302 (94%)	282 (99%)	3 (1%)	0	100	100
All	All	2280/2416 (94%)	2247 (99%)	33 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	236/250 (94%)	230 (98%)	6 (2%)	53	42
1	B	236/250 (94%)	232 (98%)	4 (2%)	66	59
1	C	236/250 (94%)	230 (98%)	6 (2%)	53	42
1	D	236/250 (94%)	232 (98%)	4 (2%)	66	59
1	E	236/250 (94%)	231 (98%)	5 (2%)	59	50
1	F	236/250 (94%)	232 (98%)	4 (2%)	66	59
1	G	236/250 (94%)	232 (98%)	4 (2%)	66	59
1	H	236/250 (94%)	232 (98%)	4 (2%)	66	59
All	All	1888/2000 (94%)	1851 (98%)	37 (2%)	60	52

5 of 37 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	33	ASN
1	E	15	LEU
1	H	193	ASN
1	D	193	ASN
1	D	211	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 88 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	224	GLN
1	E	119	ASN
1	H	111	GLN
1	D	237	HIS
1	D	284	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	1AL	A	303	-	8,11,11	1.03	0	9,14,14	4.00	4 (44%)
2	1AL	B	303	-	8,11,11	0.99	0	9,14,14	4.06	4 (44%)
2	1AL	B	304	-	8,11,11	1.01	0	9,14,14	4.06	4 (44%)
2	1AL	D	304	-	8,11,11	0.98	0	9,14,14	4.03	4 (44%)
2	1AL	E	308	-	8,11,11	0.91	0	9,14,14	4.10	4 (44%)
2	1AL	F	306	-	8,11,11	1.03	0	9,14,14	4.07	4 (44%)
2	1AL	G	307	-	8,11,11	1.02	0	9,14,14	4.00	4 (44%)
2	1AL	H	305	-	8,11,11	1.01	0	9,14,14	4.07	4 (44%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	1AL	A	303	-	-	2/8/12/12	0/0/0/0
2	1AL	B	303	-	-	2/8/12/12	0/0/0/0
2	1AL	B	304	-	-	2/8/12/12	0/0/0/0
2	1AL	D	304	-	-	2/8/12/12	0/0/0/0
2	1AL	E	308	-	-	2/8/12/12	0/0/0/0
2	1AL	F	306	-	-	2/8/12/12	0/0/0/0
2	1AL	G	307	-	-	2/8/12/12	0/0/0/0
2	1AL	H	305	-	-	2/8/12/12	0/0/0/0

There are no bond length outliers.

The worst 5 of 32 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	E	308	1AL	O2-C1-N4	-2.14	113.85	121.56
2	B	303	1AL	O2-C1-N4	-2.12	113.94	121.56
2	D	304	1AL	O2-C1-N4	-2.09	114.02	121.56
2	G	307	1AL	O2-C1-N4	-2.09	114.05	121.56
2	F	306	1AL	O2-C1-N4	-2.08	114.07	121.56

There are no chirality outliers.

5 of 16 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	H	305	1AL	O10-C9-N12-C8
2	D	304	1AL	O10-C9-N12-C8
2	E	308	1AL	O10-C9-N12-C8
2	A	303	1AL	O10-C9-N12-C8
2	F	306	1AL	O10-C9-N12-C8

There are no ring outliers.

8 monomers are involved in 23 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	303	1AL	4	0
2	B	303	1AL	2	0
2	B	304	1AL	3	0
2	D	304	1AL	2	0
2	E	308	1AL	4	0
2	F	306	1AL	2	0
2	G	307	1AL	4	0
2	H	305	1AL	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	287/302 (95%)	-0.17	3 (1%) 82 84	15, 22, 34, 49	0
1	B	287/302 (95%)	-0.18	4 (1%) 75 78	17, 24, 38, 54	0
1	C	287/302 (95%)	-0.17	3 (1%) 82 84	16, 25, 34, 54	0
1	D	287/302 (95%)	-0.16	3 (1%) 82 84	17, 24, 38, 56	0
1	E	287/302 (95%)	-0.07	6 (2%) 64 66	18, 27, 41, 56	0
1	F	287/302 (95%)	-0.18	5 (1%) 70 73	16, 23, 35, 51	0
1	G	287/302 (95%)	-0.15	6 (2%) 64 66	17, 24, 39, 54	0
1	H	287/302 (95%)	-0.15	4 (1%) 75 78	17, 25, 35, 48	0
All	All	2296/2416 (95%)	-0.15	34 (1%) 74 76	15, 24, 37, 56	0

The worst 5 of 34 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	33	ASN	4.1
1	F	11	THR	4.0
1	B	196	GLU	3.9
1	B	297	ASN	3.7
1	H	297	ASN	3.7

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	1AL	D	304	12/12	0.67	0.34	11.30	37,44,49,50	0
2	1AL	B	303	12/12	0.69	0.26	10.47	35,43,48,49	0
2	1AL	A	303	12/12	0.73	0.30	10.36	31,41,45,47	0
2	1AL	H	305	12/12	0.64	0.25	9.83	37,45,49,49	0
2	1AL	B	304	12/12	0.68	0.26	9.25	42,48,51,52	0
2	1AL	F	306	12/12	0.73	0.25	8.67	33,38,45,47	0
2	1AL	E	308	12/12	0.77	0.24	8.29	38,46,49,50	0
2	1AL	G	307	12/12	0.75	0.23	5.56	30,39,44,47	0

6.5 Other polymers [i](#)

There are no such residues in this entry.