



# Full wwPDB X-ray Structure Validation Report ⓘ

Oct 4, 2017 – 07:19 PM EDT

PDB ID : 1Z0C  
Title : Crystal Structure of A. fulgidus Lon proteolytic domain D508A mutant  
Authors : Botos, I.; Melnikov, E.E.; Cherry, S.; Kozlov, S.; Makhovskaya, O.V.; Tropea, J.E.; Gustchina, A.; Rotanova, T.V.; Wlodawer, A.  
Deposited on : unknown  
Resolution : 1.55 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20030345
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20030345


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.55 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	1088 (1.56-1.56)
Clashscore	112137	1132 (1.56-1.56)
Ramachandran outliers	110173	1110 (1.56-1.56)
Sidechain outliers	110143	1108 (1.56-1.56)
RSRZ outliers	101464	1089 (1.56-1.56)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	207	<div> <div>16%</div> <div>75%</div> <div>20%</div> <div>...</div> </div>

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 1646 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Putative protease La homolog type.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	200	Total	C	N	O	S	0	2	0
			1505	953	252	294	6			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	508	ALA	ASP	ENGINEERED	UNP O29883


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	141	Total	O	0	0
			141	141		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Putative protease La homolog type



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 65	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	81.96 Å 81.96 Å 41.50 Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	20.00 – 1.55 19.92 – 1.55	Depositor EDS
% Data completeness (in resolution range)	100.0 (20.00-1.55) 100.0 (19.92-1.55)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.28 (at 1.55 Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, $R_{free}$	0.209 , 0.256 0.209 , 0.259	Depositor DCC
$R_{free}$ test set	1181 reflections (5.35%)	DCC
Wilson B-factor (Å <sup>2</sup> )	26.2	Xtriage
Anisotropy	0.055	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 56.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.046 for h,-h-k,-l	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	1646	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	34.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 9.10% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.81	0/1531	0.86	1/2065 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	A	483	LYS	CD-CE-NZ	5.79	125.01	111.70

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1505	0	1566	38	0
2	A	141	0	0	20	0
All	All	1646	0	1566	38	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

All (38) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance ( $\text{\AA}$ )	Clash overlap ( $\text{\AA}$ )
1:A:576:LEU:HD22	1:A:581:GLU:HB3	1.46	0.94

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:563:LYS:HE3	2:A:32:HOH:O	1.72	0.88
1:A:428:ARG:HG3	2:A:138:HOH:O	1.84	0.77
1:A:576:LEU:H	1:A:576:LEU:HD12	1.54	0.72
1:A:576:LEU:O	1:A:577:ASP:HB2	1.93	0.69
1:A:580:HIS:HE1	2:A:54:HOH:O	1.78	0.66
1:A:503:GLU:HA	2:A:140:HOH:O	1.95	0.66
1:A:418:LEU:HD22	2:A:64:HOH:O	1.97	0.64
1:A:428:ARG:CG	2:A:138:HOH:O	2.43	0.64
1:A:508:ALA:HB3	2:A:8:HOH:O	1.99	0.63
1:A:419:PHE:HB3	1:A:444:PRO:HG3	1.79	0.62
1:A:574:VAL:HG12	1:A:576:LEU:HG	1.80	0.62
1:A:488:ASP:OD2	1:A:491:ASN:ND2	2.27	0.62
1:A:501:THR:HB	2:A:78:HOH:O	1.99	0.61
1:A:418:LEU:HA	2:A:70:HOH:O	2.00	0.61
1:A:435:ILE:HG13	1:A:439:ALA:HB3	1.84	0.60
1:A:416:TYR:N	2:A:100:HOH:O	2.34	0.59
1:A:580:HIS:CD2	2:A:111:HOH:O	2.57	0.58
1:A:469:ILE:HD12	1:A:509:SER:HA	1.86	0.57
1:A:598:HIS:HE1	2:A:43:HOH:O	1.88	0.56
1:A:581:GLU:HG2	2:A:139:HOH:O	2.05	0.56
1:A:532:ALA:HB2	1:A:561:LEU:HD13	1.90	0.53
1:A:508:ALA:CB	2:A:8:HOH:O	2.57	0.51
1:A:580:HIS:CE1	2:A:54:HOH:O	2.57	0.51
1:A:428:ARG:NE	2:A:138:HOH:O	2.43	0.51
1:A:441:ILE:HG22	1:A:442:VAL:O	2.13	0.49
1:A:424:TYR:HB3	1:A:526:PRO:HB2	1.96	0.47
1:A:419:PHE:CE1	1:A:559:ALA:HB1	2.50	0.47
1:A:418:LEU:O	1:A:428:ARG:NH2	2.48	0.46
1:A:458:GLY:HA3	1:A:492:MET:O	2.17	0.45
1:A:472:GLU:HG3	2:A:53:HOH:O	2.18	0.43
1:A:577:ASP:HB3	2:A:139:HOH:O	2.19	0.43
1:A:540:LYS:HE2	1:A:540:LYS:HB3	1.63	0.42
1:A:469:ILE:HA	1:A:469:ILE:HD13	1.88	0.42
1:A:487:ARG:NH2	1:A:523:GLU:OE1	2.53	0.42
1:A:471:ARG:NH2	2:A:96:HOH:O	2.43	0.42
1:A:572:ASP:CG	2:A:137:HOH:O	2.59	0.41
1:A:581:GLU:HA	1:A:582:GLY:HA2	1.63	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	198/207 (96%)	191 (96%)	6 (3%)	1 (0%)	32	9

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	577	ASP

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	166/169 (98%)	159 (96%)	7 (4%)	34	6

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	438	SER
1	A	465	ARG
1	A	511	SER
1	A	538	SER
1	A	575	LEU
1	A	576	LEU
1	A	616	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.



### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	200/207 (96%)	1.07	33 (16%) 2 2	17, 30, 54, 65	0

All (33) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	501	THR	8.7
1	A	465	ARG	7.0
1	A	502	TYR	6.1
1	A	580	HIS	5.5
1	A	618	LEU	5.4
1	A	459	ARG	4.9
1	A	466	LEU	4.9
1	A	576	LEU	4.8
1	A	578	ALA	4.7
1	A	579	GLU	4.4
1	A	457	GLU	4.2
1	A	416	TYR	3.9
1	A	418	LEU	3.9
1	A	503	GLU	3.7
1	A	575	LEU	3.7
1	A	577	ASP	3.6
1	A	574	VAL	3.5
1	A	614	LYS	3.5
1	A	581	GLU	3.4
1	A	617	GLU	2.9
1	A	571	ILE	2.8
1	A	441	ILE	2.8
1	A	453	MET	2.8
1	A	506	GLU	2.8
1	A	615	GLU	2.7
1	A	468	GLU	2.7
1	A	616	LEU	2.6

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	464	GLY	2.4
1	A	434	VAL	2.3
1	A	509	SER	2.2
1	A	422	GLU	2.2
1	A	471	ARG	2.1
1	A	481	ILE	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.