



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 13, 2017 – 07:52 am GMT

PDB ID : 2Z3Q
Title : Crystal structure of the IL-15/IL-15Ra complex
Authors : Chirifu, M.; Yamagata, Y.; Davis, S.J.; Ikemizu, S.
Deposited on : 2007-06-05
Resolution : 1.85 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtrriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	112137	2083 (1.86-1.86)
Ramachandran outliers	110173	2060 (1.86-1.86)
Sidechain outliers	110143	2060 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	119	
1	C	119	
2	B	107	
2	D	107	

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 3263 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Interleukin-15.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	114	891	560	141	182	8	0	0	0
1	C	117	910	571	146	185	8	0	0	0

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-4	ALA	-	EXPRESSION TAG	UNP P40933
A	-3	MET	-	EXPRESSION TAG	UNP P40933
A	-2	ALA	-	EXPRESSION TAG	UNP P40933
A	-1	ILE	-	EXPRESSION TAG	UNP P40933
A	0	SER	-	EXPRESSION TAG	UNP P40933
C	-4	ALA	-	EXPRESSION TAG	UNP P40933
C	-3	MET	-	EXPRESSION TAG	UNP P40933
C	-2	ALA	-	EXPRESSION TAG	UNP P40933
C	-1	ILE	-	EXPRESSION TAG	UNP P40933
C	0	SER	-	EXPRESSION TAG	UNP P40933

- Molecule 2 is a protein called Interleukin-15 receptor alpha chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	81	629	397	113	114	5	0	0	0
2	D	77	605	381	109	109	6	0	0	0

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	-4	ALA	-	EXPRESSION TAG	UNP Q13261
B	-3	MET	-	EXPRESSION TAG	UNP Q13261

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-2	ALA	-	EXPRESSION TAG	UNP Q13261
B	-1	ILE	-	EXPRESSION TAG	UNP Q13261
B	0	SER	-	EXPRESSION TAG	UNP Q13261
D	-4	ALA	-	EXPRESSION TAG	UNP Q13261
D	-3	MET	-	EXPRESSION TAG	UNP Q13261
D	-2	ALA	-	EXPRESSION TAG	UNP Q13261
D	-1	ILE	-	EXPRESSION TAG	UNP Q13261
D	0	SER	-	EXPRESSION TAG	UNP Q13261

- Molecule 3 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	48	Total O 48 48	0	0
3	B	46	Total O 46 46	0	0
3	C	53	Total O 53 53	0	0
3	D	81	Total O 81 81	0	0

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

- Molecule 1: Interleukin-15




- Molecule 1: Interleukin-15



- Molecule 2: Interleukin-15 receptor alpha chain



- Molecule 2: Interleukin-15 receptor alpha chain



4 Data and refinement statistics

Xtrriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	78.44Å 120.01Å 49.46Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	25.55 – 1.85	Depositor
% Data completeness (in resolution range)	100.0 (25.55-1.85)	Depositor
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.202 , 0.252	Depositor
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	3263	wwPDB-VP
Average B, all atoms (Å ²)	15.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.85	1/901 (0.1%)	0.80	0/1215
1	C	0.83	0/921	0.81	0/1245
2	B	0.74	0/647	0.81	0/883
2	D	0.89	1/620 (0.2%)	0.87	0/842
All	All	0.83	2/3089 (0.1%)	0.82	0/4185

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	D	0	1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	99	PHE	CE1-CZ	5.39	1.47	1.37
2	D	3	CYS	CB-SG	5.26	1.91	1.82

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	D	-2	ALA	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	891	0	883	19	0
1	C	910	0	901	12	0
2	B	629	0	630	12	0
2	D	605	0	608	19	0
3	A	48	0	0	3	0
3	B	46	0	0	2	0
3	C	53	0	0	1	0
3	D	81	0	0	1	0
All	All	3263	0	3022	59	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

The worst 5 of 59 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:112:ASN:H	1:A:113:THR:HA	1.04	1.16
1:A:112:ASN:N	1:A:113:THR:HA	1.63	1.08
2:B:47:LEU:HD13	2:B:49:LYS:HD3	1.41	1.00
2:D:-3:MET:HB2	2:D:-1:ILE:H	1.32	0.92
1:C:1:ASN:HD21	1:C:72:ASN:HD22	1.16	0.90

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	110/119 (92%)	102 (93%)	6 (6%)	2 (2%)	10 2
1	C	115/119 (97%)	110 (96%)	4 (4%)	1 (1%)	20 7

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	B	79/107 (74%)	74 (94%)	4 (5%)	1 (1%)	14	3
2	D	75/107 (70%)	71 (95%)	4 (5%)	0	100	100
All	All	379/452 (84%)	357 (94%)	18 (5%)	4 (1%)	17	5

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	19	MET
1	A	112	ASN
2	B	48	ASN
1	C	31	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	106/110 (96%)	98 (92%)	8 (8%)	16	3
1	C	108/110 (98%)	99 (92%)	9 (8%)	13	2
2	B	72/92 (78%)	68 (94%)	4 (6%)	25	8
2	D	69/92 (75%)	66 (96%)	3 (4%)	33	15
All	All	355/404 (88%)	331 (93%)	24 (7%)	18	5

5 of 24 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	49	LYS
1	C	34	SER
2	D	1	ILE
2	B	65	ARG
1	C	7	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 6 such sidechains are listed below:

Mol	Chain	Res	Type
2	B	30	ASN
1	C	72	ASN
2	B	48	ASN
2	B	11	HIS
1	C	32	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.