



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 15, 2017 – 05:11 am GMT

PDB ID : 4Z3W
Title : Active site complex BamBC of Benzoyl Coenzyme A reductase in complex with 1,5 Dienoyl-CoA
Authors : Weinert, T.; Kung, J.; Weidenweber, S.; Huwiler, S.; Boll, M.; Ermler, U.
Deposited on : 2015-04-01
Resolution : 2.21 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : trunk28620
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.21 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.





Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	4002 (2.20-2.20)
Clashscore	112137	4730 (2.20-2.20)
Ramachandran outliers	110173	4656 (2.20-2.20)
Sidechain outliers	110143	4657 (2.20-2.20)
RSRZ outliers	101464	4033 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	653	<div> <div>7%</div> <div>84%</div> <div>15%</div> <div>.</div> </div>
1	B	653	<div> <div>7%</div> <div>84%</div> <div>14%</div> <div>.</div> </div>
1	C	653	<div> <div>84%</div> <div>15%</div> <div>.</div> </div>
1	D	653	<div> <div>79%</div> <div>19%</div> <div>.</div> </div>
2	E	179	<div> <div>%</div> <div>81%</div> <div>10%</div> <div>9%</div> </div>
2	F	179	<div> <div>77%</div> <div>18%</div> <div>5%</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	G	179	
2	H	179	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	SF4	F	1002	-	-	-	X
3	SF4	G	1001	-	-	-	X
3	SF4	G	1002	-	-	-	X
3	SF4	H	1001	-	-	-	X
7	UNL	C	705	-	-	-	X

2 Entry composition

There are 9 unique types of molecules in this entry. The entry contains 27094 atoms, of which 168 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Benzoyl-CoA reductase, putative.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	652	Total	C	N	O	S	0	0	0
			5185	3311	875	965	34			
1	B	652	Total	C	N	O	S	0	1	0
			5181	3309	877	961	34			
1	C	653	Total	C	N	O	S	0	1	0
			5195	3317	876	968	34			
1	D	652	Total	C	N	O	S	0	1	0
			5193	3316	878	965	34			

- Molecule 2 is a protein called Iron-sulfur cluster-binding oxidoreductase, putative benzoyl-CoA reductase electron transfer protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	E	163	Total	C	N	O	S	0	1	0
			1241	772	215	240	14			
2	F	170	Total	C	N	O	S	0	1	0
			1325	820	227	264	14			
2	G	169	Total	C	N	O	S	0	2	0
			1322	819	229	260	14			
2	H	161	Total	C	N	O	S	0	0	0
			1221	758	213	236	14			

- Molecule 3 is IRON/SULFUR CLUSTER (three-letter code: SF4) (formula: Fe₄S₄).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	Fe	S	0	0
			8	4	4		
3	B	1	Total	Fe	S	0	0
			8	4	4		
3	C	1	Total	Fe	S	0	0
			8	4	4		
3	D	1	Total	Fe	S	0	0
			8	4	4		
3	E	1	Total	Fe	S	0	0
			8	4	4		
3	E	1	Total	Fe	S	0	0
			8	4	4		
3	E	1	Total	Fe	S	0	0
			8	4	4		
3	F	1	Total	Fe	S	0	0
			8	4	4		
3	F	1	Total	Fe	S	0	0
			8	4	4		
3	F	1	Total	Fe	S	0	0
			8	4	4		
3	G	1	Total	Fe	S	0	0
			8	4	4		
3	G	1	Total	Fe	S	0	0
			8	4	4		
3	G	1	Total	Fe	S	0	0
			8	4	4		
3	H	1	Total	Fe	S	0	0
			8	4	4		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	H	1	Total	Fe	S	0	0
			8	4	4		
3	H	1	Total	Fe	S	0	0
			8	4	4		

- Molecule 4 is PHOSPHONIC ACIDMONO-(2-AMINO-5,6-DIMERCAPTO-4-OXO-3,7,8A, 9,10,10A-HEXAHYDRO-4H-8-OXA-1,3,9,10-TETRAAZA-ANTHRACEN-7-YLMETHYL) ESTER (three-letter code: MTE) (formula: $C_{10}H_{14}N_5O_6PS_2$).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
4	A	1	Total	C	N	O	P	S	0	0
			24	10	5	6	1	2		
4	A	1	Total	C	N	O	P	S	0	0
			24	10	5	6	1	2		
4	B	1	Total	C	N	O	P	S	0	0
			24	10	5	6	1	2		
4	B	1	Total	C	N	O	P	S	0	0
			24	10	5	6	1	2		
4	C	1	Total	C	N	O	P	S	0	0
			24	10	5	6	1	2		
4	C	1	Total	C	N	O	P	S	0	0
			24	10	5	6	1	2		
4	D	1	Total	C	N	O	P	S	0	0
			24	10	5	6	1	2		
4	D	1	Total	C	N	O	P	S	0	0
			24	10	5	6	1	2		

- Molecule 5 is TUNGSTEN ION (three-letter code: W) (formula: W).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	B	1	Total W 1 1	0	0
5	A	1	Total W 1 1	0	0
5	D	1	Total W 1 1	0	0
5	C	1	Total W 1 1	0	0


- Molecule 6 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	B	1	Total Mg 1 1	0	0
6	A	1	Total Mg 1 1	0	0
6	D	1	Total Mg 1 1	0	0
6	C	1	Total Mg 1 1	0	0

- Molecule 7 is UNKNOWN LIGAND (three-letter code: UNL) (formula:).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	B	1	Total X 1 1	0	0
7	A	1	Total X 1 1	0	0
7	D	1	Total X 1 1	0	0
7	C	1	Total X 1 1	0	0

- Molecule 8 is 1,5 Dienoyl-CoA (three-letter code: 4KX) (formula: C₂₈H₄₂N₇O₁₇P₃S).



Mol	Chain	Residues	Atoms							ZeroOcc	AltConf
8	A	1	Total	C	H	N	O	P	S	0	0
			98	28	42	7	17	3	1		
8	B	1	Total	C	H	N	O	P	S	0	0
			98	28	42	7	17	3	1		
8	C	1	Total	C	H	N	O	P	S	0	0
			98	28	42	7	17	3	1		
8	D	1	Total	C	H	N	O	P	S	0	0
			98	28	42	7	17	3	1		


- Molecule 9 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	134	Total	O	0	0
			134	134		
9	B	40	Total	O	0	0
			40	40		
9	C	96	Total	O	0	0
			96	96		
9	D	96	Total	O	0	0
			96	96		
9	E	46	Total	O	0	0
			46	46		
9	F	39	Total	O	0	0
			39	39		
9	G	29	Total	O	0	0
			29	29		
9	H	27	Total	O	0	0
			27	27		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Benzoyl-CoA reductase, putative




- Molecule 1: Benzoyl-CoA reductase, putative




- Molecule 1: Benzoyl-CoA reductase, putative





- Molecule 1: Benzoyl-CoA reductase, putative

Chain D: 79% 19% .




- Molecule 2: Iron-sulfur cluster-binding oxidoreductase, putative benzoyl-CoA reductase electron transfer protein

Chain E: 81% 10% 9%




- Molecule 2: Iron-sulfur cluster-binding oxidoreductase, putative benzoyl-CoA reductase electron transfer protein


Chain F: 77% 18% 5%



- Molecule 2: Iron-sulfur cluster-binding oxidoreductase, putative benzoyl-CoA reductase electron transfer protein



- Molecule 2: Iron-sulfur cluster-binding oxidoreductase, putative benzoyl-CoA reductase electron transfer protein



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	125.34Å 116.30Å 144.16Å 90.00° 110.48° 90.00°	Depositor
Resolution (Å)	82.63 – 2.21 88.13 – 2.21	Depositor EDS
% Data completeness (in resolution range)	92.6 (82.63-2.21) 92.6 (88.13-2.21)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.05 (at 2.20Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.9_1692)	Depositor
R, R_{free}	0.178 , 0.222 0.189 , 0.227	Depositor DCC
R_{free} test set	2340 reflections (1.32%)	DCC
Wilson B-factor (Å ²)	37.3	Xtriage
Anisotropy	0.231	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 42.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	27094	wwPDB-VP
Average B, all atoms (Å ²)	58.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.68% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MG, SF4, W, UNL, 4KX, MTE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.36	0/5311	0.52	3/7179 (0.0%)
1	B	0.30	0/5310	0.53	3/7179 (0.0%)
1	C	0.35	0/5324	0.50	0/7196
1	D	0.36	0/5322	0.53	0/7193
2	E	0.38	0/1266	0.54	0/1715
2	F	0.41	0/1351	0.55	0/1829
2	G	0.37	0/1351	0.52	0/1828
2	H	0.36	0/1242	0.56	0/1681
All	All	0.35	0/26477	0.53	6/35800 (0.0%)

There are no bond length outliers.

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	576	ARG	NE-CZ-NH2	14.50	127.55	120.30
1	B	576	ARG	NE-CZ-NH1	-14.30	113.15	120.30
1	A	576	ARG	NE-CZ-NH1	10.84	125.72	120.30
1	A	576	ARG	NE-CZ-NH2	-10.82	114.89	120.30
1	B	576	ARG	CD-NE-CZ	7.31	133.84	123.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5185	0	5119	72	0
1	B	5181	0	5113	70	0
1	C	5195	0	5128	67	0
1	D	5193	0	5132	88	0
2	E	1241	0	1180	11	0
2	F	1325	0	1276	24	0
2	G	1322	0	1278	38	0
2	H	1221	0	1158	25	0
3	A	8	0	0	0	0
3	B	8	0	0	1	0
3	C	8	0	0	0	0
3	D	8	0	0	1	0
3	E	24	0	0	1	0
3	F	24	0	0	1	0
3	G	24	0	0	1	0
3	H	24	0	0	1	0
4	A	48	0	20	2	0
4	B	48	0	20	2	0
4	C	48	0	20	2	0
4	D	48	0	20	4	0
5	A	1	0	0	0	0
5	B	1	0	0	0	0
5	C	1	0	0	0	0
5	D	1	0	0	0	0
6	A	1	0	0	0	0
6	B	1	0	0	0	0
6	C	1	0	0	0	0
6	D	1	0	0	0	0
7	A	1	0	0	0	0
7	B	1	0	0	0	0
7	C	1	0	0	0	0
7	D	1	0	0	0	0
8	A	56	42	38	1	0
8	B	56	42	38	1	0
8	C	56	42	38	2	0
8	D	56	42	38	2	0
9	A	134	0	0	1	0
9	B	40	0	0	4	0
9	C	96	0	0	1	0
9	D	96	0	0	5	0
9	E	46	0	0	0	0
9	F	39	0	0	3	0
9	G	29	0	0	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
9	H	27	0	0	2	0
All	All	26926	168	25616	375	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 375 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:114:ARG:HG3	1:A:114:ARG:HH11	1.21	1.04
1:B:114:ARG:HG3	1:B:114:ARG:HH11	1.21	1.00
1:B:434:LEU:HD11	1:B:458:THR:HG23	1.54	0.89
1:A:434:LEU:HD11	1:A:458:THR:HG23	1.53	0.88
2:F:43:ASN:HB3	2:F:46:ARG:HG3	1.56	0.86

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	650/653 (100%)	626 (96%)	24 (4%)	0	100	100
1	B	651/653 (100%)	626 (96%)	25 (4%)	0	100	100
1	C	652/653 (100%)	632 (97%)	19 (3%)	1 (0%)	51	58
1	D	651/653 (100%)	625 (96%)	26 (4%)	0	100	100
2	E	160/179 (89%)	158 (99%)	1 (1%)	1 (1%)	28	29
2	F	169/179 (94%)	166 (98%)	3 (2%)	0	100	100
2	G	169/179 (94%)	163 (96%)	6 (4%)	0	100	100
2	H	157/179 (88%)	151 (96%)	5 (3%)	1 (1%)	28	29
All	All	3259/3328 (98%)	3147 (97%)	109 (3%)	3 (0%)	55	63

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	622	ASN
2	H	43	ASN
2	E	43	ASN

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	548/548 (100%)	532 (97%)	16 (3%)	48	60
1	B	546/548 (100%)	531 (97%)	15 (3%)	50	62
1	C	549/548 (100%)	534 (97%)	15 (3%)	50	62
1	D	549/548 (100%)	528 (96%)	21 (4%)	38	47
2	E	135/159 (85%)	133 (98%)	2 (2%)	70	82
2	F	150/159 (94%)	147 (98%)	3 (2%)	60	74
2	G	149/159 (94%)	145 (97%)	4 (3%)	50	62
2	H	132/159 (83%)	128 (97%)	4 (3%)	46	58
All	All	2758/2828 (98%)	2678 (97%)	80 (3%)	48	60

5 of 80 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	319	MET
1	C	599	HIS
2	G	166	ASP
1	C	321	LYS
1	C	551	TYR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 20 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	222	ASN
1	C	263	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	E	34	HIS
1	B	263	ASN
1	B	599	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 40 ligands modelled in this entry, 8 are monoatomic and 4 are unknown - leaving 28 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	SF4	A	701	1	0,12,12	0.00	-	0,24,24	0.00	-
4	MTE	A	702	5,6	21,26,26	2.68	10 (47%)	19,40,40	2.23	5 (26%)
4	MTE	A	703	5,6	21,26,26	2.74	10 (47%)	19,40,40	2.19	6 (31%)
8	4KX	A	707	-	50,59,59	0.46	0	55,87,87	0.72	2 (3%)
3	SF4	B	701	1	0,12,12	0.00	-	0,24,24	0.00	-
4	MTE	B	702	5,6	21,26,26	2.94	11 (52%)	19,40,40	1.94	4 (21%)
4	MTE	B	703	5,6	21,26,26	2.78	9 (42%)	19,40,40	2.15	4 (21%)
8	4KX	B	707	-	50,59,59	0.52	1 (2%)	55,87,87	0.73	2 (3%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	SF4	C	701	1	0,12,12	0.00	-	0,24,24	0.00	-
4	MTE	C	702	5,6	21,26,26	2.86	10 (47%)	19,40,40	1.93	4 (21%)
4	MTE	C	703	5,6	21,26,26	2.80	9 (42%)	19,40,40	2.06	2 (10%)
8	4KX	C	707	-	50,59,59	0.53	1 (2%)	55,87,87	0.90	2 (3%)
3	SF4	D	702	1	0,12,12	0.00	-	0,24,24	0.00	-
4	MTE	D	703	5,6	21,26,26	2.71	10 (47%)	19,40,40	2.73	6 (31%)
4	MTE	D	704	5,6	21,26,26	2.73	10 (47%)	19,40,40	1.91	2 (10%)
8	4KX	D	707	-	50,59,59	0.44	0	55,87,87	1.00	1 (1%)
3	SF4	E	1001	2	0,12,12	0.00	-	0,24,24	0.00	-
3	SF4	E	1002	2	0,12,12	0.00	-	0,24,24	0.00	-
3	SF4	E	1003	2	0,12,12	0.00	-	0,24,24	0.00	-
3	SF4	F	1001	2	0,12,12	0.00	-	0,24,24	0.00	-
3	SF4	F	1002	2	0,12,12	0.00	-	0,24,24	0.00	-
3	SF4	F	1003	2	0,12,12	0.00	-	0,24,24	0.00	-
3	SF4	G	1001	2	0,12,12	0.00	-	0,24,24	0.00	-
3	SF4	G	1002	2	0,12,12	0.00	-	0,24,24	0.00	-
3	SF4	G	1003	2	0,12,12	0.00	-	0,24,24	0.00	-
3	SF4	H	1001	2	0,12,12	0.00	-	0,24,24	0.00	-
3	SF4	H	1002	2	0,12,12	0.00	-	0,24,24	0.00	-
3	SF4	H	1003	2	0,12,12	0.00	-	0,24,24	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	SF4	A	701	1	-	0/0/48/48	0/6/5/5
4	MTE	A	702	5,6	-	0/6/34/34	0/3/3/3
4	MTE	A	703	5,6	-	0/6/34/34	0/3/3/3
8	4KX	A	707	-	-	0/48/79/79	0/4/4/4
3	SF4	B	701	1	-	0/0/48/48	0/6/5/5
4	MTE	B	702	5,6	-	0/6/34/34	0/3/3/3
4	MTE	B	703	5,6	-	0/6/34/34	0/3/3/3
8	4KX	B	707	-	-	0/48/79/79	0/4/4/4
3	SF4	C	701	1	-	0/0/48/48	0/6/5/5
4	MTE	C	702	5,6	-	0/6/34/34	0/3/3/3
4	MTE	C	703	5,6	-	0/6/34/34	0/3/3/3
8	4KX	C	707	-	-	0/48/79/79	0/4/4/4
3	SF4	D	702	1	-	0/0/48/48	0/6/5/5
4	MTE	D	703	5,6	-	0/6/34/34	0/3/3/3

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	MTE	D	704	5,6	-	0/6/34/34	0/3/3/3
8	4KX	D	707	-	-	0/48/79/79	0/4/4/4
3	SF4	E	1001	2	-	0/0/48/48	0/6/5/5
3	SF4	E	1002	2	-	0/0/48/48	0/6/5/5
3	SF4	E	1003	2	-	0/0/48/48	0/6/5/5
3	SF4	F	1001	2	-	0/0/48/48	0/6/5/5
3	SF4	F	1002	2	-	0/0/48/48	0/6/5/5
3	SF4	F	1003	2	-	0/0/48/48	0/6/5/5
3	SF4	G	1001	2	-	0/0/48/48	0/6/5/5
3	SF4	G	1002	2	-	0/0/48/48	0/6/5/5
3	SF4	G	1003	2	-	0/0/48/48	0/6/5/5
3	SF4	H	1001	2	-	0/0/48/48	0/6/5/5
3	SF4	H	1002	2	-	0/0/48/48	0/6/5/5
3	SF4	H	1003	2	-	0/0/48/48	0/6/5/5

The worst 5 of 81 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	703	MTE	C9-C10	-6.49	1.29	1.41
4	C	702	MTE	C9-C10	-6.32	1.29	1.41
4	C	703	MTE	C9-C10	-6.01	1.30	1.41
4	B	702	MTE	C9-C10	-5.96	1.30	1.41
4	D	704	MTE	C9-C10	-5.78	1.30	1.41

The worst 5 of 40 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	D	703	MTE	O3'-C7-C6	-7.89	103.70	108.96
4	A	703	MTE	O3'-C7-C6	-2.82	107.08	108.96
4	D	703	MTE	C10-N8-C7	-2.70	118.38	123.67
4	A	702	MTE	C10-N8-C7	-2.44	118.89	123.67
4	C	702	MTE	C10-N8-C7	-2.34	119.08	123.67

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

15 monomers are involved in 22 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	702	MTE	2	0
8	A	707	4KX	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	701	SF4	1	0
4	B	702	MTE	2	0
8	B	707	4KX	1	0
4	C	703	MTE	2	0
8	C	707	4KX	2	0
3	D	702	SF4	1	0
4	D	703	MTE	3	0
4	D	704	MTE	1	0
8	D	707	4KX	2	0
3	E	1001	SF4	1	0
3	F	1002	SF4	1	0
3	G	1002	SF4	1	0
3	H	1002	SF4	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	652/653 (99%)	-0.19	1 (0%) 94 94	28, 47, 79, 111	0
1	B	652/653 (99%)	0.45	46 (7%) 17 16	37, 83, 121, 146	0
1	C	653/653 (100%)	-0.26	3 (0%) 90 90	30, 49, 83, 119	0
1	D	652/653 (99%)	-0.26	3 (0%) 90 90	32, 54, 82, 105	0
2	E	163/179 (91%)	-0.27	1 (0%) 89 88	30, 45, 70, 117	0
2	F	170/179 (94%)	-0.29	0 100 100	30, 42, 71, 100	0
2	G	169/179 (94%)	-0.22	0 100 100	30, 48, 99, 133	0
2	H	161/179 (89%)	-0.32	1 (0%) 89 88	34, 50, 100, 129	0
All	All	3272/3328 (98%)	-0.11	55 (1%) 70 68	28, 53, 101, 146	0

The worst 5 of 55 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	394	LYS	6.0
1	B	377	PHE	5.0
1	B	366	LEU	4.9
1	B	259	TRP	4.7
1	B	285	TRP	4.5

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
7	UNL	C	705	1/-	0.97	0.43	15.12	46,46,46,46	0
3	SF4	F	1002	8/8	0.99	0.14	2.33	25,29,30,30	0
3	SF4	H	1001	8/8	0.99	0.14	2.14	24,34,36,37	0
3	SF4	G	1002	8/8	0.99	0.15	2.12	27,39,43,43	0
3	SF4	G	1001	8/8	0.99	0.15	2.01	34,37,38,41	0
3	SF4	E	1002	8/8	0.99	0.15	2.00	35,38,41,44	0
3	SF4	H	1002	8/8	0.99	0.14	1.57	41,43,53,54	0
3	SF4	F	1001	8/8	0.99	0.13	1.16	26,32,33,34	0
3	SF4	E	1001	8/8	0.99	0.14	0.94	32,32,34,38	0
3	SF4	G	1003	8/8	0.99	0.14	0.75	37,40,42,48	0
3	SF4	H	1003	8/8	0.99	0.14	0.53	40,42,44,51	0
6	MG	C	704	1/1	0.97	0.14	0.48	37,37,37,37	0
3	SF4	E	1003	8/8	0.99	0.14	0.32	27,33,35,36	0
8	4KX	C	707	56/56	0.89	0.17	0.30	61,94,124,129	0
4	MTE	A	702	24/24	0.97	0.13	0.28	14,26,32,39	0
3	SF4	F	1003	8/8	0.99	0.13	0.11	26,32,34,39	0
6	MG	D	706	1/1	0.97	0.11	0.11	32,32,32,32	0
8	4KX	A	707	56/56	0.93	0.14	0.09	31,50,87,109	98
8	4KX	D	707	56/56	0.95	0.13	0.02	36,48,67,81	0
4	MTE	B	702	24/24	0.93	0.13	-0.01	42,58,66,190	0
4	MTE	C	702	24/24	0.98	0.12	-0.03	28,31,35,39	0
4	MTE	D	703	24/24	0.97	0.11	-0.25	19,31,35,39	0
8	4KX	B	707	56/56	0.85	0.18	-0.36	69,105,135,149	98
3	SF4	C	701	8/8	0.99	0.11	-0.39	32,38,40,48	0
4	MTE	A	703	24/24	0.97	0.11	-0.50	18,25,34,35	0
4	MTE	D	704	24/24	0.98	0.10	-0.64	23,33,38,47	0
4	MTE	C	703	24/24	0.96	0.10	-0.79	33,39,42,43	0
4	MTE	B	703	24/24	0.95	0.11	-0.98	46,50,64,67	0
3	SF4	B	701	8/8	0.99	0.09	-1.00	45,50,55,62	0
3	SF4	D	702	8/8	0.98	0.10	-1.04	26,32,36,45	0
6	MG	A	705	1/1	0.97	0.10	-1.62	30,30,30,30	0
3	SF4	A	701	8/8	0.98	0.10	-2.90	23,27,32,38	0
7	UNL	D	701	1/-	0.91	0.36	-	36,36,36,36	0
5	W	C	706	1/1	0.99	0.17	-	43,43,43,43	0
6	MG	B	705	1/1	0.86	0.06	-	47,47,47,47	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
5	W	B	704	1/1	0.87	0.15	-	73,73,73,73	0
5	W	D	705	1/1	0.99	0.17	-	36,36,36,36	0
7	UNL	A	706	1/-	0.94	0.32	-	31,31,31,31	0
5	W	A	704	1/1	0.99	0.18	-	33,33,33,33	0
7	UNL	B	706	1/-	0.97	0.29	-	21,21,21,21	0

6.5 Other polymers [i](#)

There are no such residues in this entry.