



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 22, 2017 – 05:39 PM EDT

PDB ID : 4Z45
Title : Structure of OBP3 from the currant-lettuce aphid *Nasonovia ribisnigri*
Authors : Northey, T.; Venthur, H.; De Biasio, F.; Chauviac, F.-X.; Cole, A.R.; Field, L.M.; Zhou, J.-J.; Keep, N.H.
Deposited on : unknown
Resolution : 2.02 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.9-1692
EDS : rb-20029824
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20029824


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.02 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	8396 (2.04-2.00)
Clashscore	112137	9678 (2.04-2.00)
Ramachandran outliers	110173	9566 (2.04-2.00)
Sidechain outliers	110143	9565 (2.04-2.00)
RSRZ outliers	101464	8490 (2.04-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	121	<div> <div style="width: 5%; background-color: red;"></div> <div style="width: 88%; background-color: green;"></div> <div style="width: 8%; background-color: yellow;"></div> <div style="width: 7%; background-color: orange;"></div> <div style="width: 2%; background-color: grey;"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> 5% 88% 8% .. </div> </div>
1	B	121	<div> <div style="width: 5%; background-color: red;"></div> <div style="width: 84%; background-color: green;"></div> <div style="width: 11%; background-color: yellow;"></div> <div style="width: 2%; background-color: orange;"></div> <div style="width: 8%; background-color: grey;"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> 5% 84% 11% .. </div> </div>
1	D	121	<div> <div style="width: 5%; background-color: red;"></div> <div style="width: 86%; background-color: green;"></div> <div style="width: 10%; background-color: yellow;"></div> <div style="width: 2%; background-color: orange;"></div> <div style="width: 7%; background-color: grey;"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> 5% 86% 10% .. </div> </div>

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 3190 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Odorant-binding protein NribOBP3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	118	Total	C	N	O	S	0	4	0
			940	591	145	192	12			
1	B	117	Total	C	N	O	S	0	5	0
			943	597	144	190	12			
1	D	118	Total	C	N	O	S	0	4	0
			941	594	145	189	13			

There are 36 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	ALA	-	expression tag	UNP C5J8G4
A	2	SER	-	expression tag	UNP C5J8G4
A	3	MET	-	expression tag	UNP C5J8G4
A	102	ASN	SER	conflict	UNP C5J8G4
A	103	TYR	LYS	conflict	UNP C5J8G4
A	105	TYR	HIS	conflict	UNP C5J8G4
A	106	THR	ASP	conflict	UNP C5J8G4
A	107	VAL	ARG	conflict	UNP C5J8G4
A	108	MET	LYS	conflict	UNP C5J8G4
A	113	LYS	GLN	conflict	UNP C5J8G4
A	114	GLN	ASP	conflict	UNP C5J8G4
A	115	LEU	PRO	conflict	UNP C5J8G4
B	1	ALA	-	expression tag	UNP C5J8G4
B	2	SER	-	expression tag	UNP C5J8G4
B	3	MET	-	expression tag	UNP C5J8G4
B	102	ASN	SER	conflict	UNP C5J8G4
B	103	TYR	LYS	conflict	UNP C5J8G4
B	105	TYR	HIS	conflict	UNP C5J8G4
B	106	THR	ASP	conflict	UNP C5J8G4
B	107	VAL	ARG	conflict	UNP C5J8G4
B	108	MET	LYS	conflict	UNP C5J8G4
B	113	LYS	GLN	conflict	UNP C5J8G4
B	114	GLN	ASP	conflict	UNP C5J8G4

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	115	LEU	PRO	conflict	UNP C5J8G4
D	1	ALA	-	expression tag	UNP C5J8G4
D	2	SER	-	expression tag	UNP C5J8G4
D	3	MET	-	expression tag	UNP C5J8G4
D	102	ASN	SER	conflict	UNP C5J8G4
D	103	TYR	LYS	conflict	UNP C5J8G4
D	105	TYR	HIS	conflict	UNP C5J8G4
D	106	THR	ASP	conflict	UNP C5J8G4
D	107	VAL	ARG	conflict	UNP C5J8G4
D	108	MET	LYS	conflict	UNP C5J8G4
D	113	LYS	GLN	conflict	UNP C5J8G4
D	114	GLN	ASP	conflict	UNP C5J8G4
D	115	LEU	PRO	conflict	UNP C5J8G4


- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	150	Total O 150 150	0	0
2	B	126	Total O 126 126	0	0
2	D	90	Total O 90 90	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: Odorant-binding protein NribOBP3



- Molecule 1: Odorant-binding protein NribOBP3



- Molecule 1: Odorant-binding protein NribOBP3



4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α , β , γ	87.25Å 87.25Å 95.08Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	59.16 – 2.02 59.16 – 2.02	Depositor EDS
% Data completeness (in resolution range)	100.0 (59.16-2.02) 100.0 (59.16-2.02)	Depositor EDS
R_{merge}	0.14	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.38 (at 2.02Å)	Xtriage
Refinement program	REFMAC 5.8.0107	Depositor
R, R_{free}	0.164 , 0.207 0.172 , 0.211	Depositor DCC
R_{free} test set	1403 reflections (5.29%)	DCC
Wilson B-factor (Å ²)	20.9	Xtriage
Anisotropy	0.256	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 46.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.036 for -h,-k,l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3190	wwPDB-VP
Average B, all atoms (Å ²)	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.01% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	1.30	5/965 (0.5%)	1.09	5/1300 (0.4%)
1	B	1.30	7/972 (0.7%)	1.07	6/1309 (0.5%)
1	D	1.30	4/966 (0.4%)	1.11	7/1301 (0.5%)
All	All	1.30	16/2903 (0.6%)	1.09	18/3910 (0.5%)

All (16) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	8[A]	GLU	CD-OE2	8.30	1.34	1.25
1	B	8[B]	GLU	CD-OE2	8.30	1.34	1.25
1	D	105	TYR	CE1-CZ	8.07	1.49	1.38
1	A	8	GLU	CD-OE2	7.93	1.34	1.25
1	B	21[A]	GLU	CG-CD	7.02	1.62	1.51
1	B	21[B]	GLU	CG-CD	7.02	1.62	1.51
1	D	21	GLU	CD-OE1	6.96	1.33	1.25
1	B	64	GLU	CD-OE2	6.76	1.33	1.25
1	A	70	TYR	CE1-CZ	-6.68	1.29	1.38
1	A	8	GLU	CG-CD	6.39	1.61	1.51
1	D	105	TYR	CZ-OH	6.01	1.48	1.37
1	D	39	LYS	N-CA	5.61	1.57	1.46
1	A	87	TYR	CB-CG	-5.52	1.43	1.51
1	A	29	SER	CA-CB	5.13	1.60	1.52
1	B	64	GLU	CG-CD	5.02	1.59	1.51
1	B	80	GLU	CG-CD	-5.00	1.44	1.51

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	40	CYS	CA-CB-SG	-7.83	99.91	114.00
1	B	21[A]	GLU	OE1-CD-OE2	-7.49	114.31	123.30
1	B	21[B]	GLU	OE1-CD-OE2	-7.49	114.31	123.30
1	D	39	LYS	CD-CE-NZ	7.02	127.86	111.70
1	D	38	GLY	N-CA-C	6.85	130.23	113.10

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	54	ASP	CB-CG-OD1	6.61	124.25	118.30
1	A	101	CYS	CA-CB-SG	-6.56	102.19	114.00
1	B	4	ARG	NE-CZ-NH2	-6.38	117.11	120.30
1	A	4	ARG	CG-CD-NE	-6.36	98.44	111.80
1	D	45[A]	MET	CG-SD-CE	6.25	110.20	100.20
1	D	45[B]	MET	CG-SD-CE	6.25	110.20	100.20
1	B	87	TYR	CB-CG-CD1	-5.86	117.48	121.00
1	D	4	ARG	CG-CD-NE	-5.81	99.59	111.80
1	B	54	ASP	CB-CG-OD2	-5.57	113.28	118.30
1	B	4	ARG	CG-CD-NE	-5.41	100.44	111.80
1	A	55	ASP	CB-CG-OD1	5.41	123.17	118.30
1	A	23	ASP	CB-CG-OD1	5.33	123.10	118.30
1	D	55	ASP	CB-CG-OD1	5.06	122.85	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	940	0	933	3	0
1	B	943	0	938	7	0
1	D	941	0	942	15	0
2	A	150	0	0	0	0
2	B	126	0	0	0	0
2	D	90	0	0	2	0
All	All	3190	0	2813	23	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (23) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:27:VAL:HG21	1:D:41[B]:LEU:HD23	1.56	0.85

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:33:PRO:O	1:D:39:LYS:HG2	1.81	0.80
1:D:34:SER:HA	1:D:39:LYS:HE3	1.77	0.66
1:D:33:PRO:O	1:D:39:LYS:CG	2.47	0.62
1:D:41[B]:LEU:HD11	1:D:45[B]:MET:SD	2.40	0.61
1:D:33:PRO:O	1:D:39:LYS:HE2	2.03	0.58
1:D:13:TYR:HB3	1:D:45[B]:MET:HG2	1.87	0.56
1:D:41[B]:LEU:CD1	1:D:45[B]:MET:SD	2.95	0.55
1:A:20[A]:SER:OG	1:A:22:ASP:OD1	2.16	0.51
1:D:14:GLY:HA2	1:D:41[B]:LEU:HD11	1.93	0.50
1:B:13:TYR:CZ	1:B:49:LEU:HD21	2.46	0.50
1:D:33:PRO:O	1:D:39:LYS:CE	2.64	0.46
1:D:27:VAL:CG2	1:D:41[B]:LEU:HD23	2.38	0.46
1:D:41[B]:LEU:HD12	1:D:45[B]:MET:HG3	1.99	0.45
1:D:27:VAL:HG13	2:D:250:HOH:O	2.18	0.43
1:B:60:LYS:HB2	1:B:87:TYR:CD1	2.54	0.42
1:D:39:LYS:NZ	2:D:201:HOH:O	2.06	0.42
1:A:87:TYR:CD1	1:B:8[B]:GLU:OE1	2.74	0.41
1:B:16:ALA:HB1	1:B:48:LYS:HE2	2.03	0.41
1:A:87:TYR:HH	1:B:12:TYR:HD1	1.66	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	120/121 (99%)	118 (98%)	2 (2%)	0	100	100
1	B	120/121 (99%)	119 (99%)	1 (1%)	0	100	100
1	D	120/121 (99%)	117 (98%)	3 (2%)	0	100	100
All	All	360/363 (99%)	354 (98%)	6 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	109/108 (101%)	109 (100%)	0	100	100
1	B	109/108 (101%)	108 (99%)	1 (1%)	82	86
1	D	109/108 (101%)	109 (100%)	0	100	100
All	All	327/324 (101%)	326 (100%)	1 (0%)	94	95

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	77	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	D	18	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	118/121 (97%)	-0.60	1 (0%) 86 86	11, 17, 30, 68	0
1	B	117/121 (96%)	-0.50	1 (0%) 84 84	13, 24, 40, 63	0
1	D	118/121 (97%)	-0.18	6 (5%) 29 29	14, 25, 57, 66	3 (2%)
All	All	353/363 (97%)	-0.42	8 (2%) 61 61	11, 22, 50, 68	3 (0%)

All (8) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	34	SER	3.0
1	D	30	TYR	2.9
1	D	27	VAL	2.6
1	D	37	THR	2.3
1	D	1	ALA	2.2
1	B	1	ALA	2.2
1	D	33	PRO	2.1
1	A	1	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers

There are no such residues in this entry.