



Full wwPDB NMR Structure Validation Report ⓘ

Feb 12, 2017 – 08:53 pm GMT

PDB ID : 1Z6H
Title : Solution Structure of Bacillus subtilis BLAP biotinylated-form
Authors : Cui, G.; Xia, B.
Deposited on : 2005-03-22

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

Cyrange : Kirchner and Güntert (2011)
NmrClust : Kelley et al. (1996)
MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
ShiftChecker : trunk28760
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment is 95%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	125131	11601
Ramachandran outliers	121729	10391
Sidechain outliers	121581	10367

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	72	 86% 6% 8%

2 Ensemble composition and analysis

This entry contains 20 models. Model 7 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *lowest energy*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:3-A:68 (66)	0.11	7

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 3 clusters and 4 single-model clusters were found.

Cluster number	Models
1	1, 2, 4, 6, 7, 8, 12, 13, 17, 18, 19
2	3, 5, 16
3	9, 11
Single-model clusters	10; 14; 15; 20


3 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 1153 atoms, of which 585 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Biotin/Lipoyl Attachment Protein.

Mol	Chain	Residues	Atoms						Trace
1	A	72	Total	C	H	N	O	S	0
			1123	346	570	93	111	3	

- Molecule 2 is 5-(HEXAHYDRO-2-OXO-1H-THIENO[3,4-D]IMIDAZOL-6-YL)PENTANAL (three-letter code: BTI) (formula: $C_{10}H_{16}N_2O_2S$).




Mol	Chain	Residues	Atoms					
2	A	1	Total	C	H	N	O	S
			30	10	15	2	2	1

4 Residue-property plots

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: Biotin/Lipoyl Attachment Protein




4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.


4.2.1 Score per residue for model 1

- Molecule 1: Biotin/Lipoyl Attachment Protein




4.2.2 Score per residue for model 2

- Molecule 1: Biotin/Lipoyl Attachment Protein




4.2.3 Score per residue for model 3

- Molecule 1: Biotin/Lipoyl Attachment Protein




4.2.4 Score per residue for model 4

- Molecule 1: Biotin/Lipoyl Attachment Protein




4.2.5 Score per residue for model 5

- Molecule 1: Biotin/Lipoyl Attachment Protein




4.2.6 Score per residue for model 6

- Molecule 1: Biotin/Lipoyl Attachment Protein




4.2.7 Score per residue for model 7 (medoid)

- Molecule 1: Biotin/Lipoyl Attachment Protein




4.2.8 Score per residue for model 8

- Molecule 1: Biotin/Lipoyl Attachment Protein




4.2.9 Score per residue for model 9

- Molecule 1: Biotin/Lipoyl Attachment Protein




4.2.10 Score per residue for model 10

- Molecule 1: Biotin/Lipoyl Attachment Protein




4.2.11 Score per residue for model 11

- Molecule 1: Biotin/Lipoyl Attachment Protein




4.2.12 Score per residue for model 12

- Molecule 1: Biotin/Lipoyl Attachment Protein




4.2.13 Score per residue for model 13

- Molecule 1: Biotin/Lipoyl Attachment Protein




4.2.14 Score per residue for model 14

- Molecule 1: Biotin/Lipoyl Attachment Protein




4.2.15 Score per residue for model 15

- Molecule 1: Biotin/Lipoyl Attachment Protein




4.2.16 Score per residue for model 16

- Molecule 1: Biotin/Lipoyl Attachment Protein




4.2.17 Score per residue for model 17

- Molecule 1: Biotin/Lipoyl Attachment Protein




4.2.18 Score per residue for model 18

- Molecule 1: Biotin/Lipoyl Attachment Protein




4.2.19 Score per residue for model 19

- Molecule 1: Biotin/Lipoyl Attachment Protein



4.2.20 Score per residue for model 20

- Molecule 1: Biotin/Lipoyl Attachment Protein



5 Refinement protocol and experimental data overview

The models were refined using the following method: *simulated annealing, molecular dynamics*.

Of the 200 calculated structures, 20 were deposited, based on the following criterion: *structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CYANA	structure solution	1.0.6
AMBER	refinement	7.0

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	BMRB entry 6599
Number of chemical shift lists	2
Total number of shifts	935
Number of shifts mapped to atoms	920
Number of unparsed shifts	0
Number of shifts with mapping errors	15
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	95%

No validations of the models with respect to experimental NMR restraints is performed at this time.

6 Model quality [i](#)

6.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: BTI

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	0.53±0.00	0±0/514 (0.0±0.0%)	0.94±0.02	0±0/689 (0.0±0.0%)
All	All	0.53	0/10280 (0.0%)	0.94	2/13780 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	A	0.0±0.0	1.0±0.0
All	All	0	20

There are no bond-length outliers.

All unique angle outliers are listed below.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	44	ARG	NE-CZ-NH2	-7.49	116.56	120.30	2	2

There are no chirality outliers.

All unique planar outliers are listed below.

Mol	Chain	Res	Type	Group	Models (Total)
1	A	31	LEU	Peptide	20

6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	509	530	530	0±0
All	All	10480	10900	10900	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 0.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:31:LEU:HG	1:A:40:ILE:HD11	0.41	1.93	16	1

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	66/72 (92%)	62±1 (94±1%)	4±1 (6±1%)	0±0 (0±0%)	54	84
All	All	1320/1440 (92%)	1241 (94%)	77 (6%)	2 (0%)	54	84

All 2 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	12	TRP	1
1	A	35	LYS	1

6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	56/62 (90%)	52±1 (93±3%)	4±1 (7±3%)	22 68
All	All	1120/1240 (90%)	1040 (93%)	80 (7%)	22 68

All 12 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	32	GLU	20
1	A	60	ASN	20
1	A	44	ARG	9
1	A	6	GLN	9
1	A	26	GLN	8
1	A	54	LYS	4
1	A	37	GLU	4
1	A	17	LYS	2
1	A	34	MET	1
1	A	59	VAL	1
1	A	7	MET	1
1	A	35	LYS	1

6.3.3 RNA ⓘ

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.6 Ligand geometry ⓘ

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds for which Mogul statistics could be retrieved, the number of bonds that are observed in the model and the number of bonds that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length is the number of standard deviations the observed value is removed from the expected value. A bond length with $|Z| > 2$ is

considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond lengths.

Mol	Type	Chain	Res	Link	Bond lengths		
					Counts	RMSZ	#Z>2
2	BTI	A	135	1	16,16,16	0.67±0.02	0±0 (0±0%)

In the following table, the Counts columns list the number of angles for which Mogul statistics could be retrieved, the number of angles that are observed in the model and the number of angles that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond angle is the number of standard deviations the observed value is removed from the expected value. A bond angle with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond angles.

Mol	Type	Chain	Res	Link	Bond angles		
					Counts	RMSZ	#Z>2
2	BTI	A	135	1	21,21,21	2.00±0.02	1±0 (4±0%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BTI	A	135	1	-	0±0,5,27,27	0±0,2,2,2

There are no bond-length outliers.

All unique angle outliers are listed below.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
2	A	135	BTI	N2-C3-N3	5.51	104.61	108.85	16	20

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

6.7 Other polymers ⓘ

There are no such molecules in this entry.

6.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

7 Chemical shift validation [i](#)

The completeness of assignment taking into account all chemical shift lists is 95% for the well-defined parts and 94% for the entire structure.

7.1 Chemical shift list 1

File name: BMRB entry 6599

Chemical shift list name: *assigned_chem_shift_list_1*

7.1.1 Bookkeeping [i](#)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	920
Number of shifts mapped to atoms	920
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	1

7.1.2 Chemical shift referencing [i](#)

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction \pm precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	71	0.04 ± 0.24	None needed (< 0.5 ppm)
$^{13}\text{C}_\beta$	65	-0.39 ± 0.22	None needed (< 0.5 ppm)
$^{13}\text{C}'$	68	0.29 ± 0.18	None needed (< 0.5 ppm)
^{15}N	68	-0.23 ± 0.57	None needed (< 0.5 ppm)

7.1.3 Completeness of resonance assignments [i](#)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 95%, i.e. 769 atoms were assigned a chemical shift out of a possible 806. 14 out of 14 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^1H	^{13}C	^{15}N
Backbone	325/328 (99%)	130/131 (99%)	131/132 (99%)	64/65 (98%)
Sidechain	417/450 (93%)	255/260 (98%)	156/174 (90%)	6/16 (38%)

Continued on next page...

Continued from previous page...

	Total	¹ H	¹³ C	¹⁵ N
Aromatic	27/28 (96%)	15/15 (100%)	11/11 (100%)	1/2 (50%)
Overall	769/806 (95%)	400/406 (99%)	298/317 (94%)	71/83 (86%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 94%, i.e. 813 atoms were assigned a chemical shift out of a possible 867. 14 out of 14 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	¹ H	¹³ C	¹⁵ N
Backbone	346/358 (97%)	139/143 (97%)	139/144 (97%)	68/71 (96%)
Sidechain	440/481 (91%)	270/278 (97%)	162/185 (88%)	8/18 (44%)
Aromatic	27/28 (96%)	15/15 (100%)	11/11 (100%)	1/2 (50%)
Overall	813/867 (94%)	424/436 (97%)	312/340 (92%)	77/91 (85%)

7.1.4 Statistically unusual chemical shifts ⓘ


The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.

Mol	Chain	Res	Type	Atom	Shift, ppm	Expected range, ppm	Z-score
1	A	35	LYS	NZ	129.07	49.86 – 18.16	30.0

7.1.5 Random Coil Index (RCI) plots ⓘ

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain A:



7.2 Chemical shift list 2

File name: BMRB entry 6599

Chemical shift list name: *assigned_chem_shift_list_2*

7.2.1 Bookkeeping

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	15
Number of shifts mapped to atoms	0
Number of unparsed shifts	0
Number of shifts with mapping errors	15
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	0

The following assigned chemical shifts were not mapped to the molecules present in the coordinate file.

- Chain not found in structure. All 15 occurrences are reported below.

Chain	Res	Type	Atom	Shift Data		
				Value	Uncertainty	Ambiguity
UNMAPPED	1	BTI	H103	2.122	0.02	1
UNMAPPED	1	BTI	H72	1.172	0.02	1
UNMAPPED	1	BTI	H63	2.321	0.02	1
UNMAPPED	1	BTI	H82	1.013	0.02	1
UNMAPPED	1	BTI	H2	2.556	0.02	1
UNMAPPED	1	BTI	H92	1.555	0.02	1
UNMAPPED	1	BTI	H5	2.814	0.02	1
UNMAPPED	1	BTI	H73	1.46	0.02	1
UNMAPPED	1	BTI	HN2	6.093	0.02	1
UNMAPPED	1	BTI	H102	2.25	0.02	1
UNMAPPED	1	BTI	H4	2.572	0.02	1
UNMAPPED	1	BTI	H83	1.166	0.02	1
UNMAPPED	1	BTI	H62	2.002	0.02	1
UNMAPPED	1	BTI	HN3	6.717	0.02	1
UNMAPPED	1	BTI	H93	1.159	0.02	1

7.2.2 Chemical shift referencing

No chemical shift referencing corrections were calculated (not enough data).

7.2.3 Completeness of resonance assignments [i](#)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 0%, i.e. 0 atoms were assigned a chemical shift out of a possible 806. 0 out of 14 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	¹H	¹³C	¹⁵N
Backbone	0/328 (0%)	0/131 (0%)	0/132 (0%)	0/65 (0%)
Sidechain	0/450 (0%)	0/260 (0%)	0/174 (0%)	0/16 (0%)
Aromatic	0/28 (0%)	0/15 (0%)	0/11 (0%)	0/2 (0%)
Overall	0/806 (0%)	0/406 (0%)	0/317 (0%)	0/83 (0%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 0%, i.e. 0 atoms were assigned a chemical shift out of a possible 867. 0 out of 14 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	¹H	¹³C	¹⁵N
Backbone	0/358 (0%)	0/143 (0%)	0/144 (0%)	0/71 (0%)
Sidechain	0/481 (0%)	0/278 (0%)	0/185 (0%)	0/18 (0%)
Aromatic	0/28 (0%)	0/15 (0%)	0/11 (0%)	0/2 (0%)
Overall	0/867 (0%)	0/436 (0%)	0/340 (0%)	0/91 (0%)

7.2.4 Statistically unusual chemical shifts [i](#)

There are no statistically unusual chemical shifts.

7.2.5 Random Coil Index (RCI) plots [i](#)

No *random coil index* (RCI) plot could be generated from the current chemical shift list (assigned_chem_shift_list_2). RCI is only applicable to proteins.