



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 15, 2017 – 02:50 am GMT

PDB ID : 2Z6X
Title : Crystal structure of 22G, the wild-type protein of the photoswitchable GFP-like protein Dronpa
Authors : Kikuchi, A.; Jeyakanthan, J.; Taka, J.; Shiro, Y.; Mizuno, H.; Miyawaki, A.
Deposited on : 2007-08-09
Resolution : 2.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : trunk28620
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	4130 (2.30-2.30)
Clashscore	112137	4751 (2.30-2.30)
Ramachandran outliers	110173	4705 (2.30-2.30)
Sidechain outliers	110143	4704 (2.30-2.30)
RSRZ outliers	101464	4156 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	255	
1	B	255	
1	C	255	
1	D	255	
1	E	255	
1	F	255	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	G	255	
1	H	255	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
1	GYC	A	63	X	-	-	-
1	GYC	B	63	X	-	-	-
1	GYC	C	63	X	-	-	-
1	GYC	D	63	X	-	-	-
1	GYC	E	63	X	-	-	-
1	GYC	F	63	X	-	-	-
1	GYC	G	63	X	-	-	-
1	GYC	H	63	X	-	X	-

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 15008 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called photochromic protein Dronpa.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	217	Total	C	N	O	S	0	0	0
			1750	1122	295	323	10			
1	B	216	Total	C	N	O	S	0	0	0
			1743	1117	294	322	10			
1	C	215	Total	C	N	O	S	0	0	0
			1737	1114	293	320	10			
1	D	215	Total	C	N	O	S	0	0	0
			1737	1114	293	320	10			
1	E	216	Total	C	N	O	S	0	0	0
			1743	1117	294	322	10			
1	F	216	Total	C	N	O	S	0	0	0
			1743	1117	294	322	10			
1	G	215	Total	C	N	O	S	0	0	0
			1737	1114	293	320	10			
1	H	215	Total	C	N	O	S	0	0	0
			1737	1114	293	320	10			

- Molecule 2 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	113	Total	O	0	0
			113	113		
2	B	140	Total	O	0	0
			140	140		
2	C	157	Total	O	0	0
			157	157		
2	D	139	Total	O	0	0
			139	139		
2	E	148	Total	O	0	0
			148	148		
2	F	145	Total	O	0	0
			145	145		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	G	128	Total 128	O 128	0	0
2	H	111	Total 111	O 111	0	0


- Molecule 1: photochromic protein Dronpa




- Molecule 1: photochromic protein Dronpa




- Molecule 1: photochromic protein Dronpa




- Molecule 1: photochromic protein Dronpa





● Molecule 1: photochromic protein Dronpa



4 Data and refinement statistics

Property	Value	Source
Space group	P 43	Depositor
Cell constants a, b, c, α , β , γ	111.68Å 111.68Å 156.53Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.96 – 2.30 42.10 – 2.30	Depositor EDS
% Data completeness (in resolution range)	99.0 (19.96-2.30) 99.1 (42.10-2.30)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.98 (at 2.29Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.219 , 0.278 0.219 , 0.277	Depositor DCC
R_{free} test set	4269 reflections (5.08%)	DCC
Wilson B-factor (Å ²)	24.4	Xtriage
Anisotropy	0.013	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.27 , 9.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.41$, $\langle L^2 \rangle = 0.23$	Xtriage
Estimated twinning fraction	0.310 for h,-k,-l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	15008	wwPDB-VP
Average B, all atoms (Å ²)	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.80% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GYC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.40	0/1774	0.67	0/2394
1	B	0.40	0/1766	0.68	0/2382
1	C	0.43	0/1760	0.71	0/2374
1	D	0.43	0/1760	0.71	0/2374
1	E	0.42	0/1766	0.68	0/2382
1	F	0.45	0/1766	0.70	0/2382
1	G	0.41	0/1760	0.71	0/2374
1	H	0.40	0/1760	0.66	0/2374
All	All	0.42	0/14112	0.69	0/19036

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	1	0
1	B	1	0
1	C	1	0
1	D	1	0
1	E	1	0
1	F	1	0
1	G	1	0
1	H	1	0
All	All	8	0

There are no bond length outliers.

There are no bond angle outliers.

5 of 8 chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
1	A	63	GYC	CA1
1	B	63	GYC	CA1
1	C	63	GYC	CA1
1	D	63	GYC	CA1
1	E	63	GYC	CA1

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1750	0	1693	85	0
1	B	1743	0	1686	81	0
1	C	1737	0	1681	77	0
1	D	1737	0	1681	78	0
1	E	1743	0	1686	82	0
1	F	1743	0	1686	71	0
1	G	1737	0	1681	61	0
1	H	1737	0	1682	80	0
2	A	113	0	0	9	0
2	B	140	0	0	10	0
2	C	157	0	0	10	0
2	D	139	0	0	9	0
2	E	148	0	0	8	0
2	F	145	0	0	6	0
2	G	128	0	0	9	0
2	H	111	0	0	6	0
All	All	15008	0	13476	584	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 21.

The worst 5 of 584 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:H:63:GYC:HA31	1:H:63:GYC:HN2	1.15	1.08

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:138:LYS:HG2	1:C:162:LEU:HD21	1.38	1.05
1:C:187:ASP:HB3	2:C:371:HOH:O	1.59	1.01
1:H:198:LYS:HA	1:H:198:LYS:HE2	1.47	0.97
1:D:198:LYS:HB2	1:D:208:ASN:HD22	1.30	0.94

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	214/255 (84%)	199 (93%)	13 (6%)	2 (1%)	20	23
1	B	213/255 (84%)	196 (92%)	16 (8%)	1 (0%)	32	39
1	C	212/255 (83%)	198 (93%)	14 (7%)	0	100	100
1	D	212/255 (83%)	202 (95%)	10 (5%)	0	100	100
1	E	213/255 (84%)	203 (95%)	9 (4%)	1 (0%)	32	39
1	F	213/255 (84%)	204 (96%)	8 (4%)	1 (0%)	32	39
1	G	212/255 (83%)	202 (95%)	7 (3%)	3 (1%)	13	13
1	H	212/255 (83%)	195 (92%)	13 (6%)	4 (2%)	9	8
All	All	1701/2040 (83%)	1599 (94%)	90 (5%)	12 (1%)	25	30

5 of 12 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	E	167	GLY
1	H	202	LYS
1	A	200	HIS
1	G	218	GLY
1	H	167	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	186/217 (86%)	177 (95%)	9 (5%)	30	40
1	B	185/217 (85%)	173 (94%)	12 (6%)	20	26
1	C	184/217 (85%)	180 (98%)	4 (2%)	57	74
1	D	184/217 (85%)	175 (95%)	9 (5%)	29	39
1	E	185/217 (85%)	173 (94%)	12 (6%)	20	26
1	F	185/217 (85%)	179 (97%)	6 (3%)	44	60
1	G	184/217 (85%)	173 (94%)	11 (6%)	22	30
1	H	184/217 (85%)	179 (97%)	5 (3%)	50	67
All	All	1477/1736 (85%)	1409 (95%)	68 (5%)	31	42

5 of 68 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	100	ILE
1	E	94	ASN
1	G	211	GLU
1	D	145	LYS
1	E	58	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 51 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	205	ASN
1	E	94	ASN
1	H	158	ASN
1	D	216	HIS
1	E	158	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

8 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	GYC	A	63	1	22,22,23	4.44	11 (50%)	27,30,32	2.62	6 (22%)
1	GYC	B	63	1	22,22,23	4.33	11 (50%)	27,30,32	2.59	6 (22%)
1	GYC	C	63	1	22,22,23	4.37	11 (50%)	27,30,32	2.57	7 (25%)
1	GYC	D	63	1	22,22,23	4.40	10 (45%)	27,30,32	2.71	6 (22%)
1	GYC	E	63	1	22,22,23	4.60	9 (40%)	27,30,32	2.43	6 (22%)
1	GYC	F	63	1	22,22,23	4.45	13 (59%)	27,30,32	2.60	7 (25%)
1	GYC	G	63	1	22,22,23	4.52	11 (50%)	27,30,32	2.51	6 (22%)
1	GYC	H	63	1	22,22,23	4.51	10 (45%)	27,30,32	2.60	6 (22%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	GYC	A	63	1	1/1/5/7	0/9/29/30	0/2/2/2
1	GYC	B	63	1	1/1/5/7	0/9/29/30	0/2/2/2
1	GYC	C	63	1	1/1/5/7	0/9/29/30	0/2/2/2
1	GYC	D	63	1	1/1/5/7	0/9/29/30	0/2/2/2
1	GYC	E	63	1	1/1/5/7	0/9/29/30	0/2/2/2
1	GYC	F	63	1	1/1/5/7	0/9/29/30	0/2/2/2
1	GYC	G	63	1	1/1/5/7	0/9/29/30	0/2/2/2
1	GYC	H	63	1	1/1/5/7	0/9/29/30	0/2/2/2

The worst 5 of 86 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	E	63	GYC	CA1-C1	-17.52	1.26	1.51
1	H	63	GYC	CA1-C1	-16.69	1.28	1.51

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	G	63	GYC	CA1-C1	-16.54	1.28	1.51
1	D	63	GYC	CA1-C1	-16.48	1.28	1.51
1	A	63	GYC	CA1-C1	-16.04	1.28	1.51

The worst 5 of 50 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	63	GYC	CA3-N3-C2	-3.83	115.47	123.94
1	A	63	GYC	CA3-N3-C2	-3.80	115.53	123.94
1	G	63	GYC	CA3-N3-C2	-3.80	115.53	123.94
1	C	63	GYC	CA3-N3-C2	-3.74	115.65	123.94
1	F	63	GYC	CA3-N3-C2	-3.70	115.75	123.94

5 of 8 chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
1	D	63	GYC	CA1
1	G	63	GYC	CA1
1	A	63	GYC	CA1
1	C	63	GYC	CA1
1	F	63	GYC	CA1

There are no torsion outliers.

There are no ring outliers.

8 monomers are involved in 22 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	63	GYC	2	0
1	B	63	GYC	3	0
1	C	63	GYC	2	0
1	D	63	GYC	2	0
1	E	63	GYC	3	0
1	F	63	GYC	1	0
1	G	63	GYC	2	0
1	H	63	GYC	7	0

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	216/255 (84%)	-0.32	1 (0%) 90 93	15, 27, 40, 53	0
1	B	215/255 (84%)	-0.22	0 100 100	16, 29, 42, 58	0
1	C	214/255 (83%)	-0.54	1 (0%) 90 93	8, 19, 30, 41	0
1	D	214/255 (83%)	-0.44	1 (0%) 90 93	12, 21, 35, 50	0
1	E	215/255 (84%)	-0.41	1 (0%) 90 93	13, 24, 35, 56	0
1	F	215/255 (84%)	-0.46	1 (0%) 90 93	11, 22, 35, 50	0
1	G	214/255 (83%)	-0.43	1 (0%) 90 93	12, 23, 38, 51	0
1	H	214/255 (83%)	-0.30	1 (0%) 90 93	14, 29, 44, 56	0
All	All	1717/2040 (84%)	-0.39	7 (0%) 92 95	8, 24, 39, 58	0

The worst 5 of 7 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	2	SER	4.5
1	G	218	GLY	4.5
1	D	219	LEU	2.9
1	F	219	LEU	2.7
1	E	2	SER	2.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
1	GYC	D	63	21/22	0.94	0.12	-	16,21,25,28	0
1	GYC	G	63	21/22	0.91	0.13	-	26,27,29,35	0
1	GYC	A	63	21/22	0.91	0.13	-	26,34,35,46	0
1	GYC	C	63	21/22	0.94	0.12	-	15,19,22,25	0
1	GYC	F	63	21/22	0.92	0.13	-	19,25,29,32	0
1	GYC	H	63	21/22	0.91	0.13	-	23,30,33,34	0
1	GYC	B	63	21/22	0.93	0.11	-	26,32,33,39	0
1	GYC	E	63	21/22	0.95	0.11	-	23,26,28,29	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.