



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 14, 2017 – 04:46 pm GMT

PDB ID : 4ZGZ
Title : STRUCTURE OF HUMAN ANTIZYME INHIBITOR IN COMPLEX WITH
A C-TERMINAL FRAGMENT OF ANTIZYME
Authors : Chen, S.F.; Wu, H.Y.; Chan, N.L.
Deposited on : 2015-04-24
Resolution : 5.81 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.9-1692
EDS : trunk28620
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 5.81 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	1078 (7.94-3.70)
Clashscore	112137	1008 (7.82-3.80)
Ramachandran outliers	110173	1107 (7.94-3.70)
Sidechain outliers	110143	1079 (7.94-3.70)
RSRZ outliers	101464	1087 (7.94-3.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	438	<div> <div>85%</div> <div>11%</div> </div>
1	C	438	<div> <div>86%</div> <div>11%</div> </div>
2	B	128	<div> <div>70%</div> <div>27%</div> </div>
2	D	128	<div> <div>71%</div> <div>26%</div> </div>

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 7456 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Antizyme inhibitor 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	389	Total	C	N	O	S	1726	0	0
			2973	1919	470	561	23			
1	C	389	Total	C	N	O	S	1750	0	0
			2990	1927	479	562	22			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	MET	-	initiating methionine	UNP O14977
A	1	GLY	-	expression tag	UNP O14977
A	20	ASP	ASN	engineered mutation	UNP O14977
C	0	MET	-	initiating methionine	UNP O14977
C	1	GLY	-	expression tag	UNP O14977
C	20	ASP	ASN	engineered mutation	UNP O14977

- Molecule 2 is a protein called Ornithine decarboxylase antizyme 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	94	Total	C	N	O	S	434	0	0
			738	481	123	131	3			
2	D	95	Total	C	N	O	S	451	0	0
			755	490	130	132	3			

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	109	MET	-	initiating methionine	UNP P54368
B	229	LEU	-	expression tag	UNP P54368
B	230	GLU	-	expression tag	UNP P54368
B	231	HIS	-	expression tag	UNP P54368
B	232	HIS	-	expression tag	UNP P54368
B	233	HIS	-	expression tag	UNP P54368

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	234	HIS	-	expression tag	UNP P54368
B	235	HIS	-	expression tag	UNP P54368
B	236	HIS	-	expression tag	UNP P54368
D	109	MET	-	initiating methionine	UNP P54368
D	229	LEU	-	expression tag	UNP P54368
D	230	GLU	-	expression tag	UNP P54368
D	231	HIS	-	expression tag	UNP P54368
D	232	HIS	-	expression tag	UNP P54368
D	233	HIS	-	expression tag	UNP P54368
D	234	HIS	-	expression tag	UNP P54368
D	235	HIS	-	expression tag	UNP P54368
D	236	HIS	-	expression tag	UNP P54368

4 Data and refinement statistics

Property	Value	Source
Space group	P 32	Depositor
Cell constants a, b, c, α , β , γ	166.56Å 166.56Å 144.86Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	25.92 – 5.81 25.92 – 5.81	Depositor EDS
% Data completeness (in resolution range)	98.2 (25.92-5.81) 92.9 (25.92-5.81)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.05	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.12 (at 5.64Å)	Xtriage
Refinement program	PHENIX (PHENIX.REFINE:1.7.3_928)	Depositor
R, R_{free}	0.188 , 0.240 0.181 , 0.235	Depositor DCC
R_{free} test set	594 reflections (4.94%)	DCC
Wilson B-factor (Å ²)	333.2	Xtriage
Anisotropy	0.187	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.11 , 890.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.28$	Xtriage
Estimated twinning fraction	0.032 for -h,-k,l 0.058 for h,-h-k,-l 0.037 for -k,-h,-l	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	7456	wwPDB-VP
Average B, all atoms (Å ²)	55.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.02% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.23	0/3039	0.45	1/4120 (0.0%)
1	C	0.23	0/3056	0.46	0/4140
2	B	0.22	0/759	0.44	0/1028
2	D	0.24	0/776	0.52	0/1049
All	All	0.23	0/7630	0.46	1/10337 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	131	LEU	CA-CB-CG	5.83	128.71	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2973	0	2882	10	0
1	C	2990	0	2913	7	0
2	B	738	0	714	2	0
2	D	755	0	741	2	0
All	All	7456	0	7250	21	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (21) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:238:PHE:HB2	1:A:275:TYR:HB2	1.77	0.67
1:C:110:TYR:HB3	1:C:131:LEU:HD13	1.82	0.61
1:A:45:LEU:HD11	1:A:405:TYR:HB3	1.83	0.60
1:A:115:LYS:O	1:A:141:LYS:NZ	2.38	0.55
2:D:139:LEU:HG	2:D:172:VAL:HG12	1.91	0.52
2:B:153:LYS:HA	2:B:156:PHE:CE2	2.45	0.52
1:A:268:ILE:HD12	1:A:268:ILE:H	1.79	0.47
1:C:68:VAL:HG21	1:C:94:GLU:CD	2.35	0.47
1:A:58:VAL:HG12	1:A:59:VAL:HG23	1.96	0.47
1:C:62:ILE:HD12	1:C:268:ILE:HG12	1.97	0.46
1:C:26:ASP:OD1	1:C:27:ASN:N	2.49	0.45
1:A:232:LEU:HA	1:A:232:LEU:HD23	1.88	0.44
1:A:353:LEU:HG	1:A:364:ILE:HD12	1.99	0.44
1:A:58:VAL:HG13	1:A:249:ASN:HD22	1.83	0.42
1:C:287:ILE:HD11	1:C:314:TYR:HB2	2.00	0.42
1:A:29:VAL:HG22	1:A:378:TRP:CE3	2.55	0.42
2:B:208:ARG:HG2	2:B:210:ASP:OD1	2.20	0.42
2:D:141:ILE:O	2:D:141:ILE:HG13	2.19	0.42
1:C:121:LYS:HD3	1:C:121:LYS:HA	1.89	0.42
1:A:45:LEU:HD12	1:A:45:LEU:H	1.85	0.41
1:C:130:ILE:HA	1:C:150:LYS:HB2	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	379/438 (86%)	367 (97%)	12 (3%)	0	100	100
1	C	379/438 (86%)	367 (97%)	12 (3%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	B	92/128 (72%)	90 (98%)	2 (2%)	0	100	100
2	D	93/128 (73%)	91 (98%)	2 (2%)	0	100	100
All	All	943/1132 (83%)	915 (97%)	28 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	321/376 (85%)	320 (100%)	1 (0%)	94	96
1	C	324/376 (86%)	324 (100%)	0	100	100
2	B	76/111 (68%)	75 (99%)	1 (1%)	73	87
2	D	78/111 (70%)	77 (99%)	1 (1%)	73	87
All	All	799/974 (82%)	796 (100%)	3 (0%)	93	95

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	373	LEU
2	B	166	GLN
2	D	204	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	388/438 (88%)	-0.71	1 (0%) 93 91	29, 46, 79, 111	388 (100%)
1	C	389/438 (88%)	-0.69	2 (0%) 90 87	28, 42, 102, 142	389 (100%)
2	B	94/128 (73%)	-0.82	0 100 100	45, 75, 98, 105	94 (100%)
2	D	94/128 (73%)	-0.77	1 (1%) 80 75	44, 71, 98, 120	94 (100%)
All	All	965/1132 (85%)	-0.72	4 (0%) 92 89	28, 49, 96, 142	965 (100%)

All (4) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	112	SER	2.7
2	D	219	GLU	2.3
1	A	10	TYR	2.3
1	C	290	LYS	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers ⓘ

There are no such residues in this entry.