



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 14, 2017 – 10:39 am GMT

PDB ID : 3ZJU
Title : Ternary complex of E .coli leucyl-tRNA synthetase, tRNA(Leu) and the benzoxaborole AN3016 in the editing conformation
Authors : Cusack, S.; Palencia, A.; Crepin, T.; Hernandez, V.; Akama, T.; Baker, S.J.; Bu, W.; Feng, L.; Freund, Y.R.; Liu, L.; Meewan, M.; Mohan, M.; Mao, W.; Rock, F.L.; Sexton, H.; Sheoran, A.; Zhang, Y.; Zhang, Y.; Zhou, Y.; Nieman, J.A.; Anugula, M.R.; Keramane, E.M.; Savariraj, K.; Reddy, D.S.; Sharma, R.; Subedi, R.; Singh, R.; OLeary, A.; Simon, N.L.; DeMarsh, P.L.; Mushtaq, S.; Warner, M.; Livermore, D.M.; Alley, M.R.K.; Plattner, J.J.
Deposited on : 2013-01-18
Resolution : 2.40 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix)	:	1.9-1692
EDS	:	trunk28620
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	3166 (2.40-2.40)
Clashscore	112137	3674 (2.40-2.40)
Ramachandran outliers	110173	3616 (2.40-2.40)
Sidechain outliers	110143	3617 (2.40-2.40)
RSRZ outliers	101464	3195 (2.40-2.40)
RNA backbone	2435	1034 (2.86-1.94)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	880	
2	B	87	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 8359 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called LEUCYL-TRNA SYNTHETASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	820	Total	C	N	O	S	0	0	0
			6517	4147	1103	1228	39			

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-19	MET	-	EXPRESSION TAG	UNP P07813
A	-18	GLY	-	EXPRESSION TAG	UNP P07813
A	-17	SER	-	EXPRESSION TAG	UNP P07813
A	-16	SER	-	EXPRESSION TAG	UNP P07813
A	-15	HIS	-	EXPRESSION TAG	UNP P07813
A	-14	HIS	-	EXPRESSION TAG	UNP P07813
A	-13	HIS	-	EXPRESSION TAG	UNP P07813
A	-12	HIS	-	EXPRESSION TAG	UNP P07813
A	-11	HIS	-	EXPRESSION TAG	UNP P07813
A	-10	HIS	-	EXPRESSION TAG	UNP P07813
A	-9	SER	-	EXPRESSION TAG	UNP P07813
A	-8	SER	-	EXPRESSION TAG	UNP P07813
A	-7	GLY	-	EXPRESSION TAG	UNP P07813
A	-6	LEU	-	EXPRESSION TAG	UNP P07813
A	-5	VAL	-	EXPRESSION TAG	UNP P07813
A	-4	PRO	-	EXPRESSION TAG	UNP P07813
A	-3	ARG	-	EXPRESSION TAG	UNP P07813
A	-2	GLY	-	EXPRESSION TAG	UNP P07813
A	-1	SER	-	EXPRESSION TAG	UNP P07813
A	0	HIS	-	EXPRESSION TAG	UNP P07813

- Molecule 2 is a RNA chain called TRNALEU5 UAA ISOACCEPTOR.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	80	Total	B	C	N	O	P	0	0
			1724	1	771	307	565	80		

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	2	Total 2	Mg 2	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	108	Total 108	O 108	0	0
4	B	8	Total 8	O 8	0	0

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	76.72Å 120.03Å 142.81Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 2.40 29.87 – 2.40	Depositor EDS
% Data completeness (in resolution range)	98.8 (30.00-2.40) 98.8 (29.87-2.40)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.35 (at 2.39Å)	Xtriage
Refinement program	REFMAC 5.6.0085	Depositor
R, R_{free}	0.207 , 0.261 0.207 , 0.260	Depositor DCC
R_{free} test set	2645 reflections (5.40%)	DCC
Wilson B-factor (Å ²)	38.9	Xtriage
Anisotropy	0.283	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 43.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	8359	wwPDB-VP
Average B, all atoms (Å ²)	43.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.76% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, DJF

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.72	0/6671	0.60	1/9054 (0.0%)
2	B	1.08	1/1884 (0.1%)	1.41	15/2934 (0.5%)
All	All	0.81	1/8555 (0.0%)	0.87	16/11988 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	14	A	O5'-C5'	-5.21	1.34	1.42

The worst 5 of 16 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	47(A)	G	O4'-C1'-N9	8.14	114.72	108.20
2	B	28	G	C1'-O4'-C4'	-6.79	104.47	109.90
2	B	28	G	O4'-C1'-N9	6.76	113.61	108.20
2	B	12	G	C3'-C2'-C1'	-6.15	96.58	101.50
2	B	14	A	C5'-C4'-O4'	-6.14	101.73	109.10

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	461	ASP	Peptide
1	A	94	ASN	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6517	0	6379	142	1
2	B	1724	0	877	43	0
3	A	2	0	0	0	0
4	A	108	0	0	4	1
4	B	8	0	0	2	0
All	All	8359	0	7256	182	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

The worst 5 of 182 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:595:ARG:HH11	1:A:595:ARG:HG3	1.00	1.15
2:B:65:C:C6	2:B:65:C:H5''	1.92	1.04
1:A:509:LYS:O	1:A:510:GLU:HB2	1.61	0.99
1:A:695:THR:HG22	1:A:697:ASN:H	1.30	0.97
1:A:426:ARG:HH11	1:A:426:ARG:HG2	1.29	0.97

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:2028:HOH:O	4:A:2071:HOH:O[4_455]	1.59	0.61
1:A:506:PRO:O	1:A:579:GLU:OE1[4_455]	2.16	0.04

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	814/880 (92%)	766 (94%)	40 (5%)	8 (1%)	18	26

5 of 8 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	462	GLY
1	A	600	ARG
1	A	463	ILE
1	A	595	ARG
1	A	597	GLU

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	688/741 (93%)	647 (94%)	41 (6%)	22	35

5 of 41 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	450	LEU
1	A	510	GLU
1	A	837	LYS
1	A	477	THR
1	A	484	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 12 such

sidechains are listed below:

Mol	Chain	Res	Type
1	A	274	ASN
1	A	519	ASN
1	A	745	GLN
1	A	111	GLN
1	A	706	HIS

5.3.3 RNA ⓘ

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
2	B	77/87 (88%)	27 (35%)	0

5 of 27 RNA backbone outliers are listed below:

Mol	Chain	Res	Type
2	B	2	C
2	B	3	C
2	B	4	C
2	B	5	G
2	B	6	G

There are no RNA pucker outliers to report.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	DJF	B	76	2	28,41,42	0.87	0	27,61,64	2.78	4 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical

component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	DJF	B	76	2	-	0/8/52/53	0/6/6/6

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
2	B	76	DJF	N3-C2-N1	-8.58	121.39	128.86
2	B	76	DJF	O17-C16-C15	-3.19	117.14	124.00
2	B	76	DJF	C4-C5-N7	-2.99	106.52	109.41
2	B	76	DJF	O17-C16-C12	9.85	124.06	115.16

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	76	DJF	5	0

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	820/880 (93%)	0.05	33 (4%) 39 38	16, 35, 70, 128	0
2	B	79/87 (90%)	0.72	17 (21%) 1 1	28, 64, 119, 132	0
All	All	899/967 (92%)	0.11	50 (5%) 25 24	16, 36, 78, 132	0

The worst 5 of 50 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	286	ARG	5.6
2	B	31	A	5.0
1	A	287	ASN	5.0
2	B	65	C	4.7
2	B	30	G	4.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	DJF	B	76	36/37	0.97	0.14	-	29,34,45,54	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	MG	A	1861	1/1	0.87	0.15	1.49	47,47,47,47	0
3	MG	A	1862	1/1	0.80	0.12	-0.77	55,55,55,55	0

6.5 Other polymers [i](#)

There are no such residues in this entry.