



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 14, 2017 – 01:11 am GMT

PDB ID : 3ZM9
Title : The mechanism of allosteric coupling in choline kinase a1 revealed by a rationally designed inhibitor
Authors : Sahun-Roncero, M.; Rubio-Ruiz, B.; Saladino, G.; Conejo-Garcia, A.; Espinosa, A.; Velazquez-Campoy, A.; Gervasio, F.L.; Entrena, A.; Hurtado-Guerrero, R.
Deposited on : 2013-02-06
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix)	:	1.9-1692
EDS	:	trunk28620
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	5047 (1.90-1.90)
Clashscore	112137	5731 (1.90-1.90)
Ramachandran outliers	110173	5669 (1.90-1.90)
Sidechain outliers	110143	5670 (1.90-1.90)
RSRZ outliers	101464	5100 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	383	<div> <div>5%</div> <div> <div></div> <div>83%</div> <div>8%</div> <div>8%</div> </div> </div>
1	B	383	<div> <div>12%</div> <div> <div></div> <div>79%</div> <div>10%</div> <div>10%</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	QMQ	A	1458	-	-	-	X

2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 6311 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CHOLINE KINASE ALPHA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	354	Total	C	N	O	S	0	2	0
			2954	1909	498	531	16			
1	B	346	Total	C	N	O	S	0	7	0
			2903	1881	486	519	17			

- Molecule 2 is 1-(4-(4-(4-((6-AMINO-9H-PURIN-9-YL)METHYL)PHENYL)BUTYL)BENZYL)-4- (DIMETHYLAMINO)PYRIDINIUM (three-letter code: QMQ) (formula: C₃₀H₃₄N₇).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	N	0	0
			37	30	7		
2	B	1	Total	C	N	0	0
			37	30	7		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	187	Total 187	O 187	0	0
3	B	193	Total 193	O 193	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: CHOLINE KINASE ALPHA



• Molecule 1: CHOLINE KINASE ALPHA



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	55.75Å 121.21Å 131.86Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	131.86 – 1.90 19.97 – 1.90	Depositor EDS
% Data completeness (in resolution range)	97.0 (131.86-1.90) 97.1 (19.97-1.90)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.00 (at 1.90Å)	Xtriage
Refinement program	REFMAC 5.6.0117	Depositor
R, R_{free}	0.201 , 0.229 0.200 , 0.229	Depositor DCC
R_{free} test set	1722 reflections (2.55%)	DCC
Wilson B-factor (Å ²)	27.3	Xtriage
Anisotropy	0.478	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 42.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	6311	wwPDB-VP
Average B, all atoms (Å ²)	35.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.28% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: QMQ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.54	3/3036 (0.1%)	0.60	0/4088
1	B	0.57	2/2998 (0.1%)	0.63	1/4035 (0.0%)
All	All	0.56	5/6034 (0.1%)	0.61	1/8123 (0.0%)

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	420	TRP	CD2-CE2	5.20	1.47	1.41
1	A	94	TRP	CD2-CE2	5.18	1.47	1.41
1	A	420	TRP	CD2-CE2	5.14	1.47	1.41
1	A	248	TRP	CD2-CE2	5.12	1.47	1.41
1	B	94	TRP	CD2-CE2	5.08	1.47	1.41

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	339	ARG	NE-CZ-NH1	-5.01	117.79	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2954	0	2927	23	0
1	B	2903	0	2896	26	0
2	A	37	0	34	4	0
2	B	37	0	34	2	0
3	A	187	0	0	2	0
3	B	193	0	0	4	0
All	All	6311	0	5891	51	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (51) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:254[A]:GLU:OE2	1:B:290:ARG:NH2	2.16	0.78
1:A:183:MET:HE1	3:A:2056:HOH:O	1.83	0.78
1:A:126:GLN:OE1	1:A:142:LYS:HD3	1.86	0.74
1:B:192:LEU:CD1	1:B:238:MET:HE1	2.22	0.70
1:B:192:LEU:HD12	1:B:238:MET:HE1	1.74	0.69
1:A:183:MET:CE	1:A:331:PHE:HB2	2.22	0.69
1:B:116:ILE:HD11	1:B:126:GLN:HB2	1.75	0.69
1:B:357:GLU:HG2	3:B:2139:HOH:O	1.94	0.66
1:B:235:PHE:O	1:B:238:MET:HG2	2.02	0.60
1:B:135:THR:HG22	1:B:137:GLY:H	1.69	0.56
1:A:183:MET:HE3	1:A:331:PHE:HB2	1.88	0.56
1:A:183:MET:HE1	1:A:331:PHE:HB2	1.87	0.56
1:B:192:LEU:CD1	1:B:238:MET:CE	2.84	0.55
1:B:270:ARG:HH11	1:B:270:ARG:CG	2.20	0.55
1:B:190:ARG:HH11	1:B:238:MET:HE2	1.72	0.54
1:B:384:PRO:HB3	1:B:391:GLU:HG3	1.90	0.54
1:A:332:GLU:HB3	3:A:2138:HOH:O	2.08	0.53
1:B:101:GLY:O	1:B:104:ARG:HG2	2.08	0.53
1:A:183:MET:HE3	1:A:331:PHE:CB	2.39	0.53
1:A:213:ARG:NH1	2:A:1458:QMQ:HAR2	2.25	0.51
1:A:391:GLU:HA	1:A:398:LYS:HE3	1.92	0.51
1:B:332:GLU:HB3	3:B:2119:HOH:O	2.09	0.51
1:B:117:ARG:HD2	2:B:1458:QMQ:HAT2	1.94	0.50
1:A:387:GLN:O	1:A:388:ASN:HB2	2.11	0.50
1:A:354:TYR:CE1	2:A:1458:QMQ:HAO	2.48	0.48
2:B:1458:QMQ:HAM	2:B:1458:QMQ:HAB1	1.67	0.47
1:B:270:ARG:HG3	1:B:270:ARG:NH1	2.29	0.47
1:B:262:ARG:HG3	3:B:2082:HOH:O	2.13	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:357:GLU:CG	3:B:2139:HOH:O	2.59	0.46
1:A:329:ILE:HD11	2:A:1458:QMQ:HAH	1.98	0.46
1:B:110:GLU:O	1:B:130:PRO:HG3	2.16	0.46
1:A:244:LYS:HE3	1:A:336:TYR:CZ	2.49	0.46
1:B:137:GLY:HA2	1:B:138:ASP:HA	1.66	0.46
1:B:423:TRP:O	1:B:427:GLN:HG2	2.16	0.46
1:A:122:ASN:ND2	1:A:146:ARG:HH11	2.13	0.45
1:B:270:ARG:NH1	1:B:270:ARG:CG	2.77	0.45
1:A:396:GLU:HG2	1:A:396:GLU:H	1.47	0.45
1:B:113:ILE:HA	1:B:126:GLN:O	2.18	0.44
2:A:1458:QMQ:HAB1	2:A:1458:QMQ:HAM	1.74	0.43
1:A:183:MET:HG3	1:A:242:PHE:HE2	1.84	0.42
1:B:238:MET:HB2	1:B:238:MET:HE2	1.68	0.42
1:A:278:LEU:HA	1:A:278:LEU:HD12	1.91	0.42
1:A:318:ARG:HD2	1:A:324:GLN:O	2.20	0.42
1:B:274:LEU:HG	1:B:278:LEU:HD22	2.01	0.41
1:A:423:TRP:O	1:A:427:GLN:HG2	2.21	0.41
1:A:318:ARG:HG2	1:A:321:SER:HB2	2.02	0.41
1:B:449:PHE:O	1:B:453[B]:ARG:HG3	2.20	0.41
1:A:363:ARG:HB3	1:A:363:ARG:NH1	2.36	0.41
1:A:274:LEU:HG	1:A:278:LEU:HD22	2.03	0.41
1:A:116:ILE:HD11	1:A:126:GLN:HB2	2.04	0.40
1:B:217:GLU:CD	1:B:217:GLU:H	2.25	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	352/383 (92%)	345 (98%)	6 (2%)	1 (0%)	44	34
1	B	349/383 (91%)	340 (97%)	9 (3%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
All	All	701/766 (92%)	685 (98%)	15 (2%)	1 (0%)	55 45

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	388	ASN

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	320/342 (94%)	309 (97%)	11 (3%)	42 32
1	B	317/342 (93%)	303 (96%)	14 (4%)	33 22
All	All	637/684 (93%)	612 (96%)	25 (4%)	36 26

All (25) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	120	LEU
1	A	144	LEU
1	A	213	ARG
1	A	239	LYS
1	A	244	LYS
1	A	278	LEU
1	A	332	GLU
1	A	342	ASP
1	A	396	GLU
1	A	407	LEU
1	A	455	LEU
1	B	120	LEU
1	B	131	ASP
1	B	135	THR
1	B	142	LYS
1	B	199	ILE
1	B	238	MET

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	270	ARG
1	B	272	LYS
1	B	278	LEU
1	B	332	GLU
1	B	342	ASP
1	B	389	ASP
1	B	407	LEU
1	B	457	VAL

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	122	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	QMQ	A	1458	-	39,41,41	2.02	10 (25%)	46,56,56	2.02	2 (4%)
2	QMQ	B	1458	-	39,41,41	1.95	10 (25%)	46,56,56	2.18	3 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	QMQ	A	1458	-	-	0/19/19/19	0/5/5/5
2	QMQ	B	1458	-	-	0/19/19/19	0/5/5/5

All (20) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1458	QMQ	CAV-CBC	-4.99	1.39	1.51
2	A	1458	QMQ	CAV-CBC	-4.12	1.41	1.51
2	A	1458	QMQ	CAW-CBD	-3.91	1.42	1.51
2	B	1458	QMQ	CAW-CBD	-3.87	1.42	1.51
2	A	1458	QMQ	CAU-CBB	-3.50	1.41	1.51
2	B	1458	QMQ	CAU-CBB	-3.31	1.42	1.51
2	B	1458	QMQ	CAT-CBA	-3.29	1.42	1.51
2	A	1458	QMQ	CAT-CBA	-3.18	1.42	1.51
2	A	1458	QMQ	C5-C4	-2.39	1.35	1.40
2	B	1458	QMQ	C5-C4	-2.30	1.35	1.40
2	B	1458	QMQ	CAV-N9	2.78	1.54	1.49
2	B	1458	QMQ	CAO-NBK	2.79	1.41	1.34
2	A	1458	QMQ	CAO-NBK	3.04	1.41	1.34
2	B	1458	QMQ	C2-N1	3.13	1.39	1.33
2	A	1458	QMQ	C2-N1	3.31	1.40	1.33
2	A	1458	QMQ	CAV-N9	3.34	1.55	1.49
2	B	1458	QMQ	CAP-NBK	3.48	1.42	1.34
2	A	1458	QMQ	CAP-NBK	3.75	1.43	1.34
2	B	1458	QMQ	C2-N3	4.73	1.40	1.32
2	A	1458	QMQ	C2-N3	4.77	1.40	1.32

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1458	QMQ	N3-C2-N1	-13.20	117.37	128.86
2	A	1458	QMQ	N3-C2-N1	-12.53	117.94	128.86
2	B	1458	QMQ	C4-C5-N7	-2.07	107.41	109.41

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1458	QMQ	N6-C6-N1	2.02	122.76	118.77
2	B	1458	QMQ	CAV-N9-C8	4.67	131.21	125.67

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1458	QMQ	4	0
2	B	1458	QMQ	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	354/383 (92%)	0.45	20 (5%) 25 28	19, 31, 59, 87	0
1	B	346/383 (90%)	0.66	45 (13%) 4 4	15, 30, 69, 88	0
All	All	700/766 (91%)	0.55	65 (9%) 9 10	15, 31, 66, 88	0

All (65) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	120	LEU	7.6
1	A	320	ASN	7.4
1	A	457	VAL	6.6
1	B	89	ARG	5.9
1	A	150	ALA	5.8
1	B	132	THR	5.7
1	B	113	ILE	5.5
1	A	456	GLY	5.2
1	B	116	ILE	5.1
1	B	138	ASP	5.1
1	B	131	ASP	4.9
1	A	388	ASN	4.7
1	B	115	VAL	4.5
1	A	120	LEU	4.3
1	B	432[A]	SER	4.2
1	B	119	GLY	4.1
1	B	388	ASN	4.1
1	B	150	ALA	4.0
1	B	112	HIS	4.0
1	A	322	GLU	4.0
1	A	80	ASP	3.9
1	B	136	LEU	3.6
1	A	175	GLU	3.5
1	A	119	GLY	3.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	389	ASP	3.4
1	B	135	THR	3.3
1	B	87	THR	3.3
1	A	357	GLU	3.2
1	B	114	SER	3.1
1	A	317	GLY	3.0
1	B	145	LEU	2.9
1	B	258	LYS	2.9
1	A	324	GLN	2.8
1	B	121	SER	2.7
1	B	109	ASP	2.7
1	B	130	PRO	2.7
1	B	137	GLY	2.7
1	B	149	GLY	2.7
1	B	111	PHE	2.7
1	B	129	LEU	2.7
1	B	143	VAL	2.6
1	B	148	TYR	2.6
1	B	95	CYS	2.6
1	B	127	CYS	2.6
1	B	177	MET	2.6
1	A	118	GLY	2.5
1	B	107	ARG	2.5
1	A	121	SER	2.4
1	A	149	GLY	2.4
1	B	134	ALA	2.4
1	B	262	ARG	2.3
1	A	395	THR	2.3
1	A	262	ARG	2.2
1	B	421	GLY	2.1
1	B	422	LEU	2.1
1	B	396	GLU	2.1
1	B	434	GLU	2.1
1	B	117	ARG	2.1
1	B	389	ASP	2.1
1	B	176	ALA	2.1
1	A	391	GLU	2.0
1	B	106	LEU	2.0
1	B	90	ARG	2.0
1	B	99	LEU	2.0
1	B	147	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	QMQ	A	1458	37/37	0.80	0.20	2.69	33,46,57,58	0
2	QMQ	B	1458	37/37	0.90	0.13	-0.18	28,35,47,48	0

6.5 Other polymers [i](#)

There are no such residues in this entry.