



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 14, 2017 – 10:42 am GMT

PDB ID : 1ZO4  
Title : Crystal Structure Of A328S Mutant Of The Heme Domain Of P450BM-3  
Authors : Hegda, A.; Chen, B.; Haines, D.C.; Bondlela, M.; Mullin, D.; Graham, S.E.;  
Tomchick, D.R.; Machius, M.; Peterson, J.A.  
Deposited on : 2005-05-12  
Resolution : 1.46 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : 1.9-1692  
EDS : trunk28620  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : recalc28949


# 1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.46 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	1510 (1.48-1.44)
Clashscore	112137	1573 (1.48-1.44)
Ramachandran outliers	110173	1555 (1.48-1.44)
Sidechain outliers	110143	1555 (1.48-1.44)
RSRZ outliers	101464	1516 (1.48-1.44)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	473	<div> <div>7%</div> <div> <div></div> <div>85%</div> <div>11%</div> <div>.</div> </div> </div>
1	B	473	<div> <div>4%</div> <div> <div></div> <div>84%</div> <div>12%</div> <div>.</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	MES	A	1491	-	-	-	X
3	MES	B	1490	-	-	-	X
4	GOL	A	1472	-	-	X	X
4	GOL	A	1473	-	-	-	X
4	GOL	A	1482	-	-	-	X
4	GOL	A	1484	-	-	-	X
4	GOL	A	1485	-	-	-	X
4	GOL	A	1486	-	-	-	X
4	GOL	A	1487	-	-	-	X
4	GOL	A	1489	-	-	-	X
4	GOL	B	1470	-	-	-	X
4	GOL	B	1474	-	-	X	X
4	GOL	B	1475	-	-	-	X
4	GOL	B	1478	-	-	-	X
4	GOL	B	1479	-	-	-	X
4	GOL	B	1483	-	-	X	X

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 8734 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Bifunctional P-450:NADPH-P450 reductase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	456	Total	C	N	O	S	0	18	0
			3829	2440	657	714	18			
1	B	457	Total	C	N	O	S	0	19	0
			3845	2451	659	717	18			

There are 8 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	GLY	-	EXPRESSION TAG	UNP P14779
A	-1	ALA	-	EXPRESSION TAG	UNP P14779
A	0	MET	-	EXPRESSION TAG	UNP P14779
A	328	SER	ALA	ENGINEERED	UNP P14779
B	-2	GLY	-	EXPRESSION TAG	UNP P14779
B	-1	ALA	-	EXPRESSION TAG	UNP P14779
B	0	MET	-	EXPRESSION TAG	UNP P14779
B	328	SER	ALA	ENGINEERED	UNP P14779

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula:  $C_{34}H_{32}FeN_4O_4$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
2	B	1	Total 43	C 34	Fe 1	N 4	O 4	0	0

- Molecule 3 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (three-letter code: MES) (formula:  $\text{C}_6\text{H}_{13}\text{NO}_4\text{S}$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	B	1	Total	C	N	O	S	0	0
			12	6	1	4	1		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	S	0	0
			12	6	1	4	1		

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula:  $C_3H_8O_3$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		
4	B	1	Total	C	O	0	0
			6	3	3		
4	B	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		
4	B	1	Total	C	O	0	0
			6	3	3		
4	B	1	Total	C	O	0	0
			6	3	3		
4	B	1	Total	C	O	0	0
			6	3	3		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total 6	C 3	O 3	0	0
4	A	1	Total 6	C 3	O 3	0	0
4	A	1	Total 6	C 3	O 3	0	0
4	B	1	Total 6	C 3	O 3	0	0
4	A	1	Total 6	C 3	O 3	0	0
4	A	1	Total 6	C 3	O 3	0	0
4	A	1	Total 6	C 3	O 3	0	0
4	A	1	Total 6	C 3	O 3	0	0
4	B	1	Total 6	C 3	O 3	0	0
4	A	1	Total 6	C 3	O 3	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	406	Total 406	O 406	0	2
5	B	424	Total 424	O 424	0	2


### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Bifunctional P-450:NADPH-P450 reductase



- Molecule 1: Bifunctional P-450:NADPH-P450 reductase





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	58.83Å 153.21Å 61.58Å 90.00° 94.33° 90.00°	Depositor
Resolution (Å)	29.30 – 1.46 29.33 – 1.46	Depositor EDS
% Data completeness (in resolution range)	94.6 (29.30-1.46) 94.6 (29.33-1.46)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.70 (at 1.46Å)	Xtriage
Refinement program	CNS 1.0	Depositor
R, $R_{free}$	0.189 , 0.216 0.185 , 0.211	Depositor DCC
$R_{free}$ test set	8896 reflections (5.28%)	DCC
Wilson B-factor (Å <sup>2</sup> )	18.1	Xtriage
Anisotropy	0.445	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 45.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.017 for l,-k,h	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	8734	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.89% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, MES, HEM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.50	0/3914	0.71	2/5283 (0.0%)
1	B	0.51	0/3930	0.72	2/5302 (0.0%)
All	All	0.50	0/7844	0.71	4/10585 (0.0%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	328	SER	N-CA-C	-6.54	93.33	111.00
1	A	328	SER	N-CA-C	-6.34	93.89	111.00
1	A	329	PRO	N-CA-C	5.86	127.33	112.10
1	B	329	PRO	N-CA-C	5.51	126.42	112.10

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3829	0	3807	56	0
1	B	3845	0	3825	59	0
2	A	43	0	30	0	0
2	B	43	0	30	0	0
3	A	12	0	13	1	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	12	0	13	0	0
4	A	66	0	88	16	0
4	B	54	0	72	25	0
5	A	406	0	0	13	0
5	B	424	0	0	9	0
All	All	8734	0	7878	122	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 122 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:241:LYS:H	4:B:1483:GOL:H31	1.23	1.01
1:B:392:PRO:HD2	4:B:1474:GOL:H2	1.46	0.96
4:A:1472:GOL:H32	1:B:129:LYS:HE3	1.56	0.87
1:A:2:ILE:HD12	1:A:346:PRO:HD3	1.57	0.87
1:A:171:HIS:HD2	1:A:173:PHE:H	1.20	0.86

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	472/473 (100%)	464 (98%)	8 (2%)	0	100	100
1	B	474/473 (100%)	464 (98%)	10 (2%)	0	100	100
All	All	946/946 (100%)	928 (98%)	18 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	418/413 (101%)	413 (99%)	5 (1%)	75	44
1	B	419/413 (102%)	413 (99%)	6 (1%)	71	38
All	All	837/826 (101%)	826 (99%)	11 (1%)	71	40

5 of 11 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	344	GLU
1	B	47	ARG
1	B	231	ASP
1	A	182	ASP
1	B	158	PHE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 23 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	403	GLN
1	B	21	ASN
1	B	361	HIS
1	A	426	HIS
1	B	73	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

24 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
4	GOL	A	1471	-	5,5,5	0.44	0	5,5,5	0.17	0
4	GOL	A	1472	-	5,5,5	0.51	0	5,5,5	0.18	0
4	GOL	A	1473	-	5,5,5	0.42	0	5,5,5	0.19	0
4	GOL	A	1476	-	5,5,5	0.22	0	5,5,5	0.22	0
4	GOL	A	1481	-	5,5,5	0.44	0	5,5,5	0.21	0
4	GOL	A	1482	-	5,5,5	0.45	0	5,5,5	0.18	0
4	GOL	A	1484	-	5,5,5	0.35	0	5,5,5	0.18	0
4	GOL	A	1485	-	5,5,5	0.41	0	5,5,5	0.25	0
4	GOL	A	1486	-	5,5,5	0.44	0	5,5,5	0.20	0
4	GOL	A	1487	-	5,5,5	0.70	0	5,5,5	0.11	0
4	GOL	A	1489	-	5,5,5	0.49	0	5,5,5	0.19	0
3	MES	A	1491	-	12,12,12	2.32	6 (50%)	14,16,16	1.43	1 (7%)
2	HEM	A	471	1,5	28,50,50	1.59	5 (17%)	17,82,82	1.85	6 (35%)
4	GOL	B	1470	-	5,5,5	0.59	0	5,5,5	0.18	0
4	GOL	B	1474	-	5,5,5	0.55	0	5,5,5	0.29	0
4	GOL	B	1475	-	5,5,5	0.57	0	5,5,5	0.21	0
4	GOL	B	1477	-	5,5,5	0.44	0	5,5,5	0.22	0
4	GOL	B	1478	-	5,5,5	0.41	0	5,5,5	0.14	0
4	GOL	B	1479	-	5,5,5	0.46	0	5,5,5	0.12	0
4	GOL	B	1480	-	5,5,5	0.51	0	5,5,5	0.18	0
4	GOL	B	1483	-	5,5,5	0.41	0	5,5,5	0.19	0
4	GOL	B	1488	-	5,5,5	0.53	0	5,5,5	0.23	0
3	MES	B	1490	-	12,12,12	2.12	5 (41%)	14,16,16	1.38	1 (7%)
2	HEM	B	471	1,5	28,50,50	1.70	7 (25%)	17,82,82	1.74	7 (41%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	A	1471	-	-	0/4/4/4	0/0/0/0
4	GOL	A	1472	-	-	0/4/4/4	0/0/0/0
4	GOL	A	1473	-	-	0/4/4/4	0/0/0/0
4	GOL	A	1476	-	-	0/4/4/4	0/0/0/0
4	GOL	A	1481	-	-	0/4/4/4	0/0/0/0
4	GOL	A	1482	-	-	0/4/4/4	0/0/0/0
4	GOL	A	1484	-	-	0/4/4/4	0/0/0/0
4	GOL	A	1485	-	-	0/4/4/4	0/0/0/0
4	GOL	A	1486	-	-	0/4/4/4	0/0/0/0
4	GOL	A	1487	-	-	0/4/4/4	0/0/0/0
4	GOL	A	1489	-	-	0/4/4/4	0/0/0/0
3	MES	A	1491	-	-	0/6/14/14	0/1/1/1
2	HEM	A	471	1,5	-	0/6/54/54	0/0/8/8
4	GOL	B	1470	-	-	0/4/4/4	0/0/0/0
4	GOL	B	1474	-	-	0/4/4/4	0/0/0/0
4	GOL	B	1475	-	-	0/4/4/4	0/0/0/0
4	GOL	B	1477	-	-	0/4/4/4	0/0/0/0
4	GOL	B	1478	-	-	0/4/4/4	0/0/0/0
4	GOL	B	1479	-	-	0/4/4/4	0/0/0/0
4	GOL	B	1480	-	-	0/4/4/4	0/0/0/0
4	GOL	B	1483	-	-	0/4/4/4	0/0/0/0
4	GOL	B	1488	-	-	0/4/4/4	0/0/0/0
3	MES	B	1490	-	-	0/6/14/14	0/1/1/1
2	HEM	B	471	1,5	-	0/6/54/54	0/0/8/8

The worst 5 of 23 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	1491	MES	O2S-S	-3.65	1.34	1.45
3	B	1490	MES	O2S-S	-3.44	1.35	1.45
2	B	471	HEM	C3C-C2C	-2.31	1.37	1.40
2	B	471	HEM	C3B-C2B	-2.27	1.37	1.40
2	B	471	HEM	C1B-NB	2.11	1.39	1.36

The worst 5 of 15 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	471	HEM	CMD-C2D-C1D	-2.96	123.92	128.46
2	B	471	HEM	CMA-C3A-C4A	-2.53	124.58	128.46
2	B	471	HEM	CMD-C2D-C1D	-2.45	124.70	128.46

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	471	HEM	CMA-C3A-C4A	-2.12	125.21	128.46
2	B	471	HEM	CMA-C3A-C2A	2.17	129.03	124.94

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

17 monomers are involved in 42 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	1471	GOL	1	0
4	A	1472	GOL	4	0
4	A	1473	GOL	1	0
4	A	1481	GOL	1	0
4	A	1482	GOL	1	0
4	A	1484	GOL	2	0
4	A	1485	GOL	1	0
4	A	1486	GOL	3	0
4	A	1487	GOL	1	0
4	A	1489	GOL	1	0
3	A	1491	MES	1	0
4	B	1470	GOL	1	0
4	B	1474	GOL	12	0
4	B	1475	GOL	2	0
4	B	1478	GOL	3	0
4	B	1483	GOL	5	0
4	B	1488	GOL	3	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	456/473 (96%)	0.25	33 (7%) 16 17	14, 23, 45, 57	0
1	B	457/473 (96%)	0.11	19 (4%) 37 38	15, 24, 42, 54	0
All	All	913/946 (96%)	0.18	52 (5%) 24 26	14, 24, 45, 57	0

The worst 5 of 52 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	2	ILE	11.8
1	B	191	ALA	7.9
1	A	1	THR	7.5
1	B	2	ILE	7.4
1	A	191	ALA	6.9

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.



Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
4	GOL	A	1485	6/6	0.81	0.25	14.03	49,50,52,53	0
4	GOL	B	1474	6/6	0.92	0.23	13.22	21,41,44,44	0
4	GOL	A	1473	6/6	0.88	0.17	10.84	38,43,45,49	0
4	GOL	B	1475	6/6	0.83	0.19	8.90	31,43,43,43	0
4	GOL	B	1483	6/6	0.88	0.44	8.70	45,47,49,50	0
4	GOL	B	1478	6/6	0.75	0.46	8.37	45,47,50,52	0
4	GOL	A	1487	6/6	0.86	0.25	5.46	41,41,47,53	0
4	GOL	B	1470	6/6	0.87	0.21	5.31	25,44,50,50	0
4	GOL	A	1486	6/6	0.82	0.28	5.23	42,46,48,49	0
4	GOL	A	1484	6/6	0.82	0.22	4.83	40,46,52,53	0
3	MES	B	1490	12/12	0.93	0.14	4.45	36,40,42,42	0
3	MES	A	1491	12/12	0.77	0.29	4.25	48,52,59,60	0
4	GOL	A	1489	6/6	0.72	0.22	4.00	47,48,48,53	0
4	GOL	A	1472	6/6	0.84	0.21	3.89	42,45,47,48	0
4	GOL	A	1482	6/6	0.79	0.33	3.66	50,51,54,54	0
4	GOL	B	1479	6/6	0.84	0.23	2.08	35,46,50,51	0
4	GOL	A	1476	6/6	0.81	0.12	1.49	45,47,47,49	0
2	HEM	A	471	43/43	0.99	0.12	0.69	12,14,17,29	0
2	HEM	B	471	43/43	0.98	0.11	0.41	12,16,18,29	0
4	GOL	A	1481	6/6	0.64	0.25	-	47,50,52,54	0
4	GOL	B	1477	6/6	0.77	0.35	-	50,52,53,55	0
4	GOL	A	1471	6/6	0.80	0.27	-	42,47,48,51	0
4	GOL	B	1480	6/6	0.74	0.37	-	48,48,50,50	0
4	GOL	B	1488	6/6	0.81	0.22	-	45,47,48,51	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.