



# Full wwPDB X-ray Structure Validation Report ⓘ

Sep 18, 2017 – 06:37 PM EDT

PDB ID : 4ZTI  
Title : Ebola virus nucleoprotein bound to VP35 chaperoning peptide P212121  
Authors : Kirchdoerfer, R.N.; Abelson, D.M.; Saphire, E.O.  
Deposited on : unknown  
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20029824  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20029824


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	3166 (2.40-2.40)
Clashscore	112137	3674 (2.40-2.40)
Ramachandran outliers	110173	3616 (2.40-2.40)
Sidechain outliers	110143	3617 (2.40-2.40)
RSRZ outliers	101464	3195 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	381	<div> <div>3%</div> <div>82%</div> <div>6%</div> <div>12%</div> </div>
1	B	381	<div> <div>6%</div> <div>86%</div> <div>•</div> <div>12%</div> </div>

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 5508 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Polymerase cofactor VP35,Nucleoprotein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	336	Total	C	N	O	S	0	1	0
			2636	1682	465	478	11			
1	B	337	Total	C	N	O	S	0	0	0
			2649	1692	468	478	11			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-13	GLY	-	expression tag	UNP Q05127
B	-13	GLY	-	expression tag	UNP Q05127


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	114	Total	O	0	0
			114	114		
2	B	109	Total	O	0	0
			109	109		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Polymerase cofactor VP35,Nucleoprotein



- Molecule 1: Polymerase cofactor VP35,Nucleoprotein



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	94.88Å 94.95Å 112.11Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.27 – 2.40 48.27 – 2.40	Depositor EDS
% Data completeness (in resolution range)	99.9 (48.27-2.40) 100.0 (48.27-2.40)	Depositor EDS
$R_{merge}$	0.15	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.84 (at 2.39Å)	Xtriage
Refinement program	REFMAC 5.8.0073	Depositor
R, $R_{free}$	0.192 , 0.247 0.192 , 0.241	Depositor DCC
$R_{free}$ test set	2050 reflections (5.37%)	DCC
Wilson B-factor (Å <sup>2</sup> )	53.8	Xtriage
Anisotropy	0.345	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 52.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.021 for k,h,-l	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	5508	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	65.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.92% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.70	0/2686	0.71	0/3621
1	B	0.71	0/2697	0.70	0/3636
All	All	0.71	0/5383	0.70	0/7257

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2636	0	2655	13	0
1	B	2649	0	2678	8	0
2	A	114	0	0	1	0
2	B	109	0	0	1	0
All	All	5508	0	5333	20	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (20) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:310:HIS:HD2	2:A:417:HOH:O	1.84	0.60

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:179:HIS:HB3	1:B:185:ILE:HG22	1.85	0.57
1:A:1:TRP:O	1:A:5:GLN:HG2	2.07	0.54
1:A:6:LEU:HD11	1:A:16:ILE:HD12	1.91	0.52
1:A:118:GLU:HA	1:A:121:LEU:HD12	1.93	0.50
1:B:342:GLN:HB3	2:B:464:HOH:O	2.15	0.46
1:A:165:GLU:HG3	1:A:201:PHE:CE2	2.52	0.45
1:B:45:GLN:HA	1:B:87:TYR:CZ	2.51	0.45
1:A:311:GLY:O	1:B:165:GLU:HG2	2.15	0.45
1:A:211:LYS:HE3	1:A:299:LEU:O	2.17	0.45
1:B:45:GLN:HA	1:B:87:TYR:CE2	2.51	0.45
1:A:56:ILE:HG13	1:A:117:LEU:HD21	2.00	0.44
1:A:169:LEU:HD22	1:A:197:MET:HB3	1.98	0.43
1:B:5:GLN:HE22	1:B:10:ARG:HD3	1.83	0.43
1:A:45:GLN:HA	1:A:87:TYR:CZ	2.54	0.42
1:B:165:GLU:H	1:B:165:GLU:CD	2.22	0.42
1:B:118:GLU:HA	1:B:121:LEU:HD12	2.02	0.41
1:A:79:LEU:HD21	1:A:87:TYR:CE2	2.56	0.41
1:A:46:VAL:HG21	1:A:52:ILE:HG13	2.02	0.41
1:A:246:ILE:O	1:A:249:THR:HG22	2.22	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	327/381 (86%)	316 (97%)	11 (3%)	0	100	100
1	B	329/381 (86%)	319 (97%)	10 (3%)	0	100	100
All	All	656/762 (86%)	635 (97%)	21 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	278/316 (88%)	277 (100%)	1 (0%)	93	97
1	B	281/316 (89%)	280 (100%)	1 (0%)	93	97
All	All	559/632 (88%)	557 (100%)	2 (0%)	93	97

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	-3	GLU
1	B	-3	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	149	GLN
1	B	5	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.



## 5.6 Ligand geometry

There are no ligands in this entry.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	336/381 (88%)	0.10	13 (3%) 40 39	37, 58, 108, 137	0
1	B	337/381 (88%)	0.16	22 (6%) 20 18	37, 60, 110, 154	0
All	All	673/762 (88%)	0.13	35 (5%) 28 26	37, 59, 109, 154	0

All (35) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	260	ARG	5.4
1	A	360	SER	4.5
1	B	130	ILE	4.4
1	B	261	GLY	4.2
1	A	357	TYR	4.2
1	B	357	TYR	3.9
1	A	362	GLU	3.7
1	B	365	HIS	3.5
1	A	116	ARG	3.4
1	B	117	LEU	3.4
1	B	259	GLU	3.1
1	B	134	LEU	3.1
1	B	361	ARG	2.9
1	A	355	GLN	2.9
1	B	360	SER	2.8
1	A	120	LEU	2.7
1	A	134	LEU	2.6
1	B	355	GLN	2.6
1	B	133	THR	2.5
1	B	114	VAL	2.5
1	B	257	LYS	2.5
1	A	28	TYR	2.4
1	B	112	ASP	2.4
1	A	52	ILE	2.4

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	110	LYS	2.4
1	A	115	LYS	2.4
1	B	362	GLU	2.4
1	A	61	GLU	2.3
1	B	358	ALA	2.2
1	B	132	ARG	2.2
1	B	120	LEU	2.1
1	A	184	LEU	2.1
1	B	61	GLU	2.1
1	B	116	ARG	2.1
1	A	359	GLU	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.