



wwPDB X-ray Structure Validation Summary Report ⓘ

Nov 5, 2018 – 12:11 PM EST

PDB ID : 5YNC
Title : Crystal structure of Pullulanase from *Klebsiella pneumoniae* complex at 1 mM beta-cyclodextrin
Authors : Saka, N.; Iwamoto, H.; Takahashi, N.; Mizutani, K.; Mikami, B.
Deposited on : 2017-10-24
Resolution : 2.32 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.3 (157068), CSD as539be (2018)
Xtriage (Phenix) : 1.13
EDS : rb-20031633
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)
Refmac : 5.8.0158
CCP4 : 7.0 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20031633


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.32 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	111664	5225 (2.34-2.30)
Clashscore	122126	5849 (2.34-2.30)
Ramachandran outliers	120053	5790 (2.34-2.30)
Sidechain outliers	120020	5789 (2.34-2.30)
RSRZ outliers	108989	5109 (2.34-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1102	<div> <div>3%</div> <div>83%</div> <div>12%</div> <div>5%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	GLC	A	1110	-	-	-	X

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 8683 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PulA protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	1052	Total	C	N	O	S	0	7	0
			8118	5065	1392	1635	26			


There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	680	LEU	GLY	conflict	UNP W9BQ28

- Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	5	Total	Ca	0	0
			5	5		

- Molecule 3 is ACETATE ION (three-letter code: ACT) (formula: C₂H₃O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			4	2	2		
3	A	1	Total	C	O	0	0
			4	2	2		

- Molecule 4 is ALPHA-D-GLUCOSE (three-letter code: GLC) (formula: C₆H₁₂O₆).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			11	6	5		
4	A	1	Total	C	O	0	0
			11	6	5		
4	A	1	Total	C	O	0	0
			11	6	5		
4	A	1	Total	C	O	0	0
			11	6	5		
4	A	1	Total	C	O	0	0
			11	6	5		
4	A	1	Total	C	O	0	0
			11	6	5		
4	A	1	Total	C	O	0	0
			11	6	5		
4	A	1	Total	C	O	0	0
			11	6	5		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			11	6	5		
4	A	1	Total	C	O	0	0
			11	6	5		
4	A	1	Total	C	O	0	0
			11	6	5		
4	A	1	Total	C	O	0	0
			11	6	5		

- Molecule 5 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			6	3	3		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	382	Total	O	0	10
			392	392		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Pula protein



4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	88.92Å 88.92Å 299.53Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.68 – 2.32 49.68 – 2.32	Depositor EDS
% Data completeness (in resolution range)	99.7 (49.68-2.32) 99.7 (49.68-2.32)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.69 (at 2.32Å)	Xtriage
Refinement program	PHENIX (1.10.1 _2155: ???)	Depositor
R, R_{free}	0.187 , 0.254 0.187 , 0.254	Depositor DCC
R_{free} test set	2694 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å ²)	33.4	Xtriage
Anisotropy	0.545	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 45.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	8683	wwPDB-VP
Average B, all atoms (Å ²)	38.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.73% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, CA, GLC, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.42	1/8283 (0.0%)	0.58	1/11276 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	539	ASN	C-N	11.07	1.55	1.34

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	222	LYS	CD-CE-NZ	6.35	126.30	111.70

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	8118	0	7813	100	0
2	A	5	0	0	0	0
3	A	8	0	6	0	0
4	A	154	0	126	0	0
5	A	6	0	8	0	0
6	A	392	0	0	8	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	8683	0	7953	100	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 100 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:197:ILE:HD11	1:A:266:ALA:HB3	1.20	1.11
1:A:543:GLN:HG3	1:A:916:ARG:HA	1.35	1.08
1:A:222:LYS:HE2	1:A:222:LYS:HA	1.69	0.74
1:A:266:ALA:HB1	1:A:271:ILE:O	1.88	0.74
1:A:301:LEU:HD13	1:A:395:ASP:HB2	1.70	0.73

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	1057/1102 (96%)	1016 (96%)	30 (3%)	11 (1%)	17 18

5 of 11 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	208	VAL
1	A	249	ASP
1	A	643	CYS
1	A	39	ASP
1	A	66	SER

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	873/907 (96%)	855 (98%)	18 (2%)	56 73

5 of 18 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	537[A]	LYS
1	A	537[B]	LYS
1	A	925	ASP
1	A	374	TYR
1	A	482	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	711	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 22 ligands modelled in this entry, 5 are monoatomic - leaving 17 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	ACT	A	1106	-	1,3,3	0.92	0	0,3,3	0.00	-
3	ACT	A	1107	-	1,3,3	1.21	0	0,3,3	0.00	-
4	GLC	A	1108	4	11,11,12	0.63	0	15,15,17	0.87	0
4	GLC	A	1109	4	11,11,12	0.61	0	15,15,17	1.04	1 (6%)
4	GLC	A	1110	4	11,11,12	0.58	0	15,15,17	1.50	2 (13%)
4	GLC	A	1111	4	11,11,12	0.70	0	15,15,17	1.28	3 (20%)
4	GLC	A	1112	4	11,11,12	0.57	0	15,15,17	1.26	2 (13%)
4	GLC	A	1113	4	11,11,12	0.57	0	15,15,17	1.68	3 (20%)
4	GLC	A	1114	4	11,11,12	0.72	0	15,15,17	1.37	2 (13%)
4	GLC	A	1115	4	11,11,12	0.63	0	15,15,17	0.90	0
4	GLC	A	1116	4	11,11,12	0.60	0	15,15,17	0.91	0
4	GLC	A	1117	4	11,11,12	0.51	0	15,15,17	0.98	0
4	GLC	A	1118	4	11,11,12	0.58	0	15,15,17	1.03	1 (6%)
4	GLC	A	1119	4	11,11,12	0.55	0	15,15,17	1.07	0
4	GLC	A	1120	4	11,11,12	0.56	0	15,15,17	1.52	3 (20%)
4	GLC	A	1121	4	11,11,12	0.68	0	15,15,17	1.30	2 (13%)
5	GOL	A	1122	-	5,5,5	0.39	0	5,5,5	0.47	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	ACT	A	1106	-	-	0/0/0/0	0/0/0/0
3	ACT	A	1107	-	-	0/0/0/0	0/0/0/0
4	GLC	A	1108	4	-	0/2/19/22	0/1/1/1
4	GLC	A	1109	4	-	0/2/19/22	0/1/1/1
4	GLC	A	1110	4	-	0/2/19/22	0/1/1/1
4	GLC	A	1111	4	-	0/2/19/22	0/1/1/1
4	GLC	A	1112	4	-	0/2/19/22	0/1/1/1
4	GLC	A	1113	4	-	0/2/19/22	0/1/1/1
4	GLC	A	1114	4	-	0/2/19/22	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GLC	A	1115	4	-	0/2/19/22	0/1/1/1
4	GLC	A	1116	4	-	0/2/19/22	0/1/1/1
4	GLC	A	1117	4	-	0/2/19/22	0/1/1/1
4	GLC	A	1118	4	-	0/2/19/22	0/1/1/1
4	GLC	A	1119	4	-	0/2/19/22	0/1/1/1
4	GLC	A	1120	4	-	0/2/19/22	0/1/1/1
4	GLC	A	1121	4	-	0/2/19/22	0/1/1/1
5	GOL	A	1122	-	-	0/4/4/4	0/0/0/0

There are no bond length outliers.

The worst 5 of 19 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	1110	GLC	O5-C1-C2	-3.05	106.03	110.78
4	A	1113	GLC	O5-C1-C2	-2.38	107.07	110.78
4	A	1120	GLC	O5-C1-C2	-2.10	107.50	110.78
4	A	1112	GLC	C1-C2-C3	2.02	112.22	109.66
4	A	1114	GLC	O2-C2-C1	2.05	113.30	109.17

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	1052/1102 (95%)	0.13	38 (3%)	42 49	20, 35, 63, 100	0

The worst 5 of 38 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	68	THR	6.0
1	A	93	ALA	4.3
1	A	87	ALA	4.0
1	A	266	ALA	3.8
1	A	83	GLU	3.5

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	GLC	A	1111	11/12	0.74	0.38	68,80,87,90	0
4	GLC	A	1110	11/12	0.77	0.41	68,78,83,85	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	GLC	A	1114	11/12	0.79	0.30	55,66,71,71	0
3	ACT	A	1106	4/4	0.79	0.27	31,46,47,51	0
4	GLC	A	1112	11/12	0.82	0.33	71,81,86,87	0
4	GLC	A	1109	11/12	0.86	0.20	55,62,66,76	0
3	ACT	A	1107	4/4	0.86	0.15	41,44,51,53	0
4	GLC	A	1113	11/12	0.86	0.34	70,74,78,79	0
4	GLC	A	1121	11/12	0.88	0.14	49,56,62,70	0
2	CA	A	1104	1/1	0.88	0.07	65,65,65,65	0
4	GLC	A	1118	11/12	0.89	0.17	73,76,77,80	0
4	GLC	A	1108	11/12	0.89	0.22	48,60,68,70	0
4	GLC	A	1117	11/12	0.89	0.16	59,64,76,76	0
5	GOL	A	1122	6/6	0.90	0.13	34,39,40,42	0
4	GLC	A	1119	11/12	0.90	0.14	60,66,76,79	0
4	GLC	A	1116	11/12	0.90	0.13	36,45,52,57	0
2	CA	A	1103	1/1	0.91	0.07	69,69,69,69	0
2	CA	A	1102	1/1	0.93	0.06	59,59,59,59	0
4	GLC	A	1120	11/12	0.94	0.16	61,65,68,78	0
4	GLC	A	1115	11/12	0.94	0.11	36,45,49,51	0
2	CA	A	1101	1/1	0.99	0.10	25,25,25,25	0
2	CA	A	1105	1/1	1.00	0.10	31,31,31,31	0

6.5 Other polymers

There are no such residues in this entry.