



# wwPDB X-ray Structure Validation Summary Report ⓘ

May 14, 2018 – 04:38 PM EDT

PDB ID : 2A0Z  
Title : The molecular structure of toll-like receptor 3 ligand binding domain  
Authors : Bell, J.K.; Botos, I.; Hall, P.R.; Askins, J.; Shiloach, J.; Segal, D.M.; Davies, D.R.  
Deposited on : 2005-06-17  
Resolution : 2.40 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.3 (157068), CSD as539be (2018)  
Xtriage (Phenix) : 1.13  
EDS : rb-20031172  
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)  
Refmac : 5.8.0158  
CCP4 : 7.0 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20031172


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	122126	3956 (2.40-2.40)
Ramachandran outliers	120053	3897 (2.40-2.40)
Sidechain outliers	120020	3898 (2.40-2.40)
RSRZ outliers	108989	3386 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	705	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	NAG	A	2912	-	-	-	X
2	NAG	A	728	-	-	-	X
2	NAG	A	730	-	-	-	X

## 2 Entry composition

There are 10 unique types of molecules in this entry. The entry contains 6183 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Toll-like receptor 3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	671	Total	C	N	O	S	0	6	0
			5420	3459	919	1023	19			

There are 23 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	704	GLU	-	SEE REMARK 999	UNP O15455
A	705	ASN	-	SEE REMARK 999	UNP O15455
A	706	LEU	-	SEE REMARK 999	UNP O15455
A	707	TYR	-	SEE REMARK 999	UNP O15455
A	708	PHE	-	SEE REMARK 999	UNP O15455
A	709	GLN	-	SEE REMARK 999	UNP O15455
A	710	GLY	-	SEE REMARK 999	UNP O15455
A	711	ASP	-	SEE REMARK 999	UNP O15455
A	712	TYR	-	SEE REMARK 999	UNP O15455
A	713	LYS	-	SEE REMARK 999	UNP O15455
A	714	ASP	-	SEE REMARK 999	UNP O15455
A	715	ASP	-	SEE REMARK 999	UNP O15455
A	716	ASP	-	SEE REMARK 999	UNP O15455
A	717	ASP	-	SEE REMARK 999	UNP O15455
A	718	LYS	-	SEE REMARK 999	UNP O15455
A	719	GLY	-	EXPRESSION TAG	UNP O15455
A	720	SER	-	EXPRESSION TAG	UNP O15455
A	721	HIS	-	EXPRESSION TAG	UNP O15455
A	722	HIS	-	EXPRESSION TAG	UNP O15455
A	723	HIS	-	EXPRESSION TAG	UNP O15455
A	724	HIS	-	EXPRESSION TAG	UNP O15455
A	725	HIS	-	EXPRESSION TAG	UNP O15455
A	726	HIS	-	EXPRESSION TAG	UNP O15455

- Molecule 2 is N-ACETYL-D-GLUCOSAMINE (three-letter code: NAG) (formula: C<sub>8</sub>H<sub>15</sub>NO<sub>6</sub>).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			13	8	1	4		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 3 is BETA-D-MANNOSE (three-letter code: BMA) (formula:  $C_6H_{12}O_6$ ).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			11	6	5		
3	A	1	Total	C	O	0	0
			11	6	5		
3	A	1	Total	C	O	0	0
			11	6	5		

- Molecule 4 is ALPHA-L-FUCOSE (three-letter code: FUC) (formula:  $C_6H_{12}O_5$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			10	6	4		
4	A	1	Total	C	O	0	0
			10	6	4		


- Molecule 5 is ALPHA-D-MANNOSE (three-letter code: MAN) (formula:  $C_6H_{12}O_6$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			11	6	5		


- Molecule 6 is 2-(ACETYLAMINO)-2-DEOXY-A-D-GLUCOPYRANOSE (three-letter code:

NDG) (formula:  $C_8H_{15}NO_6$ ).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	A	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 7 is ALPHA-D-GLUCOSE (three-letter code: GLC) (formula:  $C_6H_{12}O_6$ ).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	C	O	0	0
			12	6	6		
7	A	1	Total	C	O	0	0
			12	6	6		

- Molecule 8 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	A	1	Total	O	S	0	0
			5	4	1		
8	A	1	Total	O	S	0	0
			5	4	1		
8	A	1	Total	O	S	0	0
			5	4	1		

- Molecule 9 is BETA-MERCAPTOETHANOL (three-letter code: BME) (formula: C<sub>2</sub>H<sub>6</sub>OS).





Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
9	A	1	Total	C	O	S	0	0
			4	2	1	1		


- Molecule 10 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
10	A	363	Total	O	0	0
			363	363		

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Toll-like receptor 3



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	141.60Å 160.79Å 122.54Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 2.40 48.75 – 2.42	Depositor EDS
% Data completeness (in resolution range)	91.1 (50.00-2.40) 93.6 (48.75-2.42)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.29 (at 2.42Å)	Xtriage
Refinement program	CNS	Depositor
R, $R_{free}$	0.206 , 0.232 0.202 , (Not available)	Depositor DCC
$R_{free}$ test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	38.7	Xtriage
Anisotropy	0.385	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 51.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	6183	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	45.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.44% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: BMA, NAG, BME, GLC, NDG, FUC, SO4, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.71	1/5533 (0.0%)	0.73	0/7511

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	198	SER	CB-OG	7.28	1.51	1.42

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5420	0	5399	15	1
2	A	279	0	244	6	0
3	A	33	0	28	0	0
4	A	20	0	20	0	0
5	A	11	0	10	0	0
6	A	14	0	13	0	0
7	A	24	0	24	0	0
8	A	15	0	0	0	0
9	A	4	0	5	0	0
10	A	363	0	0	1	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	6183	0	5743	19	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 19 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:422:ILE:H	1:A:445:GLN:HE22	1.23	0.86
2:A:5071:NAG:H61	2:A:5072:NAG:H82	1.72	0.70
1:A:230[A]:ASN:ND2	1:A:257:ASN:HD22	1.95	0.64
1:A:422:ILE:N	1:A:445:GLN:HE22	1.98	0.56
1:A:230[A]:ASN:HD22	1:A:257:ASN:HD22	1.52	0.55

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:575:ASP:OD2	1:A:575:ASP:OD2[4_555]	2.15	0.05

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	674/705 (96%)	634 (94%)	40 (6%)	0	<b>100</b> <b>100</b>

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	631/656 (96%)	622 (99%)	9 (1%)	69 84

5 of 9 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	467	LYS
1	A	613	LYS
1	A	503	GLN
1	A	243	LEU
1	A	498	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 10 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	309	ASN
1	A	311	GLN
1	A	494	ASN
1	A	229[A]	ASN
1	A	445	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry

33 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
9	BME	A	1270	1	3,3,3	0.43	0	1,2,2	0.21	0
2	NAG	A	1961	1	14,14,15	0.67	0	17,19,21	0.68	1 (5%)
2	NAG	A	2521	1,2	14,14,15	0.58	0	17,19,21	0.59	0
2	NAG	A	2522	2	14,14,15	0.57	0	17,19,21	0.65	0
2	NAG	A	2651	1,2	14,14,15	0.46	0	17,19,21	0.69	0
2	NAG	A	2652	3,2	14,14,15	0.51	0	17,19,21	0.69	0
3	BMA	A	2653	2	11,11,12	0.55	0	15,15,17	0.23	0
2	NAG	A	2751	1,2	14,14,15	0.58	0	17,19,21	0.66	0
2	NAG	A	2752	2	14,14,15	0.59	0	17,19,21	0.67	0
2	NAG	A	2911	1,2	14,14,15	0.51	0	17,19,21	0.61	0
2	NAG	A	2912	2	14,14,15	0.56	0	17,19,21	0.59	0
2	NAG	A	3981	1,8,2,4	13,13,15	1.07	1 (7%)	16,17,21	0.95	1 (6%)
2	NAG	A	3982	2	14,14,15	0.57	0	17,19,21	0.72	1 (5%)
4	FUC	A	3986	2	9,10,11	0.61	0	13,14,16	0.40	0
2	NAG	A	4131	1,2	14,14,15	0.50	0	17,19,21	0.66	0
2	NAG	A	4132	3,2	14,14,15	0.54	0	17,19,21	0.66	0
3	BMA	A	4133	3,2,5	11,11,12	0.63	0	15,15,17	0.49	0
5	MAN	A	4134	3	11,11,12	0.69	0	15,15,17	0.58	0
3	BMA	A	4135	3	11,11,12	0.65	0	15,15,17	0.45	0
2	NAG	A	5071	1,2,4	14,14,15	0.59	0	17,19,21	0.77	0
2	NAG	A	5072	2	14,14,15	0.69	0	17,19,21	0.80	0
4	FUC	A	5076	2	9,10,11	0.66	0	13,14,16	0.45	0
2	NAG	A	6361	1,6	14,14,15	1.04	1 (7%)	17,19,21	0.71	0
6	NDG	A	6362	2	14,14,15	0.66	0	17,19,21	0.65	0
2	NAG	A	727	1,2	14,14,15	0.68	0	17,19,21	0.55	0
2	NAG	A	728	2	14,14,15	0.55	0	17,19,21	0.69	0
2	NAG	A	729	1,2	14,14,15	0.55	0	17,19,21	0.66	0
2	NAG	A	730	2	14,14,15	0.65	0	17,19,21	0.51	0
8	SO4	A	750	-	4,4,4	0.31	0	6,6,6	0.12	0
8	SO4	A	751	-	4,4,4	0.32	0	6,6,6	0.08	0
8	SO4	A	752	2	4,4,4	0.44	0	6,6,6	0.24	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
7	GLC	A	753	-	12,12,12	1.65	2 (16%)	17,17,17	1.61	4 (23%)
7	GLC	A	754	-	12,12,12	1.56	1 (8%)	17,17,17	1.56	3 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
9	BME	A	1270	1	-	0/1/1/1	0/0/0/0
2	NAG	A	1961	1	-	0/6/23/26	0/1/1/1
2	NAG	A	2521	1,2	-	0/6/23/26	0/1/1/1
2	NAG	A	2522	2	-	0/6/23/26	0/1/1/1
2	NAG	A	2651	1,2	-	0/6/23/26	0/1/1/1
2	NAG	A	2652	3,2	-	2/6/23/26	0/1/1/1
3	BMA	A	2653	2	-	0/2/19/22	0/1/1/1
2	NAG	A	2751	1,2	-	0/6/23/26	0/1/1/1
2	NAG	A	2752	2	-	0/6/23/26	0/1/1/1
2	NAG	A	2911	1,2	-	0/6/23/26	0/1/1/1
2	NAG	A	2912	2	-	0/6/23/26	0/1/1/1
2	NAG	A	3981	1,8,2,4	-	0/6/19/26	0/1/1/1
2	NAG	A	3982	2	-	1/6/23/26	0/1/1/1
4	FUC	A	3986	2	-	0/0/17/20	0/1/1/1
2	NAG	A	4131	1,2	-	0/6/23/26	0/1/1/1
2	NAG	A	4132	3,2	-	0/6/23/26	0/1/1/1
3	BMA	A	4133	3,2,5	-	0/2/19/22	0/1/1/1
5	MAN	A	4134	3	-	0/2/19/22	1/1/1/1
3	BMA	A	4135	3	-	0/2/19/22	0/1/1/1
2	NAG	A	5071	1,2,4	-	0/6/23/26	0/1/1/1
2	NAG	A	5072	2	-	0/6/23/26	0/1/1/1
4	FUC	A	5076	2	-	0/0/17/20	0/1/1/1
2	NAG	A	6361	1,6	-	0/6/23/26	0/1/1/1
6	NDG	A	6362	2	-	0/6/23/26	0/1/1/1
2	NAG	A	727	1,2	-	2/6/23/26	0/1/1/1
2	NAG	A	728	2	-	0/6/23/26	0/1/1/1
2	NAG	A	729	1,2	-	0/6/23/26	0/1/1/1
2	NAG	A	730	2	-	0/6/23/26	0/1/1/1
8	SO4	A	750	-	-	0/0/0/0	0/0/0/0
8	SO4	A	751	-	-	0/0/0/0	0/0/0/0
8	SO4	A	752	2	-	0/0/0/0	0/0/0/0
7	GLC	A	753	-	-	0/2/22/22	0/1/1/1
7	GLC	A	754	-	-	0/2/22/22	0/1/1/1



All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
7	A	753	GLC	C4-C5	2.11	1.57	1.53
2	A	6361	NAG	C1-C2	2.12	1.55	1.52
2	A	3981	NAG	C1-C2	3.16	1.55	1.51
7	A	754	GLC	O5-C1	4.44	1.54	1.42
7	A	753	GLC	O5-C1	4.49	1.54	1.42

The worst 5 of 10 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	A	753	GLC	C6-C5-C4	-3.45	104.83	112.99
7	A	754	GLC	C6-C5-C4	-3.40	104.95	112.99
7	A	753	GLC	O6-C6-C5	-2.34	103.13	111.29
7	A	754	GLC	O6-C6-C5	-2.11	103.92	111.29
2	A	3981	NAG	C4-C3-C2	-2.09	110.27	112.45

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	3982	NAG	O7-C7-N2-C2
2	A	2652	NAG	C8-C7-N2-C2
2	A	727	NAG	O7-C7-N2-C2
2	A	2652	NAG	O7-C7-N2-C2
2	A	727	NAG	C8-C7-N2-C2

All (1) ring outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	4134	MAN	C1-C2-C3-C4-C5-O5

7 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	2521	NAG	1	0
2	A	3981	NAG	1	0
2	A	5071	NAG	2	0
2	A	5072	NAG	2	0
2	A	727	NAG	1	0
2	A	728	NAG	1	0
2	A	730	NAG	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	671/705 (95%)	-0.09	23 (3%) 45 43	22, 39, 76, 95	0

The worst 5 of 23 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	26	THR	8.3
1	A	693	THR	5.1
1	A	696	CYS	4.9
1	A	694	SER	4.8
1	A	692	ASP	4.6

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	NAG	A	6361	14/15	0.64	0.29	79,85,87,91	0
3	BMA	A	4135	11/12	0.65	0.36	99,99,100,100	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	NAG	A	2912	14/15	0.66	0.43	88,90,92,92	0
7	GLC	A	754	12/12	0.66	0.25	99,100,100,100	0
7	GLC	A	753	12/12	0.73	0.23	54,59,62,67	0
2	NAG	A	1961	14/15	0.74	0.23	74,78,79,81	0
2	NAG	A	730	14/15	0.75	0.51	97,99,100,100	0
3	BMA	A	2653	11/12	0.76	0.27	91,92,93,93	0
8	SO4	A	752	5/5	0.76	0.31	88,92,92,93	0
2	NAG	A	727	14/15	0.77	0.29	88,90,92,95	0
5	MAN	A	4134	11/12	0.77	0.33	97,99,100,100	0
2	NAG	A	2752	14/15	0.78	0.29	80,84,85,86	0
2	NAG	A	728	14/15	0.78	0.43	98,99,100,100	0
2	NAG	A	729	14/15	0.78	0.38	81,85,89,93	0
4	FUC	A	3986	10/11	0.78	0.35	84,85,86,87	0
2	NAG	A	2652	14/15	0.81	0.19	84,86,88,90	0
2	NAG	A	2522	14/15	0.81	0.24	76,79,80,81	0
6	NDG	A	6362	14/15	0.82	0.35	95,97,98,98	0
2	NAG	A	2521	14/15	0.82	0.17	59,65,68,72	0
2	NAG	A	3982	14/15	0.83	0.33	89,91,92,92	0
4	FUC	A	5076	10/11	0.84	0.23	73,76,77,77	0
2	NAG	A	3981	13/15	0.85	0.18	72,75,83,86	0
3	BMA	A	4133	11/12	0.86	0.26	84,91,94,97	0
2	NAG	A	2911	14/15	0.86	0.20	68,73,76,83	0
2	NAG	A	5072	14/15	0.86	0.23	73,77,79,80	0
2	NAG	A	2751	14/15	0.88	0.16	59,63,69,75	0
2	NAG	A	4132	14/15	0.91	0.14	55,59,67,77	0
2	NAG	A	2651	14/15	0.91	0.15	68,74,76,80	0
9	BME	A	1270	4/4	0.94	0.26	57,57,61,64	0
2	NAG	A	5071	14/15	0.95	0.12	37,53,66,68	0
8	SO4	A	751	5/5	0.97	0.12	67,69,69,70	0
2	NAG	A	4131	14/15	0.98	0.14	32,35,39,45	0
8	SO4	A	750	5/5	0.99	0.12	43,44,46,48	0

## 6.5 Other polymers

There are no such residues in this entry.