



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 8, 2018 – 07:43 pm GMT

PDB ID : 4A0G
Title : Structure of bifunctional DAPA aminotransferase-DTB synthetase from *Arabidopsis thaliana* in its apo form.
Authors : Cobessi, D.; Dumas, R.; Pautre, V.; Meinguet, C.; Ferrer, J.L.; Alban, C.
Deposited on : 2011-09-09
Resolution : 2.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.3 (157068), CSD as539be (2018)
Xtriage (Phenix) : 1.13
EDS : trunk30967
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)
Refmac : 5.8.0158
CCP4 : 7.0 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk30967


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	111664	4155 (2.50-2.50)
Clashscore	122126	4827 (2.50-2.50)
Ramachandran outliers	120053	4735 (2.50-2.50)
Sidechain outliers	120020	4737 (2.50-2.50)
RSRZ outliers	108989	4058 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	831	<div> <div>3%</div> <div> <div></div> <div>72%</div> <div>17%</div> <div>•</div> <div>7%</div> </div> </div>
1	B	831	<div> <div>3%</div> <div> <div></div> <div>71%</div> <div>18%</div> <div>•</div> <div>9%</div> </div> </div>
1	C	831	<div> <div>10%</div> <div> <div></div> <div>62%</div> <div>18%</div> <div>•</div> <div>18%</div> </div> </div>
1	D	831	<div> <div>6%</div> <div> <div></div> <div>64%</div> <div>19%</div> <div>•</div> <div>15%</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit crite-

ria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	SO4	A	1814	-	-	X	-
3	SO4	C	1809	-	-	X	-

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 22816 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ADENOSYLMETHIONINE-8-AMINO-7-OXONONANOATE AMINOTRANSFERASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	769	Total	C	N	O	S	0	14	0
			5943	3817	996	1098	32			
1	B	756	Total	C	N	O	S	0	7	0
			5789	3716	972	1070	31			
1	C	681	Total	C	N	O	S	0	4	0
			5160	3311	858	961	30			
1	D	709	Total	C	N	O	S	0	5	0
			5383	3457	901	996	29			

There are 80 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-19	GLY	-	expression tag	UNP B0F481
A	-18	SER	-	expression tag	UNP B0F481
A	-17	SER	-	expression tag	UNP B0F481
A	-16	HIS	-	expression tag	UNP B0F481
A	-15	HIS	-	expression tag	UNP B0F481
A	-14	HIS	-	expression tag	UNP B0F481
A	-13	HIS	-	expression tag	UNP B0F481
A	-12	HIS	-	expression tag	UNP B0F481
A	-11	HIS	-	expression tag	UNP B0F481
A	-10	SER	-	expression tag	UNP B0F481
A	-9	SER	-	expression tag	UNP B0F481
A	-8	GLY	-	expression tag	UNP B0F481
A	-7	LEU	-	expression tag	UNP B0F481
A	-6	VAL	-	expression tag	UNP B0F481
A	-5	PRO	-	expression tag	UNP B0F481
A	-4	ARG	-	expression tag	UNP B0F481
A	-3	GLY	-	expression tag	UNP B0F481
A	-2	SER	-	expression tag	UNP B0F481
A	-1	HIS	-	expression tag	UNP B0F481
A	0	MET	-	expression tag	UNP B0F481

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	-19	GLY	-	expression tag	UNP B0F481
B	-18	SER	-	expression tag	UNP B0F481
B	-17	SER	-	expression tag	UNP B0F481
B	-16	HIS	-	expression tag	UNP B0F481
B	-15	HIS	-	expression tag	UNP B0F481
B	-14	HIS	-	expression tag	UNP B0F481
B	-13	HIS	-	expression tag	UNP B0F481
B	-12	HIS	-	expression tag	UNP B0F481
B	-11	HIS	-	expression tag	UNP B0F481
B	-10	SER	-	expression tag	UNP B0F481
B	-9	SER	-	expression tag	UNP B0F481
B	-8	GLY	-	expression tag	UNP B0F481
B	-7	LEU	-	expression tag	UNP B0F481
B	-6	VAL	-	expression tag	UNP B0F481
B	-5	PRO	-	expression tag	UNP B0F481
B	-4	ARG	-	expression tag	UNP B0F481
B	-3	GLY	-	expression tag	UNP B0F481
B	-2	SER	-	expression tag	UNP B0F481
B	-1	HIS	-	expression tag	UNP B0F481
B	0	MET	-	expression tag	UNP B0F481
C	-19	GLY	-	expression tag	UNP B0F481
C	-18	SER	-	expression tag	UNP B0F481
C	-17	SER	-	expression tag	UNP B0F481
C	-16	HIS	-	expression tag	UNP B0F481
C	-15	HIS	-	expression tag	UNP B0F481
C	-14	HIS	-	expression tag	UNP B0F481
C	-13	HIS	-	expression tag	UNP B0F481
C	-12	HIS	-	expression tag	UNP B0F481
C	-11	HIS	-	expression tag	UNP B0F481
C	-10	SER	-	expression tag	UNP B0F481
C	-9	SER	-	expression tag	UNP B0F481
C	-8	GLY	-	expression tag	UNP B0F481
C	-7	LEU	-	expression tag	UNP B0F481
C	-6	VAL	-	expression tag	UNP B0F481
C	-5	PRO	-	expression tag	UNP B0F481
C	-4	ARG	-	expression tag	UNP B0F481
C	-3	GLY	-	expression tag	UNP B0F481
C	-2	SER	-	expression tag	UNP B0F481
C	-1	HIS	-	expression tag	UNP B0F481
C	0	MET	-	expression tag	UNP B0F481
D	-19	GLY	-	expression tag	UNP B0F481
D	-18	SER	-	expression tag	UNP B0F481

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	-17	SER	-	expression tag	UNP B0F481
D	-16	HIS	-	expression tag	UNP B0F481
D	-15	HIS	-	expression tag	UNP B0F481
D	-14	HIS	-	expression tag	UNP B0F481
D	-13	HIS	-	expression tag	UNP B0F481
D	-12	HIS	-	expression tag	UNP B0F481
D	-11	HIS	-	expression tag	UNP B0F481
D	-10	SER	-	expression tag	UNP B0F481
D	-9	SER	-	expression tag	UNP B0F481
D	-8	GLY	-	expression tag	UNP B0F481
D	-7	LEU	-	expression tag	UNP B0F481
D	-6	VAL	-	expression tag	UNP B0F481
D	-5	PRO	-	expression tag	UNP B0F481
D	-4	ARG	-	expression tag	UNP B0F481
D	-3	GLY	-	expression tag	UNP B0F481
D	-2	SER	-	expression tag	UNP B0F481
D	-1	HIS	-	expression tag	UNP B0F481
D	0	MET	-	expression tag	UNP B0F481

- Molecule 2 is PYRIDOXAL-5'-PHOSPHATE (three-letter code: PLP) (formula: $C_8H_{10}NO_6P$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			15	8	1	5	1		
2	B	1	Total	C	N	O	P	0	0
			15	8	1	5	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	C	1	Total	C	N	O	P	0	0
			15	8	1	5	1		
2	D	1	Total	C	N	O	P	0	0
			15	8	1	5	1		

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	C	1	Total	O	S	0	0
			5	4	1		
3	C	1	Total	O	S	0	0
			5	4	1		
3	C	1	Total	O	S	0	0
			5	4	1		
3	D	1	Total	O	S	0	0
			5	4	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	D	1	Total	O	S	0	0
			5	4	1		

- Molecule 4 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	Mg	0	0
			1	1		
4	D	1	Total	Mg	0	0
			1	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	185	Total	O	0	0
			185	185		
5	B	123	Total	O	0	0
			123	123		
5	C	57	Total	O	0	0
			57	57		
5	D	64	Total	O	0	0
			64	64		


3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: ADENOSYLMETHIONINE-8-AMINO-7-OXONONANOATE AMINOTRANSFERASE




• Molecule 1: ADENOSYLMETHIONINE-8-AMINO-7-OXONONANOATE AMINOTRANSFERASE





• Molecule 1: ADENOSYLMETHIONINE-8-AMINO-7-OXONONANOATE AMINOTRANSFERASE



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	79.44Å 80.07Å 136.94Å 99.96° 107.12° 97.25°	Depositor
Resolution (Å)	39.57 – 2.50 48.13 – 2.50	Depositor EDS
% Data completeness (in resolution range)	88.3 (39.57-2.50) 88.3 (48.13-2.50)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.28 (at 2.51Å)	Xtriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
R, R_{free}	0.177 , 0.239 0.169 , 0.230	Depositor DCC
R_{free} test set	4753 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	40.3	Xtriage
Anisotropy	0.067	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 60.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.010 for k,h,-h-k-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	22816	wwPDB-VP
Average B, all atoms (Å ²)	47.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.96% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, SO4, PLP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.60	4/6119 (0.1%)	0.68	5/8324 (0.1%)
1	B	0.61	4/5942 (0.1%)	0.70	5/8081 (0.1%)
1	C	0.48	2/5283 (0.0%)	0.66	3/7189 (0.0%)
1	D	0.48	2/5516 (0.0%)	0.69	6/7505 (0.1%)
All	All	0.55	12/22860 (0.1%)	0.68	19/31099 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	1
All	All	0	2

The worst 5 of 12 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	388	TYR	CE1-CZ	-13.53	1.21	1.38
1	A	388	TYR	CE2-CZ	-12.80	1.22	1.38
1	B	388	TYR	CE2-CZ	-12.78	1.22	1.38
1	B	388	TYR	CE1-CZ	-12.54	1.22	1.38
1	B	388	TYR	CG-CD2	-11.50	1.24	1.39

The worst 5 of 19 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	796	ARG	NE-CZ-NH2	21.46	131.03	120.30
1	C	796	ARG	NE-CZ-NH2	20.18	130.39	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	796	ARG	NE-CZ-NH1	-14.34	113.13	120.30
1	C	796	ARG	NE-CZ-NH1	-13.95	113.33	120.30
1	D	263	ARG	CG-CD-NE	10.22	133.27	111.80

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	374	PRO	Peptide
1	B	374	PRO	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5943	0	5868	161	1
1	B	5789	0	5695	149	1
1	C	5160	0	5056	145	0
1	D	5383	0	5280	147	0
2	A	15	0	6	2	0
2	B	15	0	6	2	0
2	C	15	0	6	3	0
2	D	15	0	6	1	0
3	A	15	0	0	2	0
3	B	10	0	0	0	0
3	C	15	0	0	3	0
3	D	10	0	0	0	0
4	B	1	0	0	0	0
4	D	1	0	0	0	0
5	A	185	0	0	6	0
5	B	123	0	0	3	0
5	C	57	0	0	0	0
5	D	64	0	0	1	0
All	All	22816	0	21923	558	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

The worst 5 of 558 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:263:ARG:NH2	1:D:697:TRP:CZ3	2.08	1.22
1:C:697:TRP:CZ3	1:D:263:ARG:NH2	2.11	1.17
1:C:263:ARG:NH2	1:D:697:TRP:CH2	2.19	1.11
1:A:43:GLN:O	1:A:45:SER:N	1.89	1.03
1:C:441:MET:HE1	1:C:658:ALA:HB2	1.48	0.95

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:807:GLU:OE2	1:B:769:GLU:OE1[1_455]	2.11	0.09

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	773/831 (93%)	724 (94%)	37 (5%)	12 (2%)	11	19
1	B	747/831 (90%)	699 (94%)	37 (5%)	11 (2%)	11	20
1	C	665/831 (80%)	621 (93%)	37 (6%)	7 (1%)	16	28
1	D	694/831 (84%)	644 (93%)	40 (6%)	10 (1%)	12	22
All	All	2879/3324 (87%)	2688 (93%)	151 (5%)	40 (1%)	12	22

5 of 40 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	355	ASP
1	A	357	SER
1	A	709	SER
1	B	355	ASP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	357	SER

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	637/704 (90%)	603 (95%)	34 (5%)	25	46
1	B	617/704 (88%)	585 (95%)	32 (5%)	25	47
1	C	549/704 (78%)	526 (96%)	23 (4%)	32	57
1	D	569/704 (81%)	546 (96%)	23 (4%)	34	60
All	All	2372/2816 (84%)	2260 (95%)	112 (5%)	31	52

5 of 112 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	466	VAL
1	B	707[B]	ILE
1	D	556	SER
1	B	519	ILE
1	B	655	VAL

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 13 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	89	ASN
1	C	451	ASN
1	D	89	ASN
1	C	14	HIS
1	D	14	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 16 ligands modelled in this entry, 2 are monoatomic - leaving 14 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	PLP	A	1644	1	15,15,16	1.84	3 (20%)	20,22,23	1.50	3 (15%)
3	SO4	A	1812	-	4,4,4	0.21	0	6,6,6	0.36	0
3	SO4	A	1813	-	4,4,4	0.23	0	6,6,6	0.25	0
3	SO4	A	1814	-	4,4,4	0.19	0	6,6,6	0.22	0
2	PLP	B	1644	1	15,15,16	1.76	2 (13%)	20,22,23	1.68	5 (25%)
3	SO4	B	1812	-	4,4,4	0.24	0	6,6,6	0.34	0
3	SO4	B	1813	4	4,4,4	0.30	0	6,6,6	0.60	0
2	PLP	C	1644	1	15,15,16	1.87	2 (13%)	20,22,23	1.78	5 (25%)
3	SO4	C	1807	-	4,4,4	0.16	0	6,6,6	0.17	0
3	SO4	C	1808	-	4,4,4	0.20	0	6,6,6	0.33	0
3	SO4	C	1809	-	4,4,4	0.25	0	6,6,6	0.46	0
2	PLP	D	1644	1	15,15,16	1.93	3 (20%)	20,22,23	1.69	2 (10%)
3	SO4	D	1808	-	4,4,4	0.15	0	6,6,6	0.32	0
3	SO4	D	1809	4	4,4,4	0.16	0	6,6,6	0.58	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PLP	A	1644	1	-	0/6/6/8	0/1/1/1
3	SO4	A	1812	-	-	0/0/0/0	0/0/0/0
3	SO4	A	1813	-	-	0/0/0/0	0/0/0/0
3	SO4	A	1814	-	-	0/0/0/0	0/0/0/0
2	PLP	B	1644	1	-	0/6/6/8	0/1/1/1
3	SO4	B	1812	-	-	0/0/0/0	0/0/0/0
3	SO4	B	1813	4	-	0/0/0/0	0/0/0/0
2	PLP	C	1644	1	-	0/6/6/8	0/1/1/1
3	SO4	C	1807	-	-	0/0/0/0	0/0/0/0
3	SO4	C	1808	-	-	0/0/0/0	0/0/0/0
3	SO4	C	1809	-	-	0/0/0/0	0/0/0/0
2	PLP	D	1644	1	-	0/6/6/8	0/1/1/1
3	SO4	D	1808	-	-	0/0/0/0	0/0/0/0
3	SO4	D	1809	4	-	0/0/0/0	0/0/0/0

The worst 5 of 10 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	1644	PLP	O3-C3	-5.83	1.23	1.37
2	C	1644	PLP	O3-C3	-5.78	1.23	1.37
2	A	1644	PLP	O3-C3	-5.71	1.23	1.37
2	B	1644	PLP	O3-C3	-5.57	1.24	1.37
2	D	1644	PLP	C6-N1	2.04	1.38	1.34

The worst 5 of 15 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	1644	PLP	C5-C6-N1	-2.57	119.49	123.83
2	A	1644	PLP	C5-C6-N1	-2.46	119.66	123.83
2	B	1644	PLP	C5-C6-N1	-2.19	120.13	123.83
2	C	1644	PLP	O4P-P-O1P	-2.15	100.43	106.47
2	D	1644	PLP	C5-C6-N1	-2.04	120.38	123.83

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

7 monomers are involved in 13 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1644	PLP	2	0
3	A	1814	SO4	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	1644	PLP	2	0
2	C	1644	PLP	3	0
3	C	1808	SO4	1	0
3	C	1809	SO4	2	0
2	D	1644	PLP	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2			OWAB(Å ²)	Q < 0.9
1	A	769/831 (92%)	-0.02	25 (3%)	46	50	19, 36, 74, 111	0
1	B	756/831 (90%)	-0.04	24 (3%)	47	51	19, 36, 74, 111	0
1	C	681/831 (81%)	0.43	82 (12%)	4	4	27, 53, 102, 144	0
1	D	709/831 (85%)	0.18	54 (7%)	14	14	24, 51, 90, 118	0
All	All	2915/3324 (87%)	0.13	185 (6%)	20	21	19, 43, 88, 144	0

The worst 5 of 185 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	446	PHE	6.7
1	C	448	VAL	5.9
1	C	708	THR	5.7
1	C	806	GLY	5.5
1	C	470	ARG	5.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	SO4	A	1814	5/5	0.92	0.14	56,58,73,86	5
3	SO4	C	1809	5/5	0.94	0.14	25,31,67,79	5
3	SO4	D	1809	5/5	0.94	0.13	17,31,74,99	5
4	MG	D	1810	1/1	0.95	0.06	45,45,45,45	0
3	SO4	A	1812	5/5	0.95	0.13	35,42,66,71	5
3	SO4	B	1813	5/5	0.95	0.10	32,38,71,81	5
3	SO4	C	1807	5/5	0.96	0.09	30,45,66,83	5
4	MG	B	1814	1/1	0.97	0.18	58,58,58,58	1
2	PLP	C	1644	15/16	0.98	0.13	42,56,77,91	0
2	PLP	D	1644	15/16	0.98	0.15	27,51,59,64	0
3	SO4	D	1808	5/5	0.98	0.20	36,38,46,58	5
2	PLP	A	1644	15/16	0.99	0.21	14,27,34,35	0
3	SO4	C	1808	5/5	0.99	0.18	37,47,51,66	5
3	SO4	B	1812	5/5	0.99	0.15	29,37,52,57	5
3	SO4	A	1813	5/5	0.99	0.20	39,40,59,63	0
2	PLP	B	1644	15/16	0.99	0.19	19,25,38,39	0

6.5 Other polymers [i](#)

There are no such residues in this entry.