



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2018 – 12:49 am GMT

PDB ID : 5A08  
Title : X-ray structure of the mannosyltransferase Ktr4p from *S. cerevisiae*  
Authors : Possner, D.D.D.; Guy, J.E.  
Deposited on : 2015-04-17  
Resolution : 2.21 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.3 (157068), CSD as539be (2018)  
Xtriage (Phenix) : 1.13  
EDS : trunk30967  
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)  
Refmac : 5.8.0158  
CCP4 : 7.0 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk30967


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.21 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	111664	5192 (2.24-2.20)
Clashscore	122126	5911 (2.24-2.20)
Ramachandran outliers	120053	5828 (2.24-2.20)
Sidechain outliers	120020	5829 (2.24-2.20)
RSRZ outliers	108989	5078 (2.24-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	434	<div> <div style="width: 100%; height: 10px; position: relative;"> <div style="position: absolute; top: -10px; left: 0; width: 100%; text-align: center;">%</div> <div style="position: absolute; top: 0; left: 0; width: 100%; height: 10px;"> <div style="width: 83%; height: 10px; background-color: red;"></div> <div style="width: 83%; height: 10px; background-color: orange;"></div> <div style="width: 83%; height: 10px; background-color: yellow;"></div> <div style="width: 83%; height: 10px; background-color: green;"></div> <div style="width: 83%; height: 10px; background-color: grey;"></div> </div> <div style="position: absolute; bottom: -10px; left: 0; width: 100%; text-align: center;"> <div style="width: 83%;"></div> <div style="width: 8%;"></div> <div style="width: 9%;"></div> </div> </div> </div>
1	B	434	<div> <div style="width: 100%; height: 10px; position: relative;"> <div style="position: absolute; top: -10px; left: 0; width: 100%; text-align: center;">%</div> <div style="position: absolute; top: 0; left: 0; width: 100%; height: 10px;"> <div style="width: 80%; height: 10px; background-color: red;"></div> <div style="width: 80%; height: 10px; background-color: orange;"></div> <div style="width: 80%; height: 10px; background-color: yellow;"></div> <div style="width: 80%; height: 10px; background-color: green;"></div> <div style="width: 80%; height: 10px; background-color: grey;"></div> </div> <div style="position: absolute; bottom: -10px; left: 0; width: 100%; text-align: center;"> <div style="width: 80%;"></div> <div style="width: 9%;"></div> <div style="width: 11%;"></div> </div> </div> </div>

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 7081 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called PROBABLE MANNOSYLTRANSFERASE KTR4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	394	Total	C	N	O	S	0	4	0
			3319	2116	535	650	18			
1	B	387	Total	C	N	O	S	0	3	0
			3253	2074	524	637	18			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	31	SER	-	expression tag	UNP P38131
A	32	MET	-	expression tag	UNP P38131
B	31	SER	-	expression tag	UNP P38131
B	32	MET	-	expression tag	UNP P38131

- Molecule 2 is ACETATE ION (three-letter code: ACT) (formula: C<sub>2</sub>H<sub>3</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C O 4 2 2	0	0
2	B	1	Total C O 4 2 2	0	0

- Molecule 3 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	B	3	Total Ca 3 3	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	245	Total O 245 245	0	0
4	B	253	Total O 253 253	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: PROBABLE MANNOSYLTRANSFERASE KTR4



#### • Molecule 1: PROBABLE MANNOSYLTRANSFERASE KTR4



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	60.20Å 102.37Å 156.91Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	85.74 – 2.21 49.27 – 2.21	Depositor EDS
% Data completeness (in resolution range)	99.7 (85.74-2.21) 99.7 (49.27-2.21)	Depositor EDS
$R_{merge}$	0.13	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.03 (at 2.20Å)	Xtriage
Refinement program	REFMAC 5.8.0073	Depositor
R, $R_{free}$	0.163 , 0.203 0.171 , 0.207	Depositor DCC
$R_{free}$ test set	2420 reflections (4.89%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	22.9	Xtriage
Anisotropy	0.096	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 43.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	7081	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.13% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CA, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.88	1/3429 (0.0%)	0.84	6/4654 (0.1%)
1	B	0.95	1/3358 (0.0%)	0.91	10/4557 (0.2%)
All	All	0.91	2/6787 (0.0%)	0.88	16/9211 (0.2%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	362	GLU	CD-OE1	-5.78	1.19	1.25
1	A	458	GLU	CD-OE1	5.59	1.31	1.25

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	397	ASP	CB-CG-OD2	-7.03	111.97	118.30
1	A	453	ARG	NE-CZ-NH1	6.99	123.80	120.30
1	B	453	ARG	NE-CZ-NH1	6.73	123.67	120.30
1	A	124	ARG	NE-CZ-NH2	-6.49	117.05	120.30
1	B	357	ASP	CB-CG-OD2	-6.09	112.82	118.30
1	A	124	ARG	NE-CZ-NH1	5.88	123.24	120.30
1	B	120	ASP	CB-CG-OD1	-5.87	113.02	118.30
1	B	214	ARG	NE-CZ-NH2	-5.79	117.41	120.30
1	B	357	ASP	CB-CG-OD1	5.56	123.31	118.30
1	B	124	ARG	NE-CZ-NH2	-5.50	117.55	120.30
1	A	307	MET	CG-SD-CE	5.41	108.86	100.20
1	B	124	ARG	NE-CZ-NH1	5.40	123.00	120.30
1	A	397	ASP	CB-CG-OD1	5.33	123.10	118.30
1	B	232	VAL	N-CA-CB	5.30	123.17	111.50
1	A	453	ARG	NE-CZ-NH2	-5.11	117.75	120.30
1	B	453	ARG	NE-CZ-NH2	-5.09	117.76	120.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3319	0	3071	20	0
1	B	3253	0	3004	29	0
2	A	4	0	3	0	0
2	B	4	0	3	0	0
3	B	3	0	0	0	0
4	A	245	0	0	6	0
4	B	253	0	0	7	0
All	All	7081	0	6081	49	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (49) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:222:MET:SD	1:B:381:ARG:NE	2.52	0.82
1:A:236:HIS:HD2	1:A:377:PHE:O	1.72	0.73
1:A:198:THR:O	1:A:249:HIS:HE1	1.73	0.71
1:B:182:ASP:OD2	4:B:2092:HOH:O	2.11	0.69
1:A:195:GLU:OE1	4:A:2084:HOH:O	2.12	0.66
1:A:357:ASP:OD1	4:A:2187:HOH:O	2.16	0.61
1:A:148:ASP:CG	1:A:264[B]:ASN:OD1	2.38	0.61
1:B:258:VAL:HG22	1:B:354[A]:VAL:HG13	1.83	0.60
1:B:103:LYS:CD	4:B:2239:HOH:O	2.50	0.58
1:B:116:GLU:HB2	1:B:117:PRO:CD	2.34	0.58
1:B:357:ASP:OD1	4:B:2195:HOH:O	2.18	0.56
1:B:116:GLU:HB2	1:B:117:PRO:HD2	1.88	0.56
1:B:103:LYS:HD3	4:B:2239:HOH:O	2.05	0.55
1:A:116:GLU:HB2	1:A:117:PRO:CD	2.36	0.55
1:A:244:LYS:NZ	4:A:2125:HOH:O	2.38	0.55
1:A:116:GLU:HB2	1:A:117:PRO:HD2	1.91	0.52

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:121[A]:ASN:HB2	1:B:281:LYS:HG2	1.92	0.52
1:B:336:TYR:CZ	1:B:341:GLY:HA2	2.45	0.51
1:B:222:MET:SD	1:B:381:ARG:HG3	2.51	0.51
1:A:336:TYR:CZ	1:A:341:GLY:HA2	2.46	0.51
1:A:249:HIS:HD2	1:A:251:LEU:H	1.58	0.51
1:B:103:LYS:HD2	4:B:2239:HOH:O	2.11	0.51
1:B:262:GLU:HG3	1:B:350:ASN:HB2	1.94	0.50
1:B:248:HIS:HE1	1:B:365:GLU:OE2	1.94	0.49
1:A:430:ASN:ND2	4:A:2217:HOH:O	2.46	0.49
1:B:222:MET:SD	1:B:381:ARG:CD	3.01	0.48
1:B:248:HIS:CE1	1:B:365:GLU:OE2	2.67	0.47
1:A:132:LYS:N	1:A:132:LYS:HD2	2.29	0.47
1:B:132:LYS:N	1:B:132:LYS:HD2	2.30	0.47
1:A:172:ASN:HB2	1:A:176:PHE:CZ	2.50	0.47
1:A:236:HIS:CD2	1:A:377:PHE:O	2.59	0.47
1:A:262:GLU:HG3	1:A:350:ASN:HB2	1.97	0.45
1:B:211:ASP:OD1	1:B:213:GLU:HG2	2.17	0.45
1:B:121[A]:ASN:ND2	1:B:281:LYS:HG3	2.31	0.45
1:B:103:LYS:CE	4:B:2019:HOH:O	2.65	0.44
1:B:258:VAL:HG22	1:B:354[A]:VAL:CG1	2.46	0.43
1:A:147:GLU:OE2	4:A:2063:HOH:O	2.21	0.43
1:A:220:ASP:OD2	1:A:224:LYS:NZ	2.51	0.43
1:B:121[A]:ASN:CB	1:B:281:LYS:HG2	2.49	0.42
1:B:147:GLU:HG2	4:B:2068:HOH:O	2.19	0.42
1:B:232:VAL:HG23	1:B:235:TYR:H	1.84	0.42
1:B:286:TYR:CE1	1:B:354[B]:VAL:HG22	2.54	0.42
1:B:463:MET:HG2	1:B:463:MET:O	2.19	0.42
1:A:230:ALA:O	1:A:236:HIS:HE1	2.02	0.42
1:B:172:ASN:HB2	1:B:176:PHE:CZ	2.55	0.42
1:B:275:VAL:O	1:B:279:MET:HG2	2.20	0.42
1:B:316:TYR:OH	1:B:362:GLU:OE2	2.36	0.42
1:A:93:PHE:CE1	1:A:330:SER:HA	2.55	0.41
1:A:147:GLU:HG2	4:A:2063:HOH:O	2.19	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	396/434 (91%)	388 (98%)	8 (2%)	0	100	100
1	B	386/434 (89%)	373 (97%)	13 (3%)	0	100	100
All	All	782/868 (90%)	761 (97%)	21 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	369/400 (92%)	366 (99%)	3 (1%)	83	90
1	B	362/400 (90%)	358 (99%)	4 (1%)	76	86
All	All	731/800 (91%)	724 (99%)	7 (1%)	78	87

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	190	LYS
1	A	242	TYR
1	A	388	ARG
1	B	242	TYR
1	B	244	LYS
1	B	297	GLN
1	B	388	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	203	ASN
1	A	236	HIS
1	A	249	HIS
1	A	332	ASN
1	A	335	ASN
1	A	430	ASN
1	B	155	GLN
1	B	248	HIS
1	B	335	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 3 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	ACT	A	1465	-	1,3,3	2.72	1 (100%)	0,3,3	0.00	-
2	ACT	B	1465	-	1,3,3	1.99	0	0,3,3	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ACT	A	1465	-	-	0/0/0/0	0/0/0/0
2	ACT	B	1465	-	-	0/0/0/0	0/0/0/0

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1465	ACT	CH3-C	2.72	1.52	1.48

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	394/434 (90%)	-0.45	6 (1%) 73 72	13, 25, 51, 71	0
1	B	387/434 (89%)	-0.57	3 (0%) 86 85	12, 19, 41, 81	0
All	All	781/868 (89%)	-0.51	9 (1%) 79 77	12, 22, 48, 81	0

All (9) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	71	HIS	4.1
1	B	71	HIS	3.5
1	B	231	GLU	3.3
1	A	73	ALA	2.8
1	A	226	ASN	2.8
1	B	222	MET	2.8
1	A	314	PRO	2.2
1	A	315	ASN	2.1
1	A	316	TYR	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	ACT	B	1465	4/4	0.97	0.10	21,22,22,24	0
2	ACT	A	1465	4/4	0.98	0.12	30,31,31,32	0
3	CA	B	1466	1/1	0.98	0.06	56,56,56,56	0
3	CA	B	1467	1/1	0.99	0.07	20,20,20,20	0
3	CA	B	1468	1/1	1.00	0.14	5,5,5,5	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.