



wwPDB X-ray Structure Validation Summary Report ⓘ

Jan 21, 2019 – 09:56 AM EST

PDB ID : 6A0R
Title : Homoserine dehydrogenase from *Thermus thermophilus* HB8 unliganded form
Authors : Akai, S.; Ikushiro, H.; Sawai, T.; Yano, T.; Kamiya, N.; Miyahara, I.
Deposited on : 2018-06-06
Resolution : 1.83 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.3 (157068), CSD as539be (2018)
Xtriage (Phenix) : 1.13
EDS : rb-20031633
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)
Refmac : 5.8.0158
CCP4 : 7.0 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20031633


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.83 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	111664	3313 (1.86-1.82)
Clashscore	122126	3530 (1.86-1.82)
Ramachandran outliers	120053	3495 (1.86-1.82)
Sidechain outliers	120020	3496 (1.86-1.82)
RSRZ outliers	108989	3265 (1.86-1.82)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	332	<div> <div>%</div> <div> <div></div> <div>86%</div> <div>11%</div> <div>.</div> </div> </div>
1	B	332	<div> <div>3%</div> <div> <div></div> <div>82%</div> <div>15%</div> <div>..</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	FMT	B	411	-	-	X	-

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 6155 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Homoserine dehydrogenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	B	331	Total	C	N	O	S	0	15	0
			2609	1668	461	476	4			
1	A	332	Total	C	N	O	S	0	17	0
			2644	1687	469	484	4			

- Molecule 2 is SODIUM ION (three-letter code: NA) (formula: Na).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Na	0	0
			1	1		
2	A	1	Total	Na	0	0
			1	1		

- Molecule 3 is FORMIC ACID (three-letter code: FMT) (formula: CH₂O₂).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	B	1	Total C O 3 1 2	0	0
3	B	1	Total C O 3 1 2	0	0
3	B	1	Total C O 3 1 2	0	0
3	B	1	Total C O 3 1 2	0	0
3	B	1	Total C O 3 1 2	0	0
3	B	1	Total C O 3 1 2	0	0
3	B	1	Total C O 3 1 2	0	0
3	B	1	Total C O 6 2 4	0	1
3	B	1	Total C O 3 1 2	0	0
3	B	1	Total C O 3 1 2	0	0
3	A	1	Total C O 3 1 2	0	0
3	A	1	Total C O 3 1 2	0	0
3	A	1	Total C O 3 1 2	0	0
3	A	1	Total C O 3 1 2	0	0
3	A	1	Total C O 3 1 2	0	0
3	A	1	Total C O 3 1 2	0	0
3	A	1	Total C O 3 1 2	0	0

- Molecule 4 is 3-CYCLOHEXYL-1-PROPYLSULFONIC ACID (three-letter code: CXS) (formula: C₉H₁₉NO₃S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	B	1	Total	C	N	O	S	0	0
			14	9	1	3	1		
4	B	1	Total	C	N	O	S	0	1
			14	9	1	3	1		

- Molecule 5 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	B	1	Total	C	O	0	1
			12	6	6		
5	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 6 is UNKNOWN LIGAND (three-letter code: UNL) (formula:).

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	B	1	Total	C	O	0	0
			13	11	2		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	B	379	Total	O	0	31
			393	393		
7	A	380	Total	O	0	25
			394	394		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Homoserine dehydrogenase



• Molecule 1: Homoserine dehydrogenase



4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	119.20Å 119.20Å 144.55Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	20.00 – 1.83 19.87 – 1.83	Depositor EDS
% Data completeness (in resolution range)	99.8 (20.00-1.83) 99.9 (19.87-1.83)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	6.27 (at 1.82Å)	Xtriage
Refinement program	REFMAC 5.8.0135	Depositor
R, R_{free}	0.148 , 0.177 0.160 , 0.190	Depositor DCC
R_{free} test set	5231 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å ²)	23.1	Xtriage
Anisotropy	0.019	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 61.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.015 for -h,-k,l	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	6155	wwPDB-VP
Average B, all atoms (Å ²)	29.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.05% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, FMT, CXS, UNL, NA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.36	5/2703 (0.2%)	1.26	16/3674 (0.4%)
1	B	1.48	14/2674 (0.5%)	1.32	21/3637 (0.6%)
All	All	1.42	19/5377 (0.4%)	1.29	37/7311 (0.5%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
1	B	0	4
All	All	0	6

The worst 5 of 19 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	55	GLU	CD-OE1	8.15	1.34	1.25
1	B	114	GLU	CD-OE1	-7.90	1.17	1.25
1	B	2	GLU	CB-CG	-7.11	1.38	1.52
1	A	190	ILE	N-CA	-6.84	1.32	1.46
1	B	212	GLU	CD-OE1	6.74	1.33	1.25

The worst 5 of 37 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	63	ASP	CB-CG-OD1	9.86	127.18	118.30
1	A	250	ARG	NE-CZ-NH2	8.59	124.59	120.30
1	A	277[A]	ARG	NE-CZ-NH1	7.17	123.88	120.30
1	A	277[B]	ARG	NE-CZ-NH1	7.17	123.88	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	237	ARG	NE-CZ-NH1	6.67	123.63	120.30

There are no chirality outliers.

5 of 6 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	266	ARG	Mainchain
1	B	324	GLU	Peptide
1	B	325	GLY	Peptide
1	B	328	TRP	Peptide
1	B	329	PRO	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2644	0	2668	23	0
1	B	2609	0	2641	36	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	21	0	7	0	0
3	B	33	0	12	4	0
4	B	28	0	36	0	0
5	A	6	0	8	1	0
5	B	12	0	16	1	0
6	B	13	0	0	0	0
7	A	394	0	0	2	0
7	B	393	0	0	6	0
All	All	6155	0	5388	60	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 60 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:275[B]:ARG:HG2	1:B:281[B]:GLU:HG2	1.23	1.16
1:B:250[A]:ARG:HD2	7:B:514[A]:HOH:O	1.49	1.11
1:A:275[B]:ARG:HG2	1:A:281[B]:GLU:HG2	1.56	0.85
1:B:275[B]:ARG:HG2	1:B:281[B]:GLU:CG	2.12	0.77
1:B:275[B]:ARG:CG	1:B:281[B]:GLU:HG2	2.10	0.73

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	347/332 (104%)	332 (96%)	13 (4%)	2 (1%)	27	13
1	B	344/332 (104%)	328 (95%)	12 (4%)	4 (1%)	14	4
All	All	691/664 (104%)	660 (96%)	25 (4%)	6 (1%)	21	6

5 of 6 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	325	GLY
1	B	328	TRP
1	B	329	PRO
1	B	330	GLY
1	A	267[A]	ALA

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	261/248 (105%)	250 (96%)	11 (4%)	32	14
1	B	258/248 (104%)	253 (98%)	5 (2%)	60	45
All	All	519/496 (105%)	503 (97%)	16 (3%)	49	24

5 of 16 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	183	PRO
1	A	216[A]	GLN
1	A	277[B]	ARG
1	A	99	LYS
1	A	319	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 26 ligands modelled in this entry, 1 is unknown and 2 are monoatomic - leaving 23 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	FMT	A	402	-	0,2,2	0.00	-	0,1,1	0.00	-
3	FMT	A	403	-	0,2,2	0.00	-	0,1,1	0.00	-
3	FMT	A	404	-	0,2,2	0.00	-	0,1,1	0.00	-
3	FMT	A	405	-	0,2,2	0.00	-	0,1,1	0.00	-
3	FMT	A	406	-	0,2,2	0.00	-	0,1,1	0.00	-
3	FMT	A	407	-	0,2,2	0.00	-	0,1,1	0.00	-
3	FMT	A	408	-	0,2,2	0.00	-	0,1,1	0.00	-
5	GOL	A	409	-	5,5,5	1.34	1 (20%)	5,5,5	1.63	2 (40%)
3	FMT	B	402	-	0,2,2	0.00	-	0,1,1	0.00	-
3	FMT	B	403	-	0,2,2	0.00	-	0,1,1	0.00	-
3	FMT	B	404	-	0,2,2	0.00	-	0,1,1	0.00	-
3	FMT	B	405	-	0,2,2	0.00	-	0,1,1	0.00	-
3	FMT	B	406	-	0,2,2	0.00	-	0,1,1	0.00	-
3	FMT	B	407	-	0,2,2	0.00	-	0,1,1	0.00	-
3	FMT	B	408	-	0,2,2	0.00	-	0,1,1	0.00	-
3	FMT	B	409[A]	-	0,2,2	0.00	-	0,1,1	0.00	-
3	FMT	B	409[B]	-	0,2,2	0.00	-	0,1,1	0.00	-
3	FMT	B	410	-	0,2,2	0.00	-	0,1,1	0.00	-
3	FMT	B	411	-	0,2,2	0.00	-	0,1,1	0.00	-
4	CXS	B	412	-	14,14,14	2.63	2 (14%)	17,18,18	2.02	5 (29%)
4	CXS	B	413[A]	-	14,14,14	2.71	2 (14%)	17,18,18	2.43	9 (52%)
5	GOL	B	414[A]	-	5,5,5	0.58	0	5,5,5	1.89	1 (20%)
5	GOL	B	414[B]	-	5,5,5	0.82	0	5,5,5	1.33	1 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	FMT	A	402	-	-	0/0/0/0	0/0/0/0
3	FMT	A	403	-	-	0/0/0/0	0/0/0/0
3	FMT	A	404	-	-	0/0/0/0	0/0/0/0
3	FMT	A	405	-	-	0/0/0/0	0/0/0/0
3	FMT	A	406	-	-	0/0/0/0	0/0/0/0
3	FMT	A	407	-	-	0/0/0/0	0/0/0/0
3	FMT	A	408	-	-	0/0/0/0	0/0/0/0
5	GOL	A	409	-	-	0/4/4/4	0/0/0/0
3	FMT	B	402	-	-	0/0/0/0	0/0/0/0
3	FMT	B	403	-	-	0/0/0/0	0/0/0/0
3	FMT	B	404	-	-	0/0/0/0	0/0/0/0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	FMT	B	405	-	-	0/0/0/0	0/0/0/0
3	FMT	B	406	-	-	0/0/0/0	0/0/0/0
3	FMT	B	407	-	-	0/0/0/0	0/0/0/0
3	FMT	B	408	-	-	0/0/0/0	0/0/0/0
3	FMT	B	409[A]	-	-	0/0/0/0	0/0/0/0
3	FMT	B	409[B]	-	-	0/0/0/0	0/0/0/0
3	FMT	B	410	-	-	0/0/0/0	0/0/0/0
3	FMT	B	411	-	-	0/0/0/0	0/0/0/0
4	CXS	B	412	-	-	0/8/16/16	0/1/1/1
4	CXS	B	413[A]	-	-	0/8/16/16	0/1/1/1
5	GOL	B	414[A]	-	-	0/4/4/4	0/0/0/0
5	GOL	B	414[B]	-	-	0/4/4/4	0/0/0/0

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	413[A]	CXS	C1-S	-7.33	1.67	1.77
4	B	412	CXS	C1-S	-6.21	1.68	1.77
5	A	409	GOL	O2-C2	2.56	1.51	1.43
4	B	413[A]	CXS	O2-S	6.31	1.63	1.45
4	B	412	CXS	O1-S	7.18	1.66	1.45

The worst 5 of 18 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	B	414[A]	GOL	C3-C2-C1	-3.58	97.83	111.63
4	B	412	CXS	O1-S-C1	-2.61	103.77	106.92
4	B	413[A]	CXS	O3-S-O2	-2.45	105.29	111.27
4	B	412	CXS	C8-C7-C6	-2.26	104.04	111.19
4	B	412	CXS	O3-S-C1	2.09	109.14	105.77

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	409	GOL	1	0
3	B	411	FMT	4	0
5	B	414[B]	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	332/332 (100%)	-0.53	4 (1%) 79 79	14, 23, 48, 91	0
1	B	331/332 (99%)	-0.51	10 (3%) 50 47	15, 23, 56, 90	0
All	All	663/664 (99%)	-0.52	14 (2%) 63 62	14, 23, 52, 91	0

The worst 5 of 14 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	330	GLY	4.4
1	B	47	ARG	3.9
1	B	325	GLY	3.7
1	A	332	GLU	3.6
1	B	46	PRO	2.9

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
5	GOL	B	414[A]	6/6	0.78	0.19	27,29,32,36	6
5	GOL	B	414[B]	6/6	0.78	0.19	24,30,35,40	6
4	CXS	B	412	14/14	0.83	0.18	35,48,82,93	0
5	GOL	A	409	6/6	0.87	0.21	43,48,58,59	0
3	FMT	A	407	3/3	0.88	0.18	51,51,56,57	3
4	CXS	B	413[A]	14/14	0.89	0.22	36,46,62,63	14
3	FMT	B	409[B]	3/3	0.91	0.13	25,25,30,42	3
3	FMT	B	409[A]	3/3	0.91	0.13	22,22,25,37	3
3	FMT	A	408	3/3	0.94	0.22	37,37,39,44	0
3	FMT	B	407	3/3	0.95	0.15	36,36,38,46	0
6	UNL	B	415	13/-	0.95	0.08	22,26,34,38	0
3	FMT	B	410	3/3	0.95	0.18	37,37,49,49	0
3	FMT	B	406	3/3	0.96	0.11	35,35,45,47	0
3	FMT	A	406	3/3	0.96	0.23	35,35,49,58	0
3	FMT	B	402	3/3	0.96	0.08	23,23,26,27	0
3	FMT	A	403	3/3	0.97	0.14	31,31,33,38	0
3	FMT	B	408	3/3	0.98	0.05	27,27,28,39	0
3	FMT	B	411	3/3	0.98	0.19	28,28,32,51	0
3	FMT	A	404	3/3	0.98	0.06	29,29,33,40	0
3	FMT	A	405	3/3	0.98	0.09	26,26,35,44	0
3	FMT	B	405	3/3	0.98	0.12	31,31,32,47	0
3	FMT	B	404	3/3	0.98	0.07	29,29,34,40	0
3	FMT	A	402	3/3	0.98	0.08	25,25,25,27	0
2	NA	B	401	1/1	0.99	0.03	15,15,15,15	0
3	FMT	B	403	3/3	0.99	0.04	27,27,31,36	0
2	NA	A	401	1/1	1.00	0.04	14,14,14,14	0

6.5 Other polymers [i](#)

There are no such residues in this entry.