



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 9, 2018 – 01:14 pm GMT

PDB ID : 5A23
Title : SdsA sulfatase triclinic form
Authors : De la Mora, E.; Flores-Hernandez, E.; Jakoncik, J.; Stojanoff, V.; Sanchez-Puig, N.; Moreno, A.
Deposited on : 2015-05-11
Resolution : 2.41 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	trunk30967
Percentile statistics	:	20171227.v01 (using entries in the PDB archive December 27th 2017)
Refmac	:	5.8.0158
CCP4	:	7.0 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	trunk30967


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.41 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	111664	4090 (2.44-2.40)
Clashscore	122126	4587 (2.44-2.40)
Ramachandran outliers	120053	4522 (2.44-2.40)
Sidechain outliers	120020	4523 (2.44-2.40)
RSRZ outliers	108989	3987 (2.44-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	658	<div> <div>2%</div> <div> <div></div> <div>79%</div> <div>15%</div> <div>• 5%</div> </div> </div>
1	B	658	<div> <div>3%</div> <div> <div></div> <div>77%</div> <div>17%</div> <div>• 5%</div> </div> </div>
1	C	658	<div> <div>%</div> <div> <div></div> <div>75%</div> <div>19%</div> <div>• 5%</div> </div> </div>
1	D	658	<div> <div>%</div> <div> <div></div> <div>78%</div> <div>16%</div> <div>• 5%</div> </div> </div>

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 19632 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called SDS HYDROLASE SDSA1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	628	Total	C	N	O	S	0	0	1
			4906	3096	887	913	10			
1	B	628	Total	C	N	O	S	0	0	1
			4906	3096	887	913	10			
1	C	628	Total	C	N	O	S	0	0	1
			4906	3096	887	913	10			
1	D	628	Total	C	N	O	S	0	0	1
			4906	3096	887	913	10			


- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	2	Total	Zn	0	0
			2	2		
2	A	2	Total	Zn	0	0
			2	2		
2	D	2	Total	Zn	0	0
			2	2		
2	C	2	Total	Zn	0	0
			2	2		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: SDS HYDROLASE SDSA1



• Molecule 1: SDS HYDROLASE SDSA1



• Molecule 1: SDS HYDROLASE SDSA1



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	86.90Å 86.80Å 129.03Å 107.59° 102.96° 95.59°	Depositor
Resolution (Å)	118.77 – 2.41 19.73 – 2.41	Depositor EDS
% Data completeness (in resolution range)	71.7 (118.77-2.41) 62.0 (19.73-2.41)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.27 (at 2.41Å)	Xtriage
Refinement program	REFMAC 5.8.0107	Depositor
R, R_{free}	0.253 , 0.307 0.258 , 0.308	Depositor DCC
R_{free} test set	4958 reflections (5.19%)	wwPDB-VP
Wilson B-factor (Å ²)	41.1	Xtriage
Anisotropy	0.154	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 21.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.39$, $\langle L^2 \rangle = 0.22$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	19632	wwPDB-VP
Average B, all atoms (Å ²)	46.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.32% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.97	1/5012 (0.0%)	1.07	21/6805 (0.3%)
1	B	0.82	2/5012 (0.0%)	0.95	4/6805 (0.1%)
1	C	0.85	0/5012	1.00	18/6805 (0.3%)
1	D	0.92	0/5012	1.02	18/6805 (0.3%)
All	All	0.89	3/20048 (0.0%)	1.01	61/27220 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	1
1	C	0	1
All	All	0	3

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	299	GLU	CD-OE2	5.89	1.32	1.25
1	B	299	GLU	CD-OE2	5.68	1.31	1.25
1	B	338	GLU	CD-OE1	5.33	1.31	1.25

The worst 5 of 61 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	59	ARG	NE-CZ-NH1	10.62	125.61	120.30
1	A	370	ASP	CB-CG-OD2	9.75	127.07	118.30
1	A	59	ARG	NE-CZ-NH2	-9.45	115.58	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	370	ASP	CB-CG-OD1	-9.26	109.97	118.30
1	C	217	ARG	NE-CZ-NH1	8.78	124.69	120.30

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	40	GLU	Peptide
1	B	523	ASP	Peptide
1	C	134	TRP	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4906	0	4818	52	0
1	B	4906	0	4818	100	0
1	C	4906	0	4818	87	0
1	D	4906	0	4818	66	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0
2	C	2	0	0	0	0
2	D	2	0	0	1	0
All	All	19632	0	19272	280	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 280 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:299:GLU:OE2	2:D:1001:ZN:ZN	1.17	0.91
1:C:35:ARG:C	1:C:38:GLU:HG2	2.00	0.82
1:C:35:ARG:NH2	1:C:89:ASP:OD1	2.12	0.81
1:C:35:ARG:O	1:C:38:GLU:HG2	1.82	0.78
1:B:146:ALA:HB1	1:B:181:LEU:HD21	1.65	0.77

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	622/658 (94%)	581 (93%)	37 (6%)	4 (1%)	27	38
1	B	622/658 (94%)	586 (94%)	33 (5%)	3 (0%)	31	43
1	C	622/658 (94%)	571 (92%)	49 (8%)	2 (0%)	43	57
1	D	622/658 (94%)	575 (92%)	41 (7%)	6 (1%)	17	24
All	All	2488/2632 (94%)	2313 (93%)	160 (6%)	15 (1%)	27	38

5 of 15 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	140	LEU
1	B	81	PHE
1	B	138	ASP
1	B	173	ASP
1	C	38	GLU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	499/521 (96%)	486 (97%)	13 (3%)	49	68
1	B	499/521 (96%)	487 (98%)	12 (2%)	52	70
1	C	499/521 (96%)	486 (97%)	13 (3%)	49	68

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	D	499/521 (96%)	488 (98%)	11 (2%)	55 73
All	All	1996/2084 (96%)	1947 (98%)	49 (2%)	50 69

5 of 49 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	550	ARG
1	C	62	ARG
1	D	520	VAL
1	B	614	SER
1	C	71	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 12 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	102	ASN
1	C	185	GLN
1	C	345	ASN
1	B	524	GLN
1	C	307	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 8 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	628/658 (95%)	-0.33	13 (2%) 63 60	13, 38, 73, 104	0
1	B	628/658 (95%)	-0.08	19 (3%) 50 47	23, 50, 84, 118	0
1	C	628/658 (95%)	-0.24	9 (1%) 75 73	20, 45, 75, 123	0
1	D	628/658 (95%)	-0.30	6 (0%) 82 80	18, 42, 73, 107	0
All	All	2512/2632 (95%)	-0.24	47 (1%) 66 63	13, 44, 79, 123	0

The worst 5 of 47 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	522	ARG	7.7
1	B	492	ALA	5.3
1	B	530	GLY	4.1
1	C	524	GLN	4.1
1	B	484	ALA	3.9

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	ZN	C	1002	1/1	0.97	0.16	50,50,50,50	0
2	ZN	B	1002	1/1	0.97	0.11	52,52,52,52	0
2	ZN	D	1002	1/1	0.98	0.12	48,48,48,48	1
2	ZN	C	1001	1/1	0.98	0.13	52,52,52,52	0
2	ZN	D	1001	1/1	0.99	0.14	27,27,27,27	1
2	ZN	B	1001	1/1	0.99	0.10	57,57,57,57	0
2	ZN	A	1001	1/1	0.99	0.15	39,39,39,39	0
2	ZN	A	1002	1/1	1.00	0.12	44,44,44,44	0

6.5 Other polymers [i](#)

There are no such residues in this entry.