



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2018 – 06:41 pm GMT

PDB ID : 2A3Y
Title : Pentameric crystal structure of human serum amyloid P-component bound to Bis-1,2-{{(Z)-2carboxy-2-methyl-1,3-dioxane}-5-yloxycarbamoyl}-ethane.
Authors : Ho, J.G.; Kitov, P.I.; Paszkiewicz, E.; Sadowska, J.; Bundle, D.R.; Ng, K.K.
Deposited on : 2005-06-27
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.3 (157068), CSD as539be (2018)
Xtriage (Phenix) : 1.13
EDS : trunk30967
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)
Refmac : 5.8.0158
CCP4 : 7.0 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk30967


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	111664	7193 (2.00-2.00)
Clashscore	122126	8267 (2.00-2.00)
Ramachandran outliers	120053	8166 (2.00-2.00)
Sidechain outliers	120020	8165 (2.00-2.00)
RSRZ outliers	108989	7011 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	204	<div> <div>82%</div> <div>17%</div> <div>•</div> </div>
1	B	204	<div> <div>14%</div> <div>82%</div> <div>18%</div> </div>
1	C	204	<div> <div>%</div> <div>85%</div> <div>13%</div> <div>•</div> </div>
1	D	204	<div> <div>83%</div> <div>13%</div> <div>•</div> </div>
1	E	204	<div> <div>88%</div> <div>10%</div> <div>•</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	CPJ	E	613	-	-	-	X

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 8798 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Serum amyloid P-component.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	204	Total	C	N	O	S	0	0	0
			1649	1071	272	303	3			
1	B	204	Total	C	N	O	S	0	0	0
			1649	1071	272	303	3			
1	C	204	Total	C	N	O	S	0	0	0
			1649	1071	272	303	3			
1	D	204	Total	C	N	O	S	0	0	0
			1649	1071	272	303	3			
1	E	204	Total	C	N	O	S	0	0	0
			1649	1071	272	303	3			

- Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	2	Total	Ca	0	0
			2	2		
2	A	2	Total	Ca	0	0
			2	2		
2	D	2	Total	Ca	0	0
			2	2		
2	C	2	Total	Ca	0	0
			2	2		
2	E	2	Total	Ca	0	0
			2	2		

- Molecule 3 is BIS-1,2-{[(Z)-2-CARBOXY-2-METHYL-1,3-DIOXANE]-5-YLOXYCARBA MOYL}-ETHANE (three-letter code: CPJ) (formula: C₁₆H₂₄N₂O₁₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 13 7 6	0	0
3	B	1	Total C O 13 7 6	0	0
3	C	1	Total C O 13 7 6	0	0
3	D	1	Total C O 13 7 6	0	0
3	E	1	Total C O 13 7 6	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	79	Total O 79 79	0	0
4	B	49	Total O 49 49	0	0
4	C	105	Total O 105 105	0	0
4	D	144	Total O 144 144	0	0
4	E	101	Total O 101 101	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Serum amyloid P-component




• Molecule 1: Serum amyloid P-component




• Molecule 1: Serum amyloid P-component



• Molecule 1: Serum amyloid P-component



• Molecule 1: Serum amyloid P-component



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	94.72Å 69.96Å 101.37Å 90.00° 97.09° 90.00°	Depositor
Resolution (Å)	50.60 – 2.00 50.62 – 2.00	Depositor EDS
% Data completeness (in resolution range)	94.5 (50.60-2.00) 94.5 (50.62-2.00)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	0.05	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.49 (at 2.00Å)	Xtriage
Refinement program	REFMAC 5.1.24	Depositor
R, R_{free}	0.216 , 0.250 0.219 , 0.251	Depositor DCC
R_{free} test set	4216 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	17.6	Xtriage
Anisotropy	0.751	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 50.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	8798	wwPDB-VP
Average B, all atoms (Å ²)	12.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.32% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CPJ, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.33	0/1696	0.66	4/2306 (0.2%)
1	B	0.34	0/1696	0.64	2/2306 (0.1%)
1	C	0.35	0/1696	0.69	3/2306 (0.1%)
1	D	0.37	0/1696	0.69	4/2306 (0.2%)
1	E	0.34	0/1696	0.67	2/2306 (0.1%)
All	All	0.34	0/8480	0.67	15/11530 (0.1%)

There are no bond length outliers.

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	157	LEU	CA-CB-CG	6.53	130.33	115.30
1	C	157	LEU	CA-CB-CG	5.83	128.72	115.30
1	A	157	LEU	CA-CB-CG	5.68	128.36	115.30
1	D	184	ASP	CB-CG-OD2	5.65	123.38	118.30
1	C	58	ASP	CB-CG-OD2	5.57	123.32	118.30
1	A	184	ASP	CB-CG-OD2	5.56	123.30	118.30
1	A	58	ASP	CB-CG-OD2	5.48	123.23	118.30
1	C	184	ASP	CB-CG-OD2	5.47	123.22	118.30
1	D	58	ASP	CB-CG-OD2	5.33	123.09	118.30
1	E	184	ASP	CB-CG-OD2	5.32	123.09	118.30
1	B	58	ASP	CB-CG-OD2	5.30	123.07	118.30
1	B	184	ASP	CB-CG-OD2	5.30	123.07	118.30
1	D	61	LEU	CA-CB-CG	5.26	127.40	115.30
1	E	58	ASP	CB-CG-OD2	5.17	122.95	118.30
1	A	156	ASP	CB-CG-OD2	5.10	122.89	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1649	0	1624	15	0
1	B	1649	0	1624	13	0
1	C	1649	0	1624	9	0
1	D	1649	0	1624	13	0
1	E	1649	0	1624	9	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0
2	C	2	0	0	0	0
2	D	2	0	0	0	0
2	E	2	0	0	0	0
3	A	13	0	8	0	0
3	B	13	0	8	0	0
3	C	13	0	8	0	0
3	D	13	0	8	0	0
3	E	13	0	8	0	0
4	A	79	0	0	0	0
4	B	49	0	0	0	0
4	C	105	0	0	0	0
4	D	144	0	0	1	0
4	E	101	0	0	0	0
All	All	8798	0	8160	59	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (59) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:66:GLU:HG2	4:D:757:HOH:O	1.77	0.84
1:E:30:LEU:HB2	1:E:125:VAL:HG13	1.59	0.83
1:B:95:CYS:HB2	1:B:108:TRP:HB2	1.80	0.63
1:B:52:TYR:HB3	1:B:61:LEU:O	2.00	0.61
1:A:61:LEU:HD23	1:A:75:ILE:HD13	1.85	0.59
1:D:86:GLU:OE2	1:D:109:ILE:HG21	2.02	0.58
1:A:62:LEU:HB3	1:A:74:TYR:HB2	1.87	0.56

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:52:TYR:HB3	1:E:61:LEU:O	2.07	0.54
1:A:34:THR:HG21	1:A:164:LEU:HB2	1.91	0.53
1:D:70:GLU:HG2	1:D:83:LYS:HG2	1.91	0.53
1:C:57:ARG:CZ	1:C:126:GLU:HG3	2.39	0.53
1:C:62:LEU:HB3	1:C:74:TYR:HB2	1.92	0.52
1:C:30:LEU:HB2	1:C:125:VAL:HG13	1.92	0.52
1:A:95:CYS:HB2	1:A:108:TRP:HB2	1.91	0.51
1:B:62:LEU:HB3	1:B:74:TYR:HB2	1.92	0.51
1:C:52:TYR:HB3	1:C:61:LEU:O	2.10	0.51
1:B:21:ASN:HB2	1:B:191:GLU:HB2	1.91	0.51
1:D:30:LEU:HB2	1:D:125:VAL:HG13	1.93	0.51
1:E:62:LEU:HB3	1:E:74:TYR:HB2	1.94	0.50
1:E:138:ASP:HB2	1:E:143:LYS:HB3	1.94	0.50
1:D:52:TYR:HB3	1:D:61:LEU:O	2.12	0.49
1:A:33:PHE:CZ	1:A:61:LEU:HD21	2.48	0.49
1:E:60:GLU:HG3	1:E:125:VAL:HG23	1.94	0.48
1:A:54:THR:HG21	1:A:126:GLU:HB2	1.94	0.48
1:A:21:ASN:HB2	1:A:191:GLU:HB2	1.95	0.48
1:B:33:PHE:CE2	1:B:98:TRP:HB3	2.48	0.48
1:B:30:LEU:HB2	1:B:125:VAL:CG2	2.44	0.47
1:A:52:TYR:HB3	1:A:61:LEU:HB3	1.96	0.46
1:B:35:LEU:HB2	1:B:159:MET:HG3	1.97	0.46
1:D:14:GLU:OE2	1:D:146:ARG:HD3	2.15	0.46
1:D:62:LEU:HB3	1:D:74:TYR:HB2	1.98	0.46
1:E:30:LEU:HB2	1:E:125:VAL:CG1	2.39	0.45
1:B:30:LEU:HB2	1:B:125:VAL:HG23	1.99	0.45
1:E:91:PRO:HB3	1:E:203:TRP:CE3	2.51	0.45
1:A:52:TYR:HB3	1:A:61:LEU:O	2.17	0.45
1:E:57:ARG:HE	1:E:126:GLU:HG3	1.82	0.45
1:D:74:TYR:HA	1:D:78:HIS:O	2.17	0.45
1:A:26:LEU:HD13	1:A:182:ILE:HG12	1.99	0.44
1:D:23:ILE:HD11	1:D:191:GLU:HG3	1.99	0.44
1:D:74:TYR:CZ	1:D:79:LYS:HD2	2.53	0.44
1:C:91:PRO:HB3	1:C:203:TRP:CE3	2.52	0.43
1:C:74:TYR:HA	1:C:78:HIS:O	2.17	0.43
1:D:76:GLY:O	1:D:77:ARG:HB2	2.18	0.43
1:B:106:GLU:HG2	1:B:116:LYS:HG2	2.00	0.43
1:B:8:VAL:HG12	1:B:155:GLY:HA3	2.00	0.43
1:D:91:PRO:HB3	1:D:203:TRP:CE3	2.53	0.43
1:A:88:PHE:HA	1:A:89:PRO:HA	1.79	0.43
1:A:10:VAL:HG22	1:A:153:GLU:HG2	2.00	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:31:GLN:HG2	1:C:100:SER:HB3	2.01	0.42
1:B:37:PHE:HB3	1:B:157:LEU:HD22	2.01	0.42
1:D:13:ARG:HG2	1:D:195:TYR:HD2	1.84	0.42
1:B:88:PHE:HA	1:B:89:PRO:HA	1.87	0.41
1:E:8:VAL:HG13	1:E:201:LEU:HB2	2.03	0.41
1:A:134:GLY:HA2	1:A:150:PHE:HB3	2.03	0.41
1:C:155:GLY:HA2	1:C:185:TRP:CZ2	2.56	0.41
1:C:37:PHE:CE1	1:C:49:LEU:HG	2.56	0.41
1:A:91:PRO:HB3	1:A:203:TRP:CE3	2.56	0.40
1:A:99:GLU:HB2	1:A:163:VAL:HG21	2.04	0.40
1:B:128:GLN:N	1:B:129:PRO:HD3	2.37	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	202/204 (99%)	198 (98%)	4 (2%)	0	100	100
1	B	202/204 (99%)	198 (98%)	4 (2%)	0	100	100
1	C	202/204 (99%)	197 (98%)	5 (2%)	0	100	100
1	D	202/204 (99%)	198 (98%)	4 (2%)	0	100	100
1	E	202/204 (99%)	199 (98%)	3 (2%)	0	100	100
All	All	1010/1020 (99%)	990 (98%)	20 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar

resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	181/181 (100%)	172 (95%)	9 (5%)	27	22
1	B	181/181 (100%)	169 (93%)	12 (7%)	18	13
1	C	181/181 (100%)	167 (92%)	14 (8%)	14	9
1	D	181/181 (100%)	166 (92%)	15 (8%)	12	7
1	E	181/181 (100%)	168 (93%)	13 (7%)	16	10
All	All	905/905 (100%)	842 (93%)	63 (7%)	16	11

All (63) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	4	LEU
1	A	13	ARG
1	A	22	LEU
1	A	52	TYR
1	A	57	ARG
1	A	133	LEU
1	A	146	ARG
1	A	157	LEU
1	A	183	LEU
1	B	4	LEU
1	B	22	LEU
1	B	38	ARG
1	B	49	LEU
1	B	52	TYR
1	B	55	GLN
1	B	77	ARG
1	B	130	LYS
1	B	170	LEU
1	B	171	SER
1	B	181	ASN
1	B	183	LEU
1	C	4	LEU
1	C	22	LEU
1	C	28	LYS
1	C	44	SER
1	C	49	LEU
1	C	52	TYR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	55	GLN
1	C	77	ARG
1	C	125	VAL
1	C	128	GLN
1	C	133	LEU
1	C	157	LEU
1	C	178	LEU
1	C	183	LEU
1	D	4	LEU
1	D	13	ARG
1	D	22	LEU
1	D	52	TYR
1	D	61	LEU
1	D	79	LYS
1	D	125	VAL
1	D	130	LYS
1	D	132	VAL
1	D	133	LEU
1	D	143	LYS
1	D	146	ARG
1	D	157	LEU
1	D	176	THR
1	D	183	LEU
1	E	4	LEU
1	E	22	LEU
1	E	28	LYS
1	E	49	LEU
1	E	52	TYR
1	E	57	ARG
1	E	61	LEU
1	E	82	SER
1	E	125	VAL
1	E	133	LEU
1	E	143	LYS
1	E	183	LEU
1	E	204	VAL

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	93	HIS
1	C	55	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 15 ligands modelled in this entry, 10 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	CPJ	A	610	2	10,13,31	2.27	2 (20%)	9,18,44	2.27	3 (33%)
3	CPJ	B	611	2	10,13,31	2.25	2 (20%)	9,18,44	2.18	3 (33%)
3	CPJ	C	612	2	10,13,31	2.23	2 (20%)	9,18,44	2.16	3 (33%)
3	CPJ	D	614	2	10,13,31	2.23	2 (20%)	9,18,44	2.23	3 (33%)
3	CPJ	E	613	2	10,13,31	2.21	2 (20%)	9,18,44	2.31	3 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	CPJ	A	610	2	-	0/2/21/51	0/1/1/2
3	CPJ	B	611	2	-	0/2/21/51	0/1/1/2
3	CPJ	C	612	2	-	0/2/21/51	0/1/1/2
3	CPJ	D	614	2	-	0/2/21/51	0/1/1/2

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	CPJ	E	613	2	-	0/2/21/51	0/1/1/2

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	610	CPJ	O11-C3	-3.23	1.43	1.46
3	B	611	CPJ	O11-C3	-3.21	1.43	1.46
3	C	612	CPJ	O11-C3	-3.13	1.43	1.46
3	D	614	CPJ	O11-C3	-3.02	1.43	1.46
3	E	613	CPJ	O11-C3	-2.97	1.43	1.46
3	E	613	CPJ	O11-C12	5.66	1.45	1.33
3	B	611	CPJ	O11-C12	5.68	1.45	1.33
3	C	612	CPJ	O11-C12	5.68	1.45	1.33
3	A	610	CPJ	O11-C12	5.72	1.45	1.33
3	D	614	CPJ	O11-C12	5.77	1.46	1.33

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	D	614	CPJ	O11-C12-O13	-4.46	119.84	125.53
3	A	610	CPJ	O11-C12-O13	-4.45	119.84	125.53
3	E	613	CPJ	O11-C12-O13	-4.44	119.86	125.53
3	C	612	CPJ	O11-C12-O13	-4.43	119.87	125.53
3	B	611	CPJ	O11-C12-O13	-4.31	120.03	125.53
3	C	612	CPJ	C4-O5-C6	2.72	116.31	113.94
3	C	612	CPJ	C2-O1-C6	2.74	116.33	113.94
3	B	611	CPJ	C2-O1-C6	2.81	116.39	113.94
3	D	614	CPJ	C4-O5-C6	2.84	116.42	113.94
3	B	611	CPJ	C4-O5-C6	2.93	116.50	113.94
3	A	610	CPJ	C2-O1-C6	2.95	116.52	113.94
3	A	610	CPJ	C4-O5-C6	3.04	116.59	113.94
3	E	613	CPJ	C4-O5-C6	3.13	116.67	113.94
3	D	614	CPJ	C2-O1-C6	3.17	116.71	113.94
3	E	613	CPJ	C2-O1-C6	3.32	116.84	113.94

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	204/204 (100%)	-0.25	1 (0%) 90 90	7, 12, 18, 25	0
1	B	204/204 (100%)	0.76	28 (13%) 3 2	6, 13, 20, 23	0
1	C	204/204 (100%)	-0.34	2 (0%) 82 81	4, 8, 16, 26	0
1	D	204/204 (100%)	-0.35	1 (0%) 90 90	2, 7, 16, 26	0
1	E	204/204 (100%)	-0.32	1 (0%) 90 90	4, 11, 18, 25	0
All	All	1020/1020 (100%)	-0.10	33 (3%) 47 47	2, 11, 18, 26	0

All (33) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	25	PRO	5.6
1	B	26	LEU	4.2
1	B	28	LYS	4.1
1	B	23	ILE	3.8
1	B	27	GLU	3.8
1	B	124	PHE	3.7
1	B	160	TRP	3.7
1	C	26	LEU	3.6
1	C	27	GLU	3.5
1	A	204	VAL	3.5
1	B	125	VAL	3.4
1	B	167	GLU	3.3
1	B	170	LEU	3.2
1	B	1	HIS	3.1
1	D	175	GLY	2.9
1	B	29	PRO	2.9
1	B	179	PRO	2.9
1	B	180	ALA	2.9
1	B	132	VAL	2.8
1	B	178	LEU	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	24	THR	2.6
1	B	204	VAL	2.6
1	B	57	ARG	2.6
1	B	52	TYR	2.4
1	B	98	TRP	2.3
1	B	164	LEU	2.3
1	B	166	PRO	2.3
1	B	188	LEU	2.3
1	E	204	VAL	2.3
1	B	189	ASN	2.2
1	B	126	GLU	2.0
1	B	181	ASN	2.0
1	B	128	GLN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	CPJ	B	611	13/30	0.69	0.33	115,117,118,118	0
3	CPJ	A	610	13/30	0.71	0.30	91,92,94,94	0
3	CPJ	E	613	13/30	0.76	0.41	93,94,95,96	0
3	CPJ	D	614	13/30	0.78	0.35	70,75,81,81	0
3	CPJ	C	612	13/30	0.81	0.25	82,85,88,89	0
2	CA	E	609	1/1	0.95	0.10	22,22,22,22	0
2	CA	E	608	1/1	0.96	0.08	20,20,20,20	0
2	CA	A	601	1/1	0.98	0.06	17,17,17,17	0
2	CA	D	606	1/1	0.99	0.07	9,9,9,9	0
2	CA	B	602	1/1	0.99	0.08	25,25,25,25	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	CA	A	600	1/1	0.99	0.10	16,16,16,16	0
2	CA	C	605	1/1	0.99	0.07	14,14,14,14	0
2	CA	D	607	1/1	0.99	0.07	11,11,11,11	0
2	CA	C	604	1/1	0.99	0.07	14,14,14,14	0
2	CA	B	603	1/1	0.99	0.04	22,22,22,22	0

6.5 Other polymers [i](#)

There are no such residues in this entry.