



wwPDB X-ray Structure Validation Summary Report ⓘ

Dec 17, 2018 – 09:41 PM EST

PDB ID : 6A4B
Title : Structure of TREX2 in complex with a duplex DNA with 2 nucleotide 3'-overhang
Authors : Hsiao, Y.Y.
Deposited on : 2018-06-19
Resolution : 2.70 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	rb-20031633
Percentile statistics	:	20171227.v01 (using entries in the PDB archive December 27th 2017)
Refmac	:	5.8.0158
CCP4	:	7.0 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20031633

1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R _{free}	111664	2449 (2.70-2.70)
Clashscore	122126	2756 (2.70-2.70)
Ramachandran outliers	120053	2716 (2.70-2.70)
Sidechain outliers	120020	2716 (2.70-2.70)
RSRZ outliers	108989	2376 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	256	<div><div>6%</div><div><div></div><div>70%</div><div>12%</div><div>•</div><div>17%</div></div></div>
1	B	256	<div><div>4%</div><div><div></div><div>67%</div><div>16%</div><div>•</div><div>16%</div></div></div>
1	C	256	<div><div>8%</div><div><div></div><div>72%</div><div>12%</div><div></div><div>16%</div></div></div>
1	D	256	<div><div>12%</div><div><div></div><div>71%</div><div>13%</div><div></div><div>16%</div></div></div>
1	E	256	<div><div>7%</div><div><div></div><div>71%</div><div>13%</div><div></div><div>15%</div></div></div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	256	<div> <div>2%</div> <div>71%</div> <div>13%</div> <div>15%</div> </div>
2	G	18	<div> <div>17%</div> <div>67%</div> <div>17%</div> <div>17%</div> </div>
2	H	18	<div> <div>22%</div> <div>67%</div> <div>17%</div> <div>17%</div> </div>
2	I	18	<div> <div>6%</div> <div>61%</div> <div>22%</div> <div>17%</div> </div>
2	J	18	<div> <div>11%</div> <div>89%</div> <div>11%</div> </div>
2	K	18	<div> <div>6%</div> <div>72%</div> <div>6%</div> <div>22%</div> </div>
2	L	18	<div> <div>11%</div> <div>56%</div> <div>28%</div> <div>17%</div> </div>

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 11982 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Three prime repair exonuclease 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	213	Total	C	N	O	S	0	0	0
			1661	1055	287	309	10			
1	B	216	Total	C	N	O	S	0	0	0
			1680	1067	291	312	10			
1	C	215	Total	C	N	O	S	0	0	0
			1677	1063	293	311	10			
1	D	215	Total	C	N	O	S	0	0	0
			1684	1069	295	310	10			
1	E	217	Total	C	N	O	S	0	0	0
			1690	1071	295	314	10			
1	F	217	Total	C	N	O	S	0	0	0
			1690	1071	295	314	10			

There are 120 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-19	MET	-	initiating methionine	UNP Q9R1A9
A	-18	GLY	-	expression tag	UNP Q9R1A9
A	-17	SER	-	expression tag	UNP Q9R1A9
A	-16	SER	-	expression tag	UNP Q9R1A9
A	-15	HIS	-	expression tag	UNP Q9R1A9
A	-14	HIS	-	expression tag	UNP Q9R1A9
A	-13	HIS	-	expression tag	UNP Q9R1A9
A	-12	HIS	-	expression tag	UNP Q9R1A9
A	-11	HIS	-	expression tag	UNP Q9R1A9
A	-10	HIS	-	expression tag	UNP Q9R1A9
A	-9	SER	-	expression tag	UNP Q9R1A9
A	-8	SER	-	expression tag	UNP Q9R1A9
A	-7	GLY	-	expression tag	UNP Q9R1A9
A	-6	LEU	-	expression tag	UNP Q9R1A9
A	-5	VAL	-	expression tag	UNP Q9R1A9
A	-4	PRO	-	expression tag	UNP Q9R1A9
A	-3	ARG	-	expression tag	UNP Q9R1A9

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	GLY	-	expression tag	UNP Q9R1A9
A	-1	SER	-	expression tag	UNP Q9R1A9
A	0	HIS	-	expression tag	UNP Q9R1A9
B	-19	MET	-	initiating methionine	UNP Q9R1A9
B	-18	GLY	-	expression tag	UNP Q9R1A9
B	-17	SER	-	expression tag	UNP Q9R1A9
B	-16	SER	-	expression tag	UNP Q9R1A9
B	-15	HIS	-	expression tag	UNP Q9R1A9
B	-14	HIS	-	expression tag	UNP Q9R1A9
B	-13	HIS	-	expression tag	UNP Q9R1A9
B	-12	HIS	-	expression tag	UNP Q9R1A9
B	-11	HIS	-	expression tag	UNP Q9R1A9
B	-10	HIS	-	expression tag	UNP Q9R1A9
B	-9	SER	-	expression tag	UNP Q9R1A9
B	-8	SER	-	expression tag	UNP Q9R1A9
B	-7	GLY	-	expression tag	UNP Q9R1A9
B	-6	LEU	-	expression tag	UNP Q9R1A9
B	-5	VAL	-	expression tag	UNP Q9R1A9
B	-4	PRO	-	expression tag	UNP Q9R1A9
B	-3	ARG	-	expression tag	UNP Q9R1A9
B	-2	GLY	-	expression tag	UNP Q9R1A9
B	-1	SER	-	expression tag	UNP Q9R1A9
B	0	HIS	-	expression tag	UNP Q9R1A9
C	-19	MET	-	initiating methionine	UNP Q9R1A9
C	-18	GLY	-	expression tag	UNP Q9R1A9
C	-17	SER	-	expression tag	UNP Q9R1A9
C	-16	SER	-	expression tag	UNP Q9R1A9
C	-15	HIS	-	expression tag	UNP Q9R1A9
C	-14	HIS	-	expression tag	UNP Q9R1A9
C	-13	HIS	-	expression tag	UNP Q9R1A9
C	-12	HIS	-	expression tag	UNP Q9R1A9
C	-11	HIS	-	expression tag	UNP Q9R1A9
C	-10	HIS	-	expression tag	UNP Q9R1A9
C	-9	SER	-	expression tag	UNP Q9R1A9
C	-8	SER	-	expression tag	UNP Q9R1A9
C	-7	GLY	-	expression tag	UNP Q9R1A9
C	-6	LEU	-	expression tag	UNP Q9R1A9
C	-5	VAL	-	expression tag	UNP Q9R1A9
C	-4	PRO	-	expression tag	UNP Q9R1A9
C	-3	ARG	-	expression tag	UNP Q9R1A9
C	-2	GLY	-	expression tag	UNP Q9R1A9
C	-1	SER	-	expression tag	UNP Q9R1A9

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	0	HIS	-	expression tag	UNP Q9R1A9
D	-19	MET	-	initiating methionine	UNP Q9R1A9
D	-18	GLY	-	expression tag	UNP Q9R1A9
D	-17	SER	-	expression tag	UNP Q9R1A9
D	-16	SER	-	expression tag	UNP Q9R1A9
D	-15	HIS	-	expression tag	UNP Q9R1A9
D	-14	HIS	-	expression tag	UNP Q9R1A9
D	-13	HIS	-	expression tag	UNP Q9R1A9
D	-12	HIS	-	expression tag	UNP Q9R1A9
D	-11	HIS	-	expression tag	UNP Q9R1A9
D	-10	HIS	-	expression tag	UNP Q9R1A9
D	-9	SER	-	expression tag	UNP Q9R1A9
D	-8	SER	-	expression tag	UNP Q9R1A9
D	-7	GLY	-	expression tag	UNP Q9R1A9
D	-6	LEU	-	expression tag	UNP Q9R1A9
D	-5	VAL	-	expression tag	UNP Q9R1A9
D	-4	PRO	-	expression tag	UNP Q9R1A9
D	-3	ARG	-	expression tag	UNP Q9R1A9
D	-2	GLY	-	expression tag	UNP Q9R1A9
D	-1	SER	-	expression tag	UNP Q9R1A9
D	0	HIS	-	expression tag	UNP Q9R1A9
E	-19	MET	-	initiating methionine	UNP Q9R1A9
E	-18	GLY	-	expression tag	UNP Q9R1A9
E	-17	SER	-	expression tag	UNP Q9R1A9
E	-16	SER	-	expression tag	UNP Q9R1A9
E	-15	HIS	-	expression tag	UNP Q9R1A9
E	-14	HIS	-	expression tag	UNP Q9R1A9
E	-13	HIS	-	expression tag	UNP Q9R1A9
E	-12	HIS	-	expression tag	UNP Q9R1A9
E	-11	HIS	-	expression tag	UNP Q9R1A9
E	-10	HIS	-	expression tag	UNP Q9R1A9
E	-9	SER	-	expression tag	UNP Q9R1A9
E	-8	SER	-	expression tag	UNP Q9R1A9
E	-7	GLY	-	expression tag	UNP Q9R1A9
E	-6	LEU	-	expression tag	UNP Q9R1A9
E	-5	VAL	-	expression tag	UNP Q9R1A9
E	-4	PRO	-	expression tag	UNP Q9R1A9
E	-3	ARG	-	expression tag	UNP Q9R1A9
E	-2	GLY	-	expression tag	UNP Q9R1A9
E	-1	SER	-	expression tag	UNP Q9R1A9
E	0	HIS	-	expression tag	UNP Q9R1A9
F	-19	MET	-	initiating methionine	UNP Q9R1A9

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
F	-18	GLY	-	expression tag	UNP Q9R1A9
F	-17	SER	-	expression tag	UNP Q9R1A9
F	-16	SER	-	expression tag	UNP Q9R1A9
F	-15	HIS	-	expression tag	UNP Q9R1A9
F	-14	HIS	-	expression tag	UNP Q9R1A9
F	-13	HIS	-	expression tag	UNP Q9R1A9
F	-12	HIS	-	expression tag	UNP Q9R1A9
F	-11	HIS	-	expression tag	UNP Q9R1A9
F	-10	HIS	-	expression tag	UNP Q9R1A9
F	-9	SER	-	expression tag	UNP Q9R1A9
F	-8	SER	-	expression tag	UNP Q9R1A9
F	-7	GLY	-	expression tag	UNP Q9R1A9
F	-6	LEU	-	expression tag	UNP Q9R1A9
F	-5	VAL	-	expression tag	UNP Q9R1A9
F	-4	PRO	-	expression tag	UNP Q9R1A9
F	-3	ARG	-	expression tag	UNP Q9R1A9
F	-2	GLY	-	expression tag	UNP Q9R1A9
F	-1	SER	-	expression tag	UNP Q9R1A9
F	0	HIS	-	expression tag	UNP Q9R1A9

- Molecule 2 is a DNA chain called DNA (5'-D(*GP*GP*CP*CP*CP*TP*TP*AP*GP*GP*GP*CP*CP*TP*T)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	G	15	Total	C	N	O	P	0	0	0
			303	145	53	91	14			
2	H	15	Total	C	N	O	P	0	0	0
			303	145	53	91	14			
2	I	15	Total	C	N	O	P	0	0	0
			303	145	53	91	14			
2	J	18	Total	C	N	O	P	0	0	0
			362	174	60	111	17			
2	K	14	Total	C	N	O	P	0	0	0
			283	135	51	84	13			
2	L	15	Total	C	N	O	P	0	0	0
			303	145	53	91	14			

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	G	1	Total	Mg	0	0
			1	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	D	1	Total 1	Mg 1	0	0
3	E	2	Total 2	Mg 2	0	0
3	B	1	Total 1	Mg 1	0	0
3	I	1	Total 1	Mg 1	0	0
3	C	1	Total 1	Mg 1	0	0
3	A	2	Total 2	Mg 2	0	0
3	L	1	Total 1	Mg 1	0	0
3	F	1	Total 1	Mg 1	0	0

- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	5	Total 5	O 5	0	0
4	B	5	Total 5	O 5	0	0
4	C	5	Total 5	O 5	0	0
4	D	5	Total 5	O 5	0	0
4	E	5	Total 5	O 5	0	0
4	F	2	Total 2	O 2	0	0
4	G	3	Total 3	O 3	0	0
4	L	2	Total 2	O 2	0	0

- Molecule 1: Three prime repair exonuclease 2




ALA


- Molecule 1: Three prime repair exonuclease 2




- Molecule 1: Three prime repair exonuclease 2




- Molecule 1: Three prime repair exonuclease 2




- Molecule 2: DNA (5'-D(*GP*GP*CP*CP*CP*TP*TP*AP*GP*GP*GP*CP*CP*TP*T)-3')





- Molecule 2: DNA (5'-D(*GP*GP*CP*CP*CP*TP*TP*AP*GP*GP*GP*CP*CP*TP*T)-3')





• Molecule 2: DNA (5'-D(*GP*GP*CP*CP*CP*TP*TP*AP*GP*GP*GP*CP*CP*TP*T)-3')




• Molecule 2: DNA (5'-D(*GP*GP*CP*CP*CP*TP*TP*AP*GP*GP*GP*CP*CP*TP*T)-3')



• Molecule 2: DNA (5'-D(*GP*GP*CP*CP*CP*TP*TP*AP*GP*GP*GP*CP*CP*TP*T)-3')



• Molecule 2: DNA (5'-D(*GP*GP*CP*CP*CP*TP*TP*AP*GP*GP*GP*CP*CP*TP*T)-3')



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	48.72Å 87.32Å 432.94Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.23 – 2.70 29.23 – 2.70	Depositor EDS
% Data completeness (in resolution range)	97.2 (29.23-2.70) 96.7 (29.23-2.70)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	0.08	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.08 (at 2.72Å)	Xtriage
Refinement program	PHENIX 1.9_1692	Depositor
R, R_{free}	0.224 , 0.263 0.230 , 0.266	Depositor DCC
R_{free} test set	3961 reflections (7.80%)	wwPDB-VP
Wilson B-factor (Å ²)	74.1	Xtriage
Anisotropy	0.289	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 39.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	11982	wwPDB-VP
Average B, all atoms (Å ²)	82.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.35% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.22	0/1701	0.41	0/2310
1	B	0.22	0/1723	0.39	0/2343
1	C	0.22	0/1718	0.38	0/2334
1	D	0.22	0/1726	0.39	0/2344
1	E	0.21	0/1732	0.37	0/2353
1	F	0.22	0/1731	0.38	0/2350
2	G	0.56	0/337	0.92	0/516
2	H	0.50	0/337	0.91	0/516
2	I	0.55	0/337	0.90	0/516
2	J	0.51	0/403	0.94	0/620
2	K	0.56	0/315	0.88	0/482
2	L	0.55	0/337	0.89	0/516
All	All	0.30	0/12397	0.52	0/17200

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1661	0	1631	21	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	1680	0	1643	25	0
1	C	1677	0	1641	16	0
1	D	1684	0	1647	17	0
1	E	1690	0	1654	18	0
1	F	1690	0	1653	19	0
2	G	303	0	172	3	0
2	H	303	0	172	2	0
2	I	303	0	172	3	0
2	J	362	0	206	2	0
2	K	283	0	160	1	0
2	L	303	0	172	4	0
3	A	2	0	0	0	0
3	B	1	0	0	0	0
3	C	1	0	0	0	0
3	D	1	0	0	0	0
3	E	2	0	0	0	0
3	F	1	0	0	0	0
3	G	1	0	0	0	0
3	I	1	0	0	0	0
3	L	1	0	0	0	0
4	A	5	0	0	0	0
4	B	5	0	0	0	0
4	C	5	0	0	0	0
4	D	5	0	0	0	0
4	E	5	0	0	0	0
4	F	2	0	0	0	0
4	G	3	0	0	0	0
4	L	2	0	0	0	0
All	All	11982	0	10923	118	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 118 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:125:PRO:HB2	1:E:225:MET:HG3	1.66	0.76
1:D:45:ARG:HA	1:D:51:LEU:HA	1.72	0.71
1:A:92:GLY:O	1:B:107:ARG:NH2	2.23	0.71
1:F:16:GLU:HG2	1:F:190:ALA:HB2	1.72	0.71
1:B:34:ALA:HB3	1:B:58:ASP:HB2	1.72	0.70

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	207/256 (81%)	201 (97%)	5 (2%)	1 (0%)	31	58
1	B	212/256 (83%)	203 (96%)	8 (4%)	1 (0%)	31	58
1	C	211/256 (82%)	202 (96%)	9 (4%)	0	100	100
1	D	209/256 (82%)	201 (96%)	8 (4%)	0	100	100
1	E	213/256 (83%)	204 (96%)	9 (4%)	0	100	100
1	F	211/256 (82%)	201 (95%)	10 (5%)	0	100	100
All	All	1263/1536 (82%)	1212 (96%)	49 (4%)	2 (0%)	49	77

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	93	PHE
1	A	21	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	181/214 (85%)	177 (98%)	4 (2%)	55	83
1	B	182/214 (85%)	175 (96%)	7 (4%)	36	66
1	C	181/214 (85%)	179 (99%)	2 (1%)	76	91

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	D	182/214 (85%)	180 (99%)	2 (1%)	76	91
1	E	183/214 (86%)	180 (98%)	3 (2%)	65	87
1	F	183/214 (86%)	180 (98%)	3 (2%)	65	87
All	All	1092/1284 (85%)	1071 (98%)	21 (2%)	60	85

5 of 21 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	138	LEU
1	C	138	LEU
1	E	183	GLU
1	B	137	HIS
1	F	23	MET

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 8 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	188	HIS
1	F	22	ASN
1	C	132	GLN
1	B	140	GLN
1	C	108	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry

Of 11 ligands modelled in this entry, 11 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2	OWAB(Å ²)	Q < 0.9
1	A	213/256 (83%)	0.31	15 (7%) 16 14	45, 68, 101, 122	0
1	B	216/256 (84%)	0.23	10 (4%) 32 30	44, 69, 103, 117	0
1	C	215/256 (83%)	0.46	20 (9%) 8 6	55, 79, 111, 149	0
1	D	215/256 (83%)	0.67	31 (14%) 2 1	49, 72, 116, 132	0
1	E	217/256 (84%)	0.47	17 (7%) 13 11	57, 83, 109, 131	0
1	F	217/256 (84%)	0.27	6 (2%) 53 53	61, 80, 110, 121	0
2	G	15/18 (83%)	0.52	3 (20%) 1 0	65, 98, 124, 133	0
2	H	15/18 (83%)	1.04	4 (26%) 0 0	76, 117, 142, 144	0
2	I	15/18 (83%)	0.18	1 (6%) 18 16	74, 98, 126, 145	0
2	J	18/18 (100%)	0.52	2 (11%) 5 4	84, 98, 124, 125	0
2	K	14/18 (77%)	0.32	1 (7%) 16 14	74, 97, 119, 119	0
2	L	15/18 (83%)	0.54	2 (13%) 3 2	69, 112, 137, 153	0
All	All	1385/1644 (84%)	0.41	112 (8%) 12 9	44, 78, 114, 153	0

The worst 5 of 112 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	186	ALA	8.1
1	D	227	VAL	6.0
1	D	226	TYR	5.4
1	D	14	ASP	5.0
1	C	45	ARG	4.5

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	MG	A	301	1/1	0.87	0.84	75,75,75,75	0
3	MG	A	302	1/1	0.88	0.12	75,75,75,75	0
3	MG	D	301	1/1	0.89	0.85	75,75,75,75	0
3	MG	B	301	1/1	0.90	0.16	69,69,69,69	0
3	MG	E	301	1/1	0.93	0.47	75,75,75,75	0
3	MG	L	101	1/1	0.93	0.45	69,69,69,69	0
3	MG	C	301	1/1	0.94	0.32	87,87,87,87	0
3	MG	E	302	1/1	0.94	0.06	83,83,83,83	0
3	MG	I	101	1/1	0.95	0.34	66,66,66,66	0
3	MG	F	301	1/1	0.95	0.13	76,76,76,76	0
3	MG	G	101	1/1	0.97	0.29	62,62,62,62	0

6.5 Other polymers [i](#)

There are no such residues in this entry.