



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2018 – 11:05 am GMT

PDB ID : 2A5D
Title : Structural basis for the activation of cholera toxin by human ARF6-GTP
Authors : O'Neal, C.J.; Jobling, M.G.; Holmes, R.K.; Hol, W.G.J.
Deposited on : 2005-06-30
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.3 (157068), CSD as539be (2018)
Xtriage (Phenix) : 1.13
EDS : trunk30967
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)
Refmac : 5.8.0158
CCP4 : 7.0 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk30967


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	111664	5253 (1.80-1.80)
Clashscore	122126	6077 (1.80-1.80)
Ramachandran outliers	120053	6011 (1.80-1.80)
Sidechain outliers	120020	6010 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Mol	Chain	Length	Quality of chain
1	A	175	
2	B	193	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
6	GOL	B	1496	-	X	X	-

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 3102 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ADP-ribosylation factor 6.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	163	Total	C	N	O	S	0	1	0
			1299	824	229	240	6			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MET	-	INITIATING METHIONINE	UNP P62330

- Molecule 2 is a protein called Cholera enterotoxin, A chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	187	Total	C	N	O	S	0	1	0
			1472	930	261	278	3			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	0	SER	-	CLONING ARTIFACT	UNP P01555
B	110	ASP	GLU	ENGINEERED	UNP P01555
B	112	ASP	GLU	ENGINEERED	UNP P01555
B	187	SER	CYS	ENGINEERED	UNP P01555

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Mg	0	0
			1	1		

- Molecule 4 is GUANOSINE-5'-TRIPHOSPHATE (three-letter code: GTP) (formula: C₁₀H₁₆N₅O₁₄P₃).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	P	0	0
			32	10	5	14	3		

- Molecule 5 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	B	1	Total Na 1 1	0	0

- Molecule 6 is GLYCEROL (three-letter code: GOL) (formula: $\text{C}_3\text{H}_8\text{O}_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	B	1	Total	C	O	0	0
			6	3	3		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	126	Total	O	0	0
			126	126		
7	B	165	Total	O	0	0
			165	165		

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: ADP-ribosylation factor 6

Chain A:  82% 10% • 7%



- Molecule 2: Cholera enterotoxin, A chain

Chain B:  84% 12% • •



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	44.91Å 91.45Å 98.55Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	45.74 – 1.80 45.72 – 1.80	Depositor EDS
% Data completeness (in resolution range)	93.3 (45.74-1.80) 93.3 (45.72-1.80)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.56 (at 1.79Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.167 , 0.199 0.178 , 0.206	Depositor DCC
R_{free} test set	1798 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å ²)	24.9	Xtriage
Anisotropy	0.823	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 36.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	3102	wwPDB-VP
Average B, all atoms (Å ²)	23.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.41% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GTP, GOL, MG, NA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.75	0/1331	0.85	2/1804 (0.1%)
2	B	0.89	1/1525 (0.1%)	0.88	3/2079 (0.1%)
All	All	0.83	1/2856 (0.0%)	0.87	5/3883 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	154	ASP	CB-CG	5.43	1.63	1.51

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	172	ARG	NE-CZ-NH1	9.25	124.92	120.30
2	B	172	ARG	NE-CZ-NH2	-7.89	116.36	120.30
2	B	154	ASP	CB-CG-OD1	5.97	123.67	118.30
1	A	69	LYS	CD-CE-NZ	5.52	124.39	111.70
1	A	71	ARG	NE-CZ-NH2	-5.45	117.58	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1299	0	1274	12	0
2	B	1472	0	1345	22	1
3	A	1	0	0	0	0
4	A	32	0	12	0	0
5	B	1	0	0	0	0
6	B	6	0	5	5	0
7	A	126	0	0	1	0
7	B	165	0	0	7	0
All	All	3102	0	2636	34	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (34) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:74[B]:GLN:CD	7:B:283:HOH:O	1.99	1.01
2:B:74[B]:GLN:NE2	7:B:283:HOH:O	1.95	1.00
2:B:152:ASN:H	2:B:152:ASN:HD22	1.28	0.81
2:B:171:HIS:HD2	2:B:173:ALA:H	1.25	0.80
2:B:74[B]:GLN:OE1	7:B:283:HOH:O	2.04	0.75
2:B:163:GLY:H	6:B:1496:GOL:C3	2.03	0.72
2:B:76:ILE:HD11	7:B:1389:HOH:O	1.91	0.70
2:B:76:ILE:CD1	7:B:1389:HOH:O	2.42	0.67
2:B:160:ASP:C	6:B:1496:GOL:H31	2.18	0.64
2:B:48:THR:HG22	2:B:52:PHE:H	1.62	0.63
1:A:72:PRO:HG2	6:B:1496:GOL:H11	1.83	0.60
2:B:1:ASN:HB3	2:B:151:SER:O	2.02	0.58
1:A:79:THR:HG23	7:A:1356:HOH:O	2.02	0.58
2:B:152:ASN:H	2:B:152:ASN:ND2	2.01	0.56
2:B:75:THR:HB	2:B:76:ILE:HD12	1.87	0.56
2:B:131:HIS:CE1	2:B:132:PHE:CE2	2.95	0.55
2:B:98:ASN:HD21	2:B:108:PRO:HA	1.74	0.53
2:B:171:HIS:CD2	2:B:173:ALA:H	2.16	0.52
1:A:113:ARG:HG3	1:A:114:ASP:N	2.26	0.50
2:B:3:ASP:CG	7:B:1450:HOH:O	2.50	0.50
1:A:60:ASN:ND2	1:A:62:TRP:HE1	2.09	0.49
2:B:160:ASP:O	6:B:1496:GOL:H31	2.13	0.48
1:A:60:ASN:HD21	1:A:62:TRP:HE1	1.61	0.47
2:B:48:THR:HG22	2:B:51:GLY:N	2.31	0.46
1:A:71:ARG:N	1:A:72:PRO:CD	2.80	0.45
1:A:79:THR:HG22	1:A:111:GLU:OE1	2.18	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:14:MET:HE1	1:A:173:TYR:HB3	2.01	0.42
2:B:160:ASP:O	6:B:1496:GOL:O2	2.38	0.41
1:A:53:THR:HA	1:A:57:VAL:O	2.20	0.41
2:B:54:ARG:HD3	7:B:314:HOH:O	2.19	0.41
1:A:110:ARG:O	1:A:113:ARG:HG2	2.20	0.41
2:B:135:LEU:HD12	2:B:135:LEU:N	2.35	0.41
1:A:150:TYR:HB2	1:A:168:TRP:CE2	2.56	0.41
1:A:14:MET:CE	1:A:173:TYR:HB3	2.51	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:74[B]:GLN:NE2	2:B:106:PRO:O[4_566]	2.03	0.17

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	162/175 (93%)	159 (98%)	2 (1%)	1 (1%)	27	13
2	B	186/193 (96%)	182 (98%)	4 (2%)	0	100	100
All	All	348/368 (95%)	341 (98%)	6 (2%)	1 (0%)	43	28

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	127	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	137/153 (90%)	134 (98%)	3 (2%)	55	42
2	B	149/159 (94%)	145 (97%)	4 (3%)	48	34
All	All	286/312 (92%)	279 (98%)	7 (2%)	50	38

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	19	LEU
1	A	113	ARG
1	A	169	LEU
2	B	2	ASP
2	B	48	THR
2	B	139	LEU
2	B	152	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	48	ASN
1	A	60	ASN
2	B	27	GLN
2	B	98	ASN
2	B	111	GLN
2	B	131	HIS
2	B	152	ASN
2	B	171	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
4	GTP	A	1230	3	27,34,34	1.52	2 (7%)	29,54,54	1.92	8 (27%)
6	GOL	B	1496	-	5,5,5	1.07	1 (20%)	5,5,5	6.70	5 (100%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GTP	A	1230	3	-	0/18/38/38	0/3/3/3
6	GOL	B	1496	-	-	0/4/4/4	0/0/0/0

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	B	1496	GOL	O2-C2	-2.13	1.37	1.43
4	A	1230	GTP	C6-N1	2.79	1.38	1.33
4	A	1230	GTP	PG-O3B	6.35	1.69	1.60

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	B	1496	GOL	O2-C2-C3	-10.51	61.00	109.00
6	B	1496	GOL	O2-C2-C1	-9.58	65.23	109.00
4	A	1230	GTP	N3-C2-N1	-4.73	120.47	127.41
4	A	1230	GTP	C5-C6-N1	-3.44	118.57	123.47
4	A	1230	GTP	C6-C5-C4	-3.09	117.82	120.85
4	A	1230	GTP	C1'-N9-C4	-2.84	121.73	126.64
4	A	1230	GTP	PB-O3B-PG	-2.78	123.30	132.63
6	B	1496	GOL	O1-C1-C2	2.10	120.29	110.11
4	A	1230	GTP	N2-C2-N1	2.21	120.73	117.25
4	A	1230	GTP	C2-N3-C4	2.33	117.88	115.16
6	B	1496	GOL	O3-C3-C2	2.44	121.93	110.11
6	B	1496	GOL	C3-C2-C1	3.50	125.10	111.63
4	A	1230	GTP	C6-N1-C2	4.13	122.00	116.06

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	B	1496	GOL	5	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

6.3 Carbohydrates

Unable to reproduce the depositors R factor - this section is therefore empty.

6.4 Ligands

Unable to reproduce the depositors R factor - this section is therefore empty.

6.5 Other polymers

Unable to reproduce the depositors R factor - this section is therefore empty.