



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 8, 2018 – 09:44 pm GMT

PDB ID : 3A8K
Title : Crystal Structure of ETD97N-EHred complex
Authors : Okamura-Ikeda, K.; Hosaka, H.
Deposited on : 2009-10-06
Resolution : 1.95 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.3 (157068), CSD as539be (2018)
Xtriage (Phenix) : 1.13
EDS : trunk30967
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)
Refmac : 5.8.0158
CCP4 : 7.0 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk30967


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	111664	2220 (1.96-1.96)
Clashscore	122126	2333 (1.96-1.96)
Ramachandran outliers	120053	2314 (1.96-1.96)
Sidechain outliers	120020	2314 (1.96-1.96)
RSRZ outliers	108989	2174 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	364	<div> <div>%</div> <div> <div></div> <div>87%</div> <div>13%</div> <div>.</div> </div> </div>
1	B	364	<div> <div>2%</div> <div> <div></div> <div>87%</div> <div>11%</div> <div>.</div> </div> </div>
1	C	364	<div> <div>2%</div> <div> <div></div> <div>88%</div> <div>11%</div> <div>..</div> </div> </div>
1	D	364	<div> <div>%</div> <div> <div></div> <div>85%</div> <div>13%</div> <div>..</div> </div> </div>
2	E	129	<div> <div>5%</div> <div> <div></div> <div>87%</div> <div>8%</div> <div>..</div> </div> </div>
2	F	129	<div> <div>5%</div> <div> <div></div> <div>86%</div> <div>10%</div> <div>..</div> </div> </div>

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 14698 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Aminomethyltransferase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	363	Total	C	N	O	S	0	2	0
			2829	1786	496	532	15			
1	B	363	Total	C	N	O	S	0	2	0
			2827	1786	494	532	15			
1	C	362	Total	C	N	O	S	0	0	0
			2809	1774	492	528	15			
1	D	362	Total	C	N	O	S	0	2	0
			2822	1782	495	530	15			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	97	ASN	ASP	ENGINEERED	UNP P27248
B	97	ASN	ASP	ENGINEERED	UNP P27248
C	97	ASN	ASP	ENGINEERED	UNP P27248
D	97	ASN	ASP	ENGINEERED	UNP P27248

- Molecule 2 is a protein called Glycine cleavage system H protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	E	126	Total	C	N	O	S	0	0	0
			958	603	145	206	4			
2	F	126	Total	C	N	O	S	0	0	0
			958	603	145	206	4			

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	370	Total	O	0	0
			370	370		
3	B	352	Total	O	0	0
			352	352		



Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	C	313	Total 313	O 313	0	0
3	D	298	Total 298	O 298	0	0
3	E	113	Total 113	O 113	0	0
3	F	49	Total 49	O 49	0	0



- Molecule 2: Glycine cleavage system H protein



- Molecule 2: Glycine cleavage system H protein



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	54.12Å 88.41Å 97.86Å 91.56° 102.42° 89.59°	Depositor
Resolution (Å)	43.97 – 1.95 43.97 – 1.95	Depositor EDS
% Data completeness (in resolution range)	98.1 (43.97-1.95) 98.1 (43.97-1.95)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.21 (at 1.95Å)	Xtriage
Refinement program	REFMAC 5.5.0102	Depositor
R, R_{free}	0.171 , 0.229 0.171 , 0.228	Depositor DCC
R_{free} test set	6363 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å ²)	21.7	Xtriage
Anisotropy	0.174	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 52.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.059 for -h,k,-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	14698	wwPDB-VP
Average B, all atoms (Å ²)	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.27% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: LA2

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.12	7/2893 (0.2%)	0.98	5/3917 (0.1%)
1	B	1.11	0/2891	0.99	8/3914 (0.2%)
1	C	1.09	3/2867 (0.1%)	0.97	3/3884 (0.1%)
1	D	1.06	3/2886 (0.1%)	0.94	5/3909 (0.1%)
2	E	0.92	0/954	0.87	2/1301 (0.2%)
2	F	0.84	0/954	0.88	2/1301 (0.2%)
All	All	1.07	13/13445 (0.1%)	0.96	25/18226 (0.1%)

The worst 5 of 13 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	118	GLU	CG-CD	7.03	1.62	1.51
1	C	100	VAL	CB-CG1	-6.35	1.39	1.52
1	C	238	GLU	CB-CG	6.16	1.63	1.52
1	D	153	ALA	CA-CB	5.93	1.65	1.52
1	A	322	TYR	CD2-CE2	5.80	1.48	1.39

The worst 5 of 25 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	266	ARG	NE-CZ-NH2	-13.68	113.46	120.30
1	D	266	ARG	NE-CZ-NH2	-10.34	115.13	120.30
1	C	266	ARG	NE-CZ-NH1	10.22	125.41	120.30
1	B	62	ARG	NE-CZ-NH1	8.86	124.73	120.30
1	B	208	ARG	NE-CZ-NH1	8.50	124.55	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2829	0	2792	28	0
1	B	2827	0	2792	30	0
1	C	2809	0	2768	25	0
1	D	2822	0	2785	35	0
2	E	958	0	909	9	0
2	F	958	0	909	6	0
3	A	370	0	0	5	0
3	B	352	0	0	6	0
3	C	313	0	0	5	0
3	D	298	0	0	7	0
3	E	113	0	0	2	0
3	F	49	0	0	0	0
All	All	14698	0	12955	130	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 130 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:298:ARG:HH12	1:D:340:GLN:HE21	1.12	0.97
1:C:97:ASN:HB2	3:C:698:HOH:O	1.71	0.89
1:D:180[A]:ASP:OD2	1:D:202:LYS:HD2	1.74	0.87
1:D:26:PRO:HG2	3:D:845:HOH:O	1.81	0.81
1:C:146:VAL:HG12	1:C:151:ALA:HB1	1.63	0.80

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	363/364 (100%)	353 (97%)	10 (3%)	0	100	100
1	B	363/364 (100%)	354 (98%)	7 (2%)	2 (1%)	27	15
1	C	360/364 (99%)	349 (97%)	10 (3%)	1 (0%)	43	32
1	D	362/364 (100%)	352 (97%)	8 (2%)	2 (1%)	27	15
2	E	123/129 (95%)	119 (97%)	3 (2%)	1 (1%)	21	10
2	F	123/129 (95%)	119 (97%)	4 (3%)	0	100	100
All	All	1694/1714 (99%)	1646 (97%)	42 (2%)	6 (0%)	36	24

5 of 6 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	E	126	GLU
1	D	359	GLY
1	C	42	ASP
1	B	42	ASP
1	B	342	ARG

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	293/292 (100%)	287 (98%)	6 (2%)	58	51
1	B	293/292 (100%)	287 (98%)	6 (2%)	58	51
1	C	291/292 (100%)	284 (98%)	7 (2%)	52	43
1	D	293/292 (100%)	283 (97%)	10 (3%)	40	27
2	E	100/103 (97%)	96 (96%)	4 (4%)	34	20
2	F	100/103 (97%)	95 (95%)	5 (5%)	27	13
All	All	1370/1374 (100%)	1332 (97%)	38 (3%)	47	35

5 of 38 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	295	LEU
1	D	266	ARG
2	F	16	LEU
1	D	100	VAL
1	D	269	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 21 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	37	HIS
1	C	3	GLN
1	D	2	GLN
1	B	21	HIS
1	D	32	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	LA2	E	64	2	18,19,20	0.65	0	14,21,23	1.61	1 (7%)
2	LA2	F	64	2	18,19,20	0.65	0	14,21,23	1.52	2 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	LA2	E	64	2	-	0/16/20/22	0/0/0/0
2	LA2	F	64	2	-	0/16/20/22	0/0/0/0

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	E	64	LA2	CB-CA-C	-4.55	103.18	111.85
2	F	64	LA2	C7-C8-S8	-3.53	110.06	113.74
2	F	64	LA2	CB-CA-C	-2.89	106.34	111.85

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	363/364 (99%)	-0.25	2 (0%) 89 93	10, 17, 38, 55	0
1	B	363/364 (99%)	-0.16	7 (1%) 66 75	11, 18, 39, 57	1 (0%)
1	C	362/364 (99%)	-0.20	9 (2%) 57 66	13, 22, 43, 56	1 (0%)
1	D	362/364 (99%)	-0.08	5 (1%) 75 82	13, 22, 42, 63	0
2	E	125/129 (96%)	0.01	7 (5%) 24 33	17, 26, 48, 68	0
2	F	125/129 (96%)	0.39	6 (4%) 30 41	22, 36, 49, 57	0
All	All	1700/1714 (99%)	-0.12	36 (2%) 63 72	10, 22, 44, 68	2 (0%)

The worst 5 of 36 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	363	ALA	5.6
1	C	302	ALA	4.3
2	F	20	ALA	4.2
1	C	275	HIS	3.9
1	A	275	HIS	3.5

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	LA2	F	64	20/21	0.96	0.09	23,31,39,44	0
2	LA2	E	64	20/21	0.97	0.09	19,24,32,43	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.