



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2018 – 05:44 am GMT

PDB ID : 4A8O  
Title : Non-Catalytic Ions Direct the RNA-Dependent RNA Polymerase of Bacterial dsRNA virus phi6 from De Novo Initiation to Elongation  
Authors : Wright, S.; Poranen, M.M.; Bamford, D.H.; Stuart, D.I.; Grimes, J.M.  
Deposited on : 2011-11-21  
Resolution : 2.67 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	trunk30967
Percentile statistics	:	20171227.v01 (using entries in the PDB archive December 27th 2017)
Refmac	:	5.8.0158
CCP4	:	7.0 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	trunk30967


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.67 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	111664	3333 (2.70-2.66)
Clashscore	122126	3672 (2.70-2.66)
Ramachandran outliers	120053	3620 (2.70-2.66)
Sidechain outliers	120020	3620 (2.70-2.66)
RSRZ outliers	108989	3248 (2.70-2.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	665	<div> <div>%</div> <div> <div></div> <div>91%</div> <div>8%</div> </div> </div>
1	B	665	<div> <div>%</div> <div> <div></div> <div>92%</div> <div>8%</div> </div> </div>
1	C	665	<div> <div>2%</div> <div> <div></div> <div>92%</div> <div>7%</div> </div> </div>

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 15798 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called RNA-DIRECTED RNA POLYMERASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	664	Total	C	N	O	S	0	0	0
			5265	3342	915	976	32			
1	B	664	Total	C	N	O	S	0	0	0
			5265	3342	915	976	32			
1	C	664	Total	C	N	O	S	0	0	0
			5265	3342	915	976	32			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	456	MET	ILE	conflict	UNP P11124
A	634	GLN	GLU	conflict	UNP P11124
B	456	MET	ILE	conflict	UNP P11124
B	634	GLN	GLU	conflict	UNP P11124
C	456	MET	ILE	conflict	UNP P11124
C	634	GLN	GLU	conflict	UNP P11124


- Molecule 2 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Mn	0	0
			1	1		
2	A	1	Total	Mn	0	0
			1	1		
2	C	1	Total	Mn	0	0
			1	1		


### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### ● Molecule 1: RNA-DIRECTED RNA POLYMERASE



#### ● Molecule 1: RNA-DIRECTED RNA POLYMERASE



#### ● Molecule 1: RNA-DIRECTED RNA POLYMERASE



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	106.19Å 92.45Å 141.04Å 90.00° 101.47° 90.00°	Depositor
Resolution (Å)	39.09 – 2.67 69.12 – 2.67	Depositor EDS
% Data completeness (in resolution range)	(Not available) (39.09-2.67) 97.4 (69.12-2.67)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.51 (at 2.65Å)	Xtriage
Refinement program	BUSTER 2.9.2	Depositor
R, $R_{free}$	0.215 , 0.225 0.215 , 0.223	Depositor DCC
$R_{free}$ test set	3738 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	52.2	Xtriage
Anisotropy	0.871	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 26.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	15798	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	60.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.25% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.41	0/5396	0.62	1/7297 (0.0%)
1	B	0.40	0/5396	0.62	0/7297
1	C	0.39	0/5396	0.63	0/7297
All	All	0.40	0/16188	0.62	1/21891 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	206	GLN	N-CA-C	-5.30	96.70	111.00

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5265	0	5167	30	0
1	B	5265	0	5167	27	0
1	C	5265	0	5167	18	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
All	All	15798	0	15501	66	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (66) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:251:ARG:HH11	1:A:255:GLN:HE22	1.41	0.69
1:B:251:ARG:HH11	1:B:255:GLN:HE22	1.40	0.67
1:C:251:ARG:HH11	1:C:255:GLN:HE22	1.42	0.67
1:B:606:ALA:HB3	1:B:609:ALA:HB2	1.80	0.63
1:A:608:GLN:HE22	1:B:593:ARG:CZ	2.12	0.63
1:A:600:ARG:HH12	1:B:581:GLU:HG3	1.66	0.61
1:A:600:ARG:NH1	1:B:581:GLU:HG3	2.15	0.60
1:A:593:ARG:HG2	1:B:42:GLY:HA2	1.83	0.60
1:C:391:LEU:HD22	1:C:398:THR:HG22	1.86	0.58
1:A:392:SER:O	1:A:398:THR:HG21	2.05	0.57
1:A:151:THR:HG22	1:A:163:LYS:HG3	1.89	0.53
1:C:651:VAL:O	1:C:655:GLU:HB2	2.10	0.52
1:A:651:VAL:O	1:A:655:GLU:HB2	2.10	0.52
1:A:203:TYR:HE1	1:A:271:THR:HG22	1.75	0.51
1:B:537:ARG:HG3	1:B:537:ARG:HH11	1.75	0.51
1:C:203:TYR:HE1	1:C:271:THR:HG22	1.75	0.51
1:B:203:TYR:HE1	1:B:271:THR:HG22	1.76	0.50
1:A:392:SER:O	1:A:398:THR:CG2	2.60	0.50
1:B:145:ILE:HD12	1:B:164:ILE:HD13	1.94	0.50
1:A:407:SER:HA	1:A:448:GLN:HE22	1.77	0.50
1:B:407:SER:HA	1:B:448:GLN:HE22	1.77	0.50
1:B:651:VAL:O	1:B:655:GLU:HB2	2.11	0.50
1:A:47:ASP:OD1	1:A:49:ARG:HD3	2.12	0.49
1:A:534:SER:O	1:B:48:PRO:HG2	2.13	0.49
1:A:94:PRO:HB3	1:A:269:ARG:HG3	1.95	0.48
1:C:407:SER:HA	1:C:448:GLN:HE22	1.77	0.48
1:A:145:ILE:HD12	1:A:164:ILE:HD13	1.95	0.48
1:A:203:TYR:CE1	1:A:271:THR:HG22	2.48	0.48
1:A:70:ASP:OD2	1:A:74:ARG:HD2	2.13	0.48
1:C:70:ASP:OD2	1:C:74:ARG:HD2	2.14	0.48
1:B:70:ASP:OD2	1:B:74:ARG:HD2	2.14	0.47
1:C:203:TYR:CE1	1:C:271:THR:HG22	2.48	0.47
1:B:391:LEU:HD22	1:B:398:THR:HG22	1.96	0.47
1:B:301:THR:HG23	1:B:448:GLN:HG3	1.96	0.47
1:C:145:ILE:HD12	1:C:164:ILE:HD13	1.95	0.47
1:A:72:TYR:CE2	1:A:476:LYS:HD3	2.50	0.47
1:A:537:ARG:HD2	1:B:49:ARG:HG3	1.95	0.47

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:203:TYR:CE1	1:B:271:THR:HG22	2.49	0.47
1:C:301:THR:HG23	1:C:448:GLN:HG3	1.96	0.47
1:C:86:GLY:O	1:C:89:HIS:HD2	1.98	0.47
1:C:94:PRO:HB3	1:C:269:ARG:HG3	1.97	0.46
1:B:47:ASP:OD1	1:B:49:ARG:HD3	2.15	0.46
1:B:86:GLY:O	1:B:89:HIS:HD2	1.99	0.46
1:A:301:THR:HG23	1:A:448:GLN:HG3	1.98	0.45
1:A:86:GLY:O	1:A:89:HIS:HD2	1.99	0.45
1:A:44:LEU:HD21	1:B:257:GLY:HA3	1.98	0.45
1:B:72:TYR:CE2	1:B:476:LYS:HD3	2.52	0.45
1:A:217:THR:HG23	1:A:219:LYS:H	1.81	0.45
1:B:94:PRO:HB3	1:B:269:ARG:HG3	1.99	0.45
1:C:47:ASP:OD1	1:C:49:ARG:HD3	2.18	0.44
1:A:427:LYS:HE3	1:C:12:ILE:HG21	2.00	0.44
1:C:72:TYR:CE2	1:C:476:LYS:HD3	2.51	0.44
1:A:328:HIS:HD2	1:A:329:ASP:OD1	2.00	0.44
1:A:163:LYS:HE3	1:A:163:LYS:HB2	1.68	0.43
1:A:608:GLN:NE2	1:B:593:ARG:CZ	2.80	0.43
1:B:217:THR:HG23	1:B:219:LYS:H	1.84	0.43
1:C:217:THR:HG23	1:C:219:LYS:H	1.83	0.43
1:A:153:ILE:HA	1:A:154:PRO:HA	1.87	0.43
1:B:406:MET:HB3	1:B:410:TYR:CE2	2.54	0.43
1:C:658:LEU:HG	1:C:662:MET:HG3	2.01	0.42
1:B:475:LEU:HD21	1:B:482:PRO:HG3	1.99	0.42
1:A:94:PRO:CB	1:A:269:ARG:HG3	2.50	0.41
1:A:475:LEU:HD21	1:A:482:PRO:HG3	2.02	0.41
1:C:395:GLN:HB3	1:C:398:THR:HG23	2.03	0.41
1:B:497:LEU:HA	1:B:497:LEU:HD12	1.99	0.40
1:C:475:LEU:HD21	1:C:482:PRO:HG3	2.03	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	662/665 (100%)	641 (97%)	19 (3%)	2 (0%)	43	69
1	B	662/665 (100%)	640 (97%)	19 (3%)	3 (0%)	31	56
1	C	662/665 (100%)	640 (97%)	18 (3%)	4 (1%)	27	51
All	All	1986/1995 (100%)	1921 (97%)	56 (3%)	9 (0%)	31	56

All (9) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	607	ARG
1	B	607	ARG
1	C	393	SER
1	A	136	SER
1	B	136	SER
1	B	393	SER
1	C	136	SER
1	C	607	ARG
1	C	2	ARG

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	557/558 (100%)	538 (97%)	19 (3%)	40	68
1	B	557/558 (100%)	540 (97%)	17 (3%)	43	71
1	C	557/558 (100%)	541 (97%)	16 (3%)	45	73
All	All	1671/1674 (100%)	1619 (97%)	52 (3%)	43	71

All (52) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	3	ARG
1	A	30	ARG
1	A	43	LEU
1	A	63	ASP

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	78	ASN
1	A	81	ARG
1	A	82	THR
1	A	102	LEU
1	A	113	ASP
1	A	163	LYS
1	A	269	ARG
1	A	384	ASN
1	A	391	LEU
1	A	497	LEU
1	A	506	ARG
1	A	608	GLN
1	A	629	GLN
1	A	662	MET
1	A	664	ARG
1	B	3	ARG
1	B	30	ARG
1	B	43	LEU
1	B	63	ASP
1	B	78	ASN
1	B	82	THR
1	B	102	LEU
1	B	113	ASP
1	B	269	ARG
1	B	324	ASP
1	B	384	ASN
1	B	404	LEU
1	B	497	LEU
1	B	506	ARG
1	B	607	ARG
1	B	629	GLN
1	B	664	ARG
1	C	3	ARG
1	C	43	LEU
1	C	63	ASP
1	C	78	ASN
1	C	82	THR
1	C	113	ASP
1	C	269	ARG
1	C	384	ASN
1	C	391	LEU
1	C	392	SER

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	C	404	LEU
1	C	497	LEU
1	C	576	TRP
1	C	582	SER
1	C	629	GLN
1	C	664	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (35) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	15	GLN
1	A	26	GLN
1	A	78	ASN
1	A	89	HIS
1	A	91	ASN
1	A	255	GLN
1	A	309	ASN
1	A	328	HIS
1	A	448	GLN
1	A	525	GLN
1	A	608	GLN
1	A	626	ASN
1	A	634	GLN
1	B	15	GLN
1	B	26	GLN
1	B	78	ASN
1	B	89	HIS
1	B	255	GLN
1	B	309	ASN
1	B	328	HIS
1	B	448	GLN
1	B	525	GLN
1	B	626	ASN
1	B	634	GLN
1	C	15	GLN
1	C	26	GLN
1	C	78	ASN
1	C	89	HIS
1	C	255	GLN
1	C	309	ASN
1	C	328	HIS
1	C	448	GLN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	C	525	GLN
1	C	626	ASN
1	C	634	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

Of 3 ligands modelled in this entry, 3 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	664/665 (99%)	0.03	5 (0%) 86 86	42, 55, 80, 108	0
1	B	664/665 (99%)	0.17	8 (1%) 79 79	43, 58, 85, 136	0
1	C	664/665 (99%)	0.18	14 (2%) 63 63	38, 59, 89, 157	0
All	All	1992/1995 (99%)	0.13	27 (1%) 75 75	38, 57, 84, 157	0

All (27) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	607	ARG	11.4
1	C	608	GLN	9.2
1	C	606	ALA	8.9
1	B	608	GLN	5.4
1	C	610	GLY	4.4
1	B	604	SER	4.3
1	B	664	ARG	4.1
1	B	1	PRO	3.7
1	B	607	ARG	3.7
1	C	576	TRP	3.4
1	C	609	ALA	3.1
1	C	642	HIS	3.1
1	C	603	ALA	3.0
1	A	148	GLY	2.9
1	C	604	SER	2.8
1	A	607	ARG	2.8
1	C	315	LYS	2.7
1	C	537	ARG	2.6
1	A	664	ARG	2.5
1	B	630	TYR	2.5
1	C	600	ARG	2.5
1	B	537	ARG	2.4
1	C	611	LEU	2.2

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	561	CYS	2.1
1	A	2	ARG	2.1
1	C	655	GLU	2.1
1	B	581	GLU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	MN	A	1665	1/1	0.49	0.16	191,191,191,191	0
2	MN	C	1665	1/1	0.72	0.32	165,165,165,165	0
2	MN	B	1665	1/1	0.93	0.19	114,114,114,114	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.