



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2018 – 04:50 pm GMT

PDB ID : 1AXE  
Title : CRYSTAL STRUCTURE OF THE ACTIVE-SITE MUTANT PHE93->TRP  
OF HORSE LIVER ALCOHOL DEHYDROGENASE IN COMPLEX WITH  
NAD AND INHIBITOR TRIFLUOROETHANOL  
Authors : Colby, T.D.; Chin, J.K.; Goldstein, B.M.  
Deposited on : 1997-10-15  
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.3 (157068), CSD as539be (2018)  
Xtriage (Phenix) : 1.13  
EDS : trunk30967  
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)  
Refmac : 5.8.0158  
CCP4 : 7.0 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk30967


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	111664	7193 (2.00-2.00)
Clashscore	122126	8267 (2.00-2.00)
Ramachandran outliers	120053	8166 (2.00-2.00)
Sidechain outliers	120020	8165 (2.00-2.00)
RSRZ outliers	108989	7011 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	374	<div> <div style="width: 100%; height: 10px; position: relative;"> <div style="position: absolute; left: 0; top: -10px; width: 5px; height: 5px; background-color: red;"></div> <div style="position: absolute; left: 87%; top: -10px; width: 5px; height: 5px; background-color: yellow;"></div> <div style="position: absolute; left: 95%; top: -10px; width: 5px; height: 5px; background-color: orange;"></div> <div style="position: absolute; left: 98%; top: -10px; width: 5px; height: 5px; background-color: green;"></div> </div> <div style="display: flex; justify-content: space-between; width: 100%;"> <span>87%</span> <span>12%</span> </div> </div>
1	B	374	<div> <div style="width: 100%; height: 10px; position: relative;"> <div style="position: absolute; left: 84%; top: -10px; width: 5px; height: 5px; background-color: yellow;"></div> <div style="position: absolute; left: 95%; top: -10px; width: 5px; height: 5px; background-color: orange;"></div> <div style="position: absolute; left: 98%; top: -10px; width: 5px; height: 5px; background-color: green;"></div> </div> <div style="display: flex; justify-content: space-between; width: 100%;"> <span>84%</span> <span>14%</span> </div> </div>

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 5796 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ALCOHOL DEHYDROGENASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	374	Total	C	N	O	S	0	0	0
			2788	1771	473	521	23			
1	B	374	Total	C	N	O	S	0	0	0
			2788	1771	473	521	23			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	93	TRP	PHE	ENGINEERED	UNP P00327
B	93	TRP	PHE	ENGINEERED	UNP P00327

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	2	Total	Zn	0	0
			2	2		
2	A	2	Total	Zn	0	0
			2	2		

- Molecule 3 is NICOTINAMIDE-ADENINE-DINUCLEOTIDE (three-letter code: NAD) (formula: C<sub>21</sub>H<sub>27</sub>N<sub>7</sub>O<sub>14</sub>P<sub>2</sub>).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total 44	C 21	N 7	O 14	P 2	0	0
3	B	1	Total 44	C 21	N 7	O 14	P 2	0	0

- Molecule 4 is TRIFLUOROETHANOL (three-letter code: ETF) (formula:  $\text{C}_2\text{H}_3\text{F}_3\text{O}$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total 6	C 2	F 3	O 1	0	0
4	B	1	Total 6	C 2	F 3	O 1	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	61	Total 61	O 61	0	0
5	B	55	Total 55	O 55	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: ALCOHOL DEHYDROGENASE



#### • Molecule 1: ALCOHOL DEHYDROGENASE



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	51.82Å 44.58Å 93.28Å 103.10° 87.59° 70.55°	Depositor
Resolution (Å)	10.00 – 2.00 9.99 – 2.00	Depositor EDS
% Data completeness (in resolution range)	67.6 (10.00-2.00) 80.4 (9.99-2.00)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	0.06	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.40 (at 2.01Å)	Xtriage
Refinement program	X-PLOR 3.1	Depositor
R, $R_{free}$	0.203 , 0.269 0.185 , 0.230	Depositor DCC
$R_{free}$ test set	4128 reflections (10.06%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	22.1	Xtriage
Anisotropy	0.650	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 55.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.51$ , $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	5796	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	23.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.64% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, ETF, NAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.50	0/2841	0.73	0/3841
1	B	0.47	0/2841	0.71	0/3841
All	All	0.49	0/5682	0.72	0/7682

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2788	0	2849	32	0
1	B	2788	0	2849	39	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0
3	A	44	0	26	2	0
3	B	44	0	26	4	0
4	A	6	0	2	1	0
4	B	6	0	2	1	0
5	A	61	0	0	1	0
5	B	55	0	0	2	0
All	All	5796	0	5754	69	0



The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (69) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:101:ARG:HE	1:B:283:GLN:HG3	1.47	0.78
1:A:336:MET:HA	1:A:336:MET:HE2	1.67	0.75
1:A:185:LYS:HE3	5:A:460:HOH:O	1.91	0.70
1:A:101:ARG:NE	1:B:283:GLN:HG3	2.07	0.69
1:B:283:GLN:HE22	1:B:285:ALA:HB3	1.61	0.64
1:A:283:GLN:NE2	1:A:285:ALA:H	1.96	0.63
1:A:187:ALA:HB2	1:A:290:VAL:HG21	1.82	0.61
1:B:166:LEU:HA	1:B:169:VAL:HG22	1.84	0.59
1:A:89:VAL:HG12	1:A:159:LYS:HA	1.85	0.59
1:A:271:ARG:O	1:A:275:MET:HG3	2.03	0.58
1:B:31:PRO:HD3	1:B:37:ARG:HB2	1.85	0.57
1:B:345:LEU:O	1:B:369:ARG:HB2	2.05	0.56
1:B:175:GLY:HA2	1:B:203:VAL:HG22	1.87	0.56
1:B:61:LEU:HD12	1:B:63:VAL:HG12	1.87	0.55
3:A:403:NAD:C4N	4:A:404:ETF:H21	2.36	0.55
1:B:187:ALA:HB2	1:B:290:VAL:HG21	1.88	0.54
1:A:23:ILE:HD11	1:A:355:ILE:CG2	2.36	0.54
1:B:100:CYS:HB2	1:B:112:LEU:HD12	1.90	0.54
1:B:283:GLN:NE2	1:B:285:ALA:H	2.04	0.54
1:B:10:LYS:HA	1:B:24:GLU:O	2.09	0.53
1:B:304:ASN:ND2	1:B:306:MET:H	2.08	0.52
1:B:250:ILE:HD12	1:B:253:VAL:HB	1.91	0.51
1:A:355:ILE:HG13	1:A:372:LEU:HD11	1.92	0.51
1:B:336:MET:HA	1:B:336:MET:CE	2.41	0.51
1:B:181:GLY:O	1:B:185:LYS:HB3	2.11	0.50
1:B:304:ASN:HD22	1:B:304:ASN:C	2.14	0.50
3:B:403:NAD:C4N	4:B:404:ETF:H21	2.42	0.49
1:A:283:GLN:HE22	1:A:285:ALA:HB3	1.78	0.49
1:B:26:VAL:HG12	1:B:132:CYS:HB2	1.95	0.49
1:A:26:VAL:HG12	1:A:132:CYS:HB2	1.95	0.48
1:A:283:GLN:HE22	1:A:285:ALA:H	1.60	0.48
1:A:23:ILE:HD11	1:A:355:ILE:HG22	1.94	0.48
1:B:294:VAL:HG23	3:B:403:NAD:H1D	1.95	0.47
1:A:63:VAL:HG22	1:A:64:ILE:N	2.30	0.47
1:A:70:ALA:HB1	1:A:166:LEU:HD22	1.96	0.47
1:A:359:PHE:O	1:A:363:ARG:HG3	2.15	0.47
1:B:56:THR:HG23	1:B:296:PRO:HA	1.98	0.46

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:242:ASN:OD1	1:B:244:GLN:HG2	2.15	0.46
1:B:61:LEU:HD12	1:B:63:VAL:CG1	2.46	0.46
1:A:83:VAL:HG12	1:A:159:LYS:HB2	1.98	0.46
1:B:271:ARG:HB2	1:B:274:THR:OG1	2.15	0.46
1:A:285:ALA:HB1	5:B:412:HOH:O	2.16	0.46
1:B:195:CYS:HA	1:B:264:PHE:O	2.17	0.45
1:A:348:HIS:O	1:A:370:THR:HA	2.17	0.45
1:B:174:CYS:SG	3:B:403:NAD:H5N	2.57	0.45
1:B:226:LYS:HG2	5:B:416:HOH:O	2.17	0.45
1:B:272:LEU:HD22	1:B:301:LEU:HB3	1.99	0.45
1:B:343:ASP:HB2	1:B:344:PRO:HD3	1.98	0.44
1:A:101:ARG:CZ	1:B:283:GLN:HG3	2.47	0.44
1:A:351:PRO:HG2	1:A:354:LYS:HG2	1.98	0.44
1:B:304:ASN:HD22	1:B:305:PRO:N	2.16	0.44
1:A:231:LYS:HE3	1:A:344:PRO:O	2.17	0.44
1:A:16:GLU:HB2	1:A:19:LYS:HG3	2.00	0.43
1:A:100:CYS:HB2	1:A:112:LEU:HD12	2.00	0.43
1:A:23:ILE:HD11	1:A:355:ILE:HG21	2.00	0.43
1:B:2:THR:HB	1:B:7:ILE:HD11	2.00	0.43
1:A:135:LYS:HA	1:A:136:PRO:HD3	1.92	0.42
1:A:96:GLN:HB3	1:A:325:LYS:HB2	2.00	0.42
1:B:244:GLN:H	1:B:244:GLN:HG2	1.53	0.42
1:A:101:ARG:NH2	1:B:283:GLN:HG3	2.35	0.41
1:B:9:CYS:HB2	1:B:148:GLN:OE1	2.20	0.41
1:A:90:ILE:HA	1:A:91:PRO:HD3	1.87	0.41
1:A:174:CYS:SG	3:A:403:NAD:H5N	2.60	0.41
1:A:250:ILE:HD12	1:A:253:VAL:HB	2.02	0.41
1:B:70:ALA:HB1	1:B:166:LEU:HD22	2.02	0.41
1:B:135:LYS:HA	1:B:136:PRO:HD3	1.86	0.41
1:B:292:VAL:O	3:B:403:NAD:H2N	2.21	0.41
1:B:283:GLN:HE22	1:B:285:ALA:H	1.69	0.41
1:B:63:VAL:HG22	1:B:64:ILE:N	2.35	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	372/374 (100%)	353 (95%)	18 (5%)	1 (0%)	43	39
1	B	372/374 (100%)	347 (93%)	24 (6%)	1 (0%)	43	39
All	All	744/748 (100%)	700 (94%)	42 (6%)	2 (0%)	43	39

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	174	CYS
1	A	174	CYS

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	308/308 (100%)	301 (98%)	7 (2%)	53	56
1	B	308/308 (100%)	297 (96%)	11 (4%)	38	36
All	All	616/616 (100%)	598 (97%)	18 (3%)	45	45

All (18) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	117	SER
1	A	239	GLU
1	A	268	VAL
1	A	283	GLN
1	A	304	ASN
1	A	309	LEU
1	A	338	LYS
1	B	61	LEU
1	B	191	GLN
1	B	231	LYS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	239	GLU
1	B	244	GLN
1	B	283	GLN
1	B	304	ASN
1	B	309	LEU
1	B	330	LYS
1	B	336	MET
1	B	353	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	283	GLN
1	A	300	ASN
1	A	304	ASN
1	B	283	GLN
1	B	300	ASN
1	B	304	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 4 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	NAD	A	403	-	40,48,48	0.85	0	44,73,73	1.84	8 (18%)
4	ETF	A	404	2	5,5,5	0.41	0	7,7,7	0.64	0
3	NAD	B	403	-	40,48,48	0.94	2 (5%)	44,73,73	2.07	7 (15%)
4	ETF	B	404	2	5,5,5	0.17	0	7,7,7	0.57	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAD	A	403	-	-	0/22/62/62	0/5/5/5
4	ETF	A	404	2	-	0/3/3/3	0/0/0/0
3	NAD	B	403	-	-	0/22/62/62	0/5/5/5
4	ETF	B	404	2	-	0/3/3/3	0/0/0/0

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	403	NAD	O4D-C1D	2.34	1.44	1.41
3	B	403	NAD	C6N-N1N	2.36	1.41	1.35

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	403	NAD	N3A-C2A-N1A	-9.42	120.80	128.86
3	A	403	NAD	N3A-C2A-N1A	-8.53	121.57	128.86
3	A	403	NAD	O2B-C2B-C3B	-2.74	103.05	111.83
3	A	403	NAD	O5D-PN-O1N	-2.24	100.30	109.07
3	B	403	NAD	O7N-C7N-C3N	-2.05	117.06	119.62
3	B	403	NAD	C2N-C3N-C4N	2.02	120.59	118.26
3	A	403	NAD	O3D-C3D-C2D	2.09	118.53	111.83
3	B	403	NAD	C2A-N1A-C6A	2.22	122.52	118.75
3	A	403	NAD	C2A-N1A-C6A	2.30	122.65	118.75
3	B	403	NAD	O2A-PA-O1A	2.50	124.86	112.14
3	A	403	NAD	O3B-C3B-C4B	2.51	118.34	111.06
3	B	403	NAD	N6A-C6A-N1A	2.84	124.47	118.57
3	A	403	NAD	N6A-C6A-N1A	2.89	124.56	118.57

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	403	NAD	C4A-C5A-N7A	3.68	112.96	109.41
3	B	403	NAD	C4A-C5A-N7A	5.20	114.43	109.41

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	403	NAD	2	0
4	A	404	ETF	1	0
3	B	403	NAD	4	0
4	B	404	ETF	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	374/374 (100%)	-0.44	3 (0%) 86 85	8, 20, 36, 60	0
1	B	374/374 (100%)	-0.38	1 (0%) 93 93	11, 23, 42, 69	0
All	All	748/748 (100%)	-0.41	4 (0%) 90 90	8, 21, 40, 69	0

All (4) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	1	SER	3.6
1	A	5	LYS	3.2
1	A	1	SER	2.9
1	A	118	MET	2.5

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	ETF	A	404	6/6	0.91	0.11	22,27,32,32	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
4	ETF	B	404	6/6	0.91	0.12	23,34,37,39	0
2	ZN	B	402	1/1	0.96	0.04	22,22,22,22	0
3	NAD	A	403	44/44	0.96	0.08	2,14,24,26	0
2	ZN	A	401	1/1	0.97	0.05	23,23,23,23	0
2	ZN	A	402	1/1	0.97	0.06	22,22,22,22	0
3	NAD	B	403	44/44	0.97	0.07	9,15,18,20	0
2	ZN	B	401	1/1	0.99	0.02	25,25,25,25	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.