



# Full wwPDB X-ray Structure Validation Report ⓘ

Jul 18, 2019 – 02:02 AM EDT

PDB ID : 2B0S  
Title : Crystal structure analysis of anti-HIV-1 V3 Fab 2219 in complex with MN peptide  
Authors : Stanfield, R.L.; Gorny, M.K.; Zolla-Pazner, S.; Wilson, I.A.  
Deposited on : 2005-09-14  
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.0 (224370), CSD as540be (2019)  
Xtriage (Phenix) : 1.13  
EDS : 2.3.2  
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)  
Refmac : 5.8.0158  
CCP4 : 7.0 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.3.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	111664	4477 (2.30-2.30)
Clashscore	122126	5072 (2.30-2.30)
Ramachandran outliers	120053	5022 (2.30-2.30)
Sidechain outliers	120020	5021 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ .

Mol	Chain	Length	Quality of chain
1	L	215	
2	H	226	
3	P	19	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	ACY	H	502	-	-	X	-

## 2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 3660 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Fab 2219, light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	L	215	Total	C	N	O	S	0	0	0
			1616	1011	274	327	4			

- Molecule 2 is a protein called Fab 2219, heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	H	226	Total	C	N	O	S	1	0	0
			1699	1070	277	345	7			

- Molecule 3 is a protein called MN peptide of Exterior membrane glycoprotein GP120.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	P	16	Total	C	N	O	0	0	0
			134	86	29	19			

There is a discrepancy between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
P	323	ALA	-	INSERTION	UNP P05877

- Molecule 4 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C<sub>2</sub>H<sub>6</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	L	1	Total	C	O	0	0
			4	2	2		
4	H	1	Total	C	O	0	0
			4	2	2		

- Molecule 5 is ACETIC ACID (three-letter code: ACY) (formula:  $C_2H_4O_2$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	L	1	Total	C	O	0	0
			4	2	2		
5	H	1	Total	C	O	0	0
			4	2	2		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	H	1	Total	C	O	0	0
			4	2	2		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	L	71	Total	O	0	0
			71	71		
6	H	110	Total	O	0	0
			110	110		
6	P	10	Total	O	0	0
			10	10		

### 3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Fab 2219, light chain

Chain L:  90% 9%



- Molecule 2: Fab 2219, heavy chain

Chain H:  88% 12%



- Molecule 3: MN peptide of Exterior membrane glycoprotein GP120

Chain P:  58% 21% 5% 16%



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	60.47Å 60.47Å 275.19Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	69.01 – 2.30 45.42 – 2.00	Depositor EDS
% Data completeness (in resolution range)	99.8 (69.01-2.30) 99.4 (45.42-2.00)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.09	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.10 (at 2.00Å)	Xtriage
Refinement program	REFMAC 5.2, CNS	Depositor
R, $R_{free}$	0.217 , 0.267 0.225 , 0.271	Depositor DCC
$R_{free}$ test set	1759 reflections (4.95%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	20.7	Xtriage
Anisotropy	0.197	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 41.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.51$ , $\langle L^2 \rangle = 0.35$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	3660	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	34.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.24% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ACY, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	L	0.60	0/1660	0.65	0/2269
2	H	1.41	2/1744 (0.1%)	0.99	3/2373 (0.1%)
3	P	1.59	1/137 (0.7%)	1.13	2/181 (1.1%)
All	All	1.12	3/3541 (0.1%)	0.85	5/4823 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	H	1	1
3	P	0	1
All	All	1	2

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	H	229	SER	C-O	51.31	2.20	1.23
3	P	309	ILE	C-N	16.36	1.62	1.33
2	H	92	CYS	CB-SG	-7.59	1.69	1.82

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	H	229	SER	CA-C-O	-31.21	54.57	120.10
3	P	309	ILE	O-C-N	-6.79	111.66	123.20
2	H	129	LYS	N-CA-C	6.13	127.56	111.00
2	H	101	ASP	CB-CG-OD1	5.75	123.47	118.30
3	P	306	ARG	NE-CZ-NH2	-5.02	117.79	120.30



All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	H	129	LYS	CA

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	H	134	SER	Peptide
3	P	309	ILE	Mainchain

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	L	1616	0	1554	21	0
2	H	1699	0	1624	9	0
3	P	134	0	143	4	0
4	H	4	0	6	0	0
4	L	4	0	6	0	0
5	H	8	0	6	2	0
5	L	4	0	3	1	0
6	H	110	0	0	1	0
6	L	71	0	0	1	0
6	P	10	0	0	1	0
All	All	3660	0	3342	32	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (32) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:L:27(B):ASN:HD22	1:L:28:VAL:H	1.16	0.91
1:L:31:ASN:OD1	3:P:315:ARG:NH2	2.14	0.80
1:L:27(B):ASN:ND2	1:L:28:VAL:H	1.90	0.69
5:L:503:ACY:H2	6:L:708:HOH:O	1.95	0.66
1:L:27(B):ASN:HD22	1:L:28:VAL:N	1.94	0.62

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:H:186:SER:HB2	5:H:502:ACY:H1	1.81	0.61
2:H:36:TRP:CE3	2:H:80:LEU:HD22	2.36	0.60
1:L:167:GLN:HE21	1:L:170:ASN:HD21	1.53	0.56
3:P:315:ARG:NH1	6:P:15:HOH:O	2.38	0.56
2:H:4:LEU:HD22	2:H:24:THR:HG22	1.88	0.54
2:H:46:GLU:OE1	6:H:716:HOH:O	2.19	0.53
1:L:26:THR:H	1:L:27(B):ASN:ND2	2.07	0.52
1:L:170:ASN:ND2	1:L:172:LYS:H	2.08	0.51
2:H:100(B):SER:O	3:P:308:HIS:HE1	1.94	0.51
1:L:31:ASN:O	1:L:66:LYS:NZ	2.45	0.49
1:L:35:TRP:HB2	1:L:48:ILE:HB	1.95	0.48
2:H:32:TYR:CZ	3:P:304:ARG:HG3	2.50	0.47
1:L:120:PRO:HD3	1:L:132:LEU:CD2	2.46	0.46
1:L:19:ILE:HD12	1:L:19:ILE:C	2.36	0.46
1:L:1:GLN:HA	1:L:26:THR:HG21	1.97	0.45
1:L:26:THR:H	1:L:27(B):ASN:HD21	1.63	0.45
1:L:212:THR:HG22	1:L:212:THR:O	2.17	0.44
1:L:178:TYR:CG	5:H:502:ACY:H3	2.53	0.44
2:H:173:THR:HA	2:H:189:SER:HA	2.00	0.43
1:L:1:GLN:NE2	1:L:27(A):SER:OG	2.44	0.43
2:H:166:LEU:HD21	2:H:191:VAL:HG21	2.01	0.43
1:L:170:ASN:O	1:L:171:ASN:HB2	2.19	0.42
2:H:147:TYR:CE1	2:H:152:VAL:HG23	2.55	0.42
1:L:170:ASN:HD22	1:L:170:ASN:C	2.23	0.41
1:L:166:LYS:HD2	1:L:171:ASN:HA	2.02	0.41
1:L:167:GLN:OE1	1:L:174:ALA:HB2	2.21	0.41
1:L:120:PRO:HD3	1:L:132:LEU:HD22	2.03	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	L	213/215 (99%)	205 (96%)	8 (4%)	0	100	100
2	H	224/226 (99%)	214 (96%)	6 (3%)	4 (2%)	9	8
3	P	14/19 (74%)	14 (100%)	0	0	100	100
All	All	451/460 (98%)	433 (96%)	14 (3%)	4 (1%)	19	22

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	H	128	SER
2	H	127	SER
2	H	130	SER
2	H	100(E)	ASP

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	L	180/180 (100%)	176 (98%)	4 (2%)	55	72
2	H	191/191 (100%)	186 (97%)	5 (3%)	49	66
3	P	13/15 (87%)	13 (100%)	0	100	100
All	All	384/386 (100%)	375 (98%)	9 (2%)	53	71

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	L	27(B)	ASN
1	L	55	SER
1	L	152	SER
1	L	170	ASN
2	H	58	ARG
2	H	62	SER
2	H	116	THR
2	H	198	LEU
2	H	216	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	L	1	GLN
1	L	27(B)	ASN
1	L	126	GLN
1	L	170	ASN
2	H	79	HIS
2	H	216	ASN
3	P	308	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
5	ACY	H	501	-	1,3,3	0.90	0	0,3,3	0.00	-
5	ACY	H	502	-	1,3,3	0.39	0	0,3,3	0.00	-
4	EDO	H	703	-	3,3,3	0.52	0	2,2,2	0.18	0
5	ACY	L	503	-	1,3,3	1.33	0	0,3,3	0.00	-
4	EDO	L	702	-	3,3,3	0.55	0	2,2,2	0.19	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	EDO	H	703	-	-	1/1/1/1	-
4	EDO	L	702	-	-	1/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	H	703	EDO	O1-C1-C2-O2
4	L	702	EDO	O1-C1-C2-O2

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
3	P	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	P	309:ILE	C	312:GLY	N	1.62

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

### 6.3 Carbohydrates [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

### 6.4 Ligands [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

### 6.5 Other polymers [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.