



Full wwPDB X-ray Structure Validation Report ⓘ

Dec 16, 2019 – 10:25 AM EST

PDB ID : 6B0F
Title : ESTROGEN RECEPTOR ALPHA LIGAND BINDING DOMAIN IN COMPLEX WITH LSZ102
Authors : Kirby, C.A.; Baird, J.
Deposited on : 2017-09-14
Resolution : 2.86 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.0 (224370), CSD as540be (2019)
Xtriage (Phenix) : 1.13
EDS : 2.4
buster-report : 1.1.7 (2018)
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)
Refmac : 5.8.0158
CCP4 : 7.0 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.4


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 2.86 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	111664	2715 (2.90-2.82)
Clashscore	122126	2976 (2.90-2.82)
Ramachandran outliers	120053	2913 (2.90-2.82)
Sidechain outliers	120020	2916 (2.90-2.82)
RSRZ outliers	108989	2654 (2.90-2.82)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	254	
1	B	254	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 3199 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Estrogen receptor.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	204	Total	C	N	O	S	0	0	0
			1624	1046	276	287	15			
1	B	185	Total	C	N	O	S	0	0	0
			1474	945	253	262	14			

There are 8 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	300	GLY	-	expression tag	UNP P03372
A	381	SER	CYS	engineered mutation	UNP P03372
A	417	SER	CYS	engineered mutation	UNP P03372
A	530	SER	CYS	engineered mutation	UNP P03372
B	300	GLY	-	expression tag	UNP P03372
B	381	SER	CYS	engineered mutation	UNP P03372
B	417	SER	CYS	engineered mutation	UNP P03372
B	530	SER	CYS	engineered mutation	UNP P03372

- Molecule 2 is LSZ102 (three-letter code: C6V) (formula: C₂₅H₁₇F₃O₄S) (labeled as "Ligand of Interest" by author).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total 33	C 25	F 3	O 4	S 1	0	0
2	B	1	Total 33	C 25	F 3	O 4	S 1	0	0

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: $\text{C}_3\text{H}_8\text{O}_3$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	10	Total	O	0	0
			10	10		
4	B	7	Total	O	0	0
			7	7		

- Molecule 1: Estrogen receptor



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	105.82Å 52.10Å 97.02Å 90.00° 114.33° 90.00°	Depositor
Resolution (Å)	45.84 – 2.86 45.84 – 2.86	Depositor EDS
% Data completeness (in resolution range)	98.6 (45.84-2.86) 98.7 (45.84-2.86)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.61 (at 2.86Å)	Xtriage
Refinement program	BUSTER 2.11.7	Depositor
R, R_{free}	0.227 , 0.299 0.234 , 0.300	Depositor DCC
R_{free} test set	538 reflections (4.80%)	wwPDB-VP
Wilson B-factor (Å ²)	49.4	Xtriage
Anisotropy	0.715	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 79.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	3199	wwPDB-VP
Average B, all atoms (Å ²)	51.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 9.42% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: C6V, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.50	0/1650	0.68	0/2225
1	B	0.50	0/1496	0.65	0/2014
All	All	0.50	0/3146	0.67	0/4239

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1624	0	1681	16	0
1	B	1474	0	1519	14	0
2	A	33	0	0	0	0
2	B	33	0	0	0	0
3	A	12	0	16	1	0
3	B	6	0	8	1	0
4	A	10	0	0	0	0
4	B	7	0	0	0	0
All	All	3199	0	3224	27	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (27) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:327:LEU:HD22	3:B:602:GOL:H12	1.55	0.87
1:B:388:MET:HG2	1:B:428:LEU:HD21	1.56	0.85
1:A:394:ARG:HE	3:A:602:GOL:H32	1.61	0.65
1:A:392:VAL:HG11	1:A:431:THR:HG22	1.86	0.58
1:A:382:ALA:HB2	1:A:456:SER:HB3	1.87	0.57
1:B:359:ASN:HA	1:B:362:LYS:HD3	1.88	0.56
1:A:448:LEU:HA	1:A:451:ILE:HD12	1.90	0.53
1:A:490:MET:HB3	1:A:500:GLN:HG2	1.92	0.52
1:B:395:SER:OG	1:B:402:LEU:HA	2.11	0.51
1:B:370:LEU:HB2	1:B:375:GLN:HG3	1.93	0.51
1:A:508:LEU:HD13	1:B:505:ALA:HA	1.94	0.49
1:A:392:VAL:HG13	1:A:432:SER:HA	1.94	0.49
1:B:328:TYR:HA	1:B:405:ALA:HB1	1.96	0.47
1:B:382:ALA:HB2	1:B:456:SER:HB3	1.97	0.47
1:A:381:SER:HB2	1:A:456:SER:HB2	1.97	0.46
1:A:505:ALA:HA	1:B:508:LEU:HD13	1.98	0.46
1:A:339:GLU:HG3	1:A:418:VAL:HG22	1.99	0.44
1:A:323:GLU:HA	1:A:324:PRO:HD3	1.86	0.44
1:B:474:HIS:O	1:B:478:VAL:HG23	2.19	0.43
1:A:329:SER:H	1:A:407:ASN:HD21	1.67	0.42
1:B:379:LEU:O	1:B:383:TRP:HB3	2.20	0.42
1:B:507:LEU:O	1:B:510:ILE:HB	2.20	0.42
1:B:324:PRO:HB3	1:B:356:HIS:HB3	2.02	0.41
1:A:391:LEU:HD23	1:A:428:LEU:HD22	2.03	0.41
1:A:497:LEU:HD22	1:B:497:LEU:HD22	2.02	0.40
1:A:396:MET:HG2	1:A:436:ARG:HB2	2.03	0.40
1:A:511:LEU:HD23	1:A:514:ILE:HD12	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	194/254 (76%)	188 (97%)	6 (3%)	0	100	100
1	B	175/254 (69%)	170 (97%)	4 (2%)	1 (1%)	27	57
All	All	369/508 (73%)	358 (97%)	10 (3%)	1 (0%)	43	71

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	365	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	183/228 (80%)	178 (97%)	5 (3%)	48	77
1	B	165/228 (72%)	160 (97%)	5 (3%)	44	75
All	All	348/456 (76%)	338 (97%)	10 (3%)	45	76

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	345	LEU
1	A	379	LEU
1	A	475	ILE
1	A	532	ASN
1	A	544	LEU
1	B	355	VAL
1	B	395	SER
1	B	409	LEU
1	B	495	LEU
1	B	497	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	C6V	A	601	-	28,36,36	2.29	6 (21%)	32,53,53	1.40	5 (15%)
3	GOL	A	602	-	5,5,5	0.07	0	5,5,5	0.18	0
3	GOL	A	603	-	5,5,5	0.07	0	5,5,5	0.17	0
2	C6V	B	601	-	28,36,36	2.23	7 (25%)	32,53,53	1.37	4 (12%)
3	GOL	B	602	-	5,5,5	0.21	0	5,5,5	0.32	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	C6V	A	601	-	-	5/12/19/19	0/4/4/4
3	GOL	A	602	-	-	0/4/4/4	-
3	GOL	A	603	-	-	0/4/4/4	-
2	C6V	B	601	-	-	5/12/19/19	0/4/4/4

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	B	602	-	-	3/4/4/4	-

All (13) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	601	C6V	C10-C6	8.51	1.51	1.40
2	A	601	C6V	C10-C6	8.48	1.51	1.40
2	A	601	C6V	C13-C9	3.79	1.54	1.48
2	A	601	C6V	C30-C24	3.45	1.57	1.51
2	B	601	C6V	C13-C9	2.98	1.52	1.48
2	B	601	C6V	C30-C24	2.90	1.56	1.51
2	B	601	C6V	O12-C10	-2.41	1.34	1.39
2	B	601	C6V	C7-C2	2.25	1.41	1.36
2	A	601	C6V	C7-C2	2.25	1.41	1.36
2	B	601	C6V	C4-C3	2.21	1.41	1.37
2	A	601	C6V	C4-C3	2.17	1.41	1.37
2	A	601	C6V	O12-C10	-2.08	1.35	1.39
2	B	601	C6V	C13-C24	2.07	1.44	1.41

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	601	C6V	C24-C25-C26	4.44	123.66	117.58
2	B	601	C6V	C28-C13-C24	3.08	121.04	118.51
2	B	601	C6V	C24-C25-C26	2.98	121.65	117.58
2	A	601	C6V	C27-C26-C25	-2.66	119.84	123.30
2	A	601	C6V	C25-C24-C13	-2.38	116.47	119.67
2	A	601	C6V	C2-C7-C6	-2.26	118.05	121.17
2	B	601	C6V	C25-C24-C30	2.24	119.04	115.13
2	A	601	C6V	C25-C24-C30	2.12	118.83	115.13
2	B	601	C6V	C25-C24-C13	-2.06	116.90	119.67

There are no chirality outliers.

All (13) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	601	C6V	C28-C13-C9-S8
2	B	601	C6V	C13-C24-C30-F31
2	B	601	C6V	C13-C24-C30-F33
2	A	601	C6V	C28-C13-C9-S8
2	A	601	C6V	C13-C24-C30-F31

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
2	A	601	C6V	C25-C24-C30-F31
3	B	602	GOL	O1-C1-C2-C3
3	B	602	GOL	C1-C2-C3-O3
2	B	601	C6V	C25-C24-C30-F31
2	B	601	C6V	C25-C24-C30-F33
3	B	602	GOL	O1-C1-C2-O2
2	A	601	C6V	C20-C1-C17-C16
2	A	601	C6V	C20-C1-C17-C18


There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	602	GOL	1	0
3	B	602	GOL	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	204/254 (80%)	0.34	8 (3%) 39 33	33, 50, 81, 108	0
1	B	185/254 (72%)	0.25	3 (1%) 72 70	30, 48, 72, 94	0
All	All	389/508 (76%)	0.29	11 (2%) 53 48	30, 49, 77, 108	0

All (11) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	418	VAL	3.7
1	A	530	SER	3.6
1	B	365	PRO	3.0
1	A	501	HIS	2.8
1	B	411	ASP	2.8
1	B	345	LEU	2.7
1	A	410	LEU	2.4
1	A	438	MET	2.2
1	A	419	GLU	2.2
1	A	429	LEU	2.1
1	A	537	TYR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	GOL	A	603	6/6	0.80	0.24	57,62,64,66	0
3	GOL	A	602	6/6	0.85	0.21	66,67,69,69	0
2	C6V	B	601	33/33	0.94	0.25	41,47,58,61	0
2	C6V	A	601	33/33	0.95	0.30	38,45,50,54	0
3	GOL	B	602	6/6	0.95	0.29	34,35,41,45	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around C6V B 601:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around C6V A 601:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.