



# wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 9, 2018 – 02:03 am GMT

PDB ID : 3B5N  
Title : Structure of the yeast plasma membrane SNARE complex  
Authors : Strop, P.; Brunger, A.T.  
Deposited on : 2007-10-26  
Resolution : 1.60 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : trunk30967  
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)  
Refmac : 5.8.0158  
CCP4 : 7.0 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk30967


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	111664	2957 (1.60-1.60)
Clashscore	122126	3202 (1.60-1.60)
Ramachandran outliers	120053	3117 (1.60-1.60)
Sidechain outliers	120020	3116 (1.60-1.60)
RSRZ outliers	108989	2883 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	61	<div> <div>2%</div> <div> <div></div> <div>95%</div> <div>5%</div> </div> </div>
1	E	61	<div> <div></div> <div> <div>87%</div> <div>8%</div> <div>5%</div> </div> </div>
1	I	61	<div> <div>25%</div> <div> <div></div> <div>79%</div> <div>16%</div> <div>5%</div> </div> </div>
2	B	69	<div> <div>3%</div> <div> <div></div> <div>97%</div> <div></div> </div> </div>
2	F	69	<div> <div>3%</div> <div> <div></div> <div>91%</div> <div>7%</div> <div></div> </div> </div>
2	J	69	<div> <div>16%</div> <div> <div></div> <div>67%</div> <div>25%</div> <div>7%</div> </div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	C	70	<div><div>%</div><div><div></div><div>87%</div><div>13%</div></div></div>
3	G	70	<div><div>4%</div><div><div></div><div>83%</div><div>16%</div><div></div></div><div></div></div>
3	K	70	<div><div>11%</div><div><div></div><div>77%</div><div>16%</div><div>7%</div></div><div></div></div>
4	D	64	<div><div></div><div><div></div><div>95%</div><div>5%</div></div></div>
4	H	64	<div><div></div><div><div></div><div>83%</div><div>14%</div><div></div></div><div></div></div>
4	L	64	<div><div>11%</div><div><div></div><div>81%</div><div>11%</div><div>5%</div><div></div></div><div></div></div>

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 6656 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Synaptobrevin homolog 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	61	Total	C	N	O	S	0	0	0
			473	284	93	94	2			
1	E	58	Total	C	N	O	S	0	0	0
			452	273	87	90	2			
1	I	58	Total	C	N	O	S	0	0	0
			452	273	87	90	2			

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	26	GLY	-	EXPRESSION TAG	UNP P31109
E	26	GLY	-	EXPRESSION TAG	UNP P31109
I	26	GLY	-	EXPRESSION TAG	UNP P31109

- Molecule 2 is a protein called Protein SSO1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	69	Total	C	N	O	S	0	0	0
			545	333	94	116	2			
2	F	68	Total	C	N	O	S	0	0	0
			533	327	90	114	2			
2	J	64	Total	C	N	O	S	0	0	0
			509	312	86	109	2			

- Molecule 3 is a protein called Protein transport protein SEC9.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	70	Total	C	N	O	S	0	1	0
			542	329	101	108	4			
3	G	69	Total	C	N	O	S	0	2	0
			532	324	97	107	4			

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	K	65	Total	C	N	O	S	0	1	0
			503	305	93	101	4			

There are 9 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	431	GLY	-	EXPRESSION TAG	UNP P40357
C	432	SER	-	EXPRESSION TAG	UNP P40357
C	500	GLN	-	EXPRESSION TAG	UNP P40357
G	431	GLY	-	EXPRESSION TAG	UNP P40357
G	432	SER	-	EXPRESSION TAG	UNP P40357
G	500	GLN	-	EXPRESSION TAG	UNP P40357
K	431	GLY	-	EXPRESSION TAG	UNP P40357
K	432	SER	-	EXPRESSION TAG	UNP P40357
K	500	GLN	-	EXPRESSION TAG	UNP P40357

- Molecule 4 is a protein called Protein transport protein SEC9.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	D	64	Total	C	N	O	S	0	0	0
			511	305	94	109	3			
4	H	63	Total	C	N	O	S	0	1	0
			514	308	96	107	3			
4	L	62	Total	C	N	O	S	0	0	0
			498	297	92	106	3			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
D	587	GLY	-	EXPRESSION TAG	UNP P40357
D	588	SER	-	EXPRESSION TAG	UNP P40357
H	587	GLY	-	EXPRESSION TAG	UNP P40357
H	588	SER	-	EXPRESSION TAG	UNP P40357
L	587	GLY	-	EXPRESSION TAG	UNP P40357
L	588	SER	-	EXPRESSION TAG	UNP P40357

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	46	Total	O	0	0
			46	46		

*Continued on next page...*


*Continued from previous page...*

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	59	Total 59	O 59	0	0
5	C	53	Total 53	O 53	0	0
5	D	57	Total 57	O 57	0	0
5	E	54	Total 54	O 54	0	0
5	F	78	Total 78	O 78	0	0
5	G	58	Total 58	O 58	0	0
5	H	72	Total 72	O 72	0	0
5	I	25	Total 25	O 25	0	0
5	J	19	Total 19	O 19	0	0
5	K	28	Total 28	O 28	0	0
5	L	43	Total 43	O 43	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Synaptobrevin homolog 1




- Molecule 1: Synaptobrevin homolog 1




- Molecule 1: Synaptobrevin homolog 1



- Molecule 2: Protein SSO1




- Molecule 2: Protein SSO1




- Molecule 2: Protein SSO1








- Molecule 3: Protein transport protein SEC9



- Molecule 3: Protein transport protein SEC9




- Molecule 3: Protein transport protein SEC9





- Molecule 4: Protein transport protein SEC9



- Molecule 4: Protein transport protein SEC9



- Molecule 4: Protein transport protein SEC9





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	76.96Å 48.11Å 110.27Å 90.00° 97.77° 90.00°	Depositor
Resolution (Å)	34.57 – 1.60 34.57 – 1.60	Depositor EDS
% Data completeness (in resolution range)	85.0 (34.57-1.60) 85.0 (34.57-1.60)	Depositor EDS
$R_{merge}$	0.05	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.89 (at 1.60Å)	Xtriage
Refinement program	PHENIX	Depositor
R, $R_{free}$	0.206 , 0.248 0.210 , 0.249	Depositor DCC
$R_{free}$ test set	4522 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	23.4	Xtriage
Anisotropy	0.597	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.38 , 56.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	6656	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	44.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 9.09% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.27	0/475	0.39	0/633
1	E	0.31	0/454	0.45	0/606
1	I	0.24	0/454	0.39	0/606
2	B	0.28	0/547	0.38	0/734
2	F	0.33	0/535	0.41	0/720
2	J	0.23	0/511	0.35	0/687
3	C	0.29	0/546	0.39	0/725
3	G	0.31	0/539	0.38	0/718
3	K	0.25	0/507	0.37	0/675
4	D	0.34	0/511	0.41	0/682
4	H	0.31	0/517	0.45	0/691
4	L	0.26	0/498	0.35	0/666
All	All	0.29	0/6094	0.39	0/8143

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	473	0	473	3	0
1	E	452	0	452	6	0
1	I	452	0	452	12	0
2	B	545	0	536	3	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	F	533	0	523	5	0
2	J	509	0	497	20	0
3	C	542	0	562	9	0
3	G	532	0	552	16	0
3	K	503	0	517	10	0
4	D	511	0	509	3	0
4	H	514	0	519	21	0
4	L	498	0	495	13	0
5	A	46	0	0	0	0
5	B	59	0	0	3	0
5	C	53	0	0	4	0
5	D	57	0	0	0	0
5	E	54	0	0	1	0
5	F	78	0	0	0	0
5	G	58	0	0	3	0
5	H	72	0	0	3	0
5	I	25	0	0	1	0
5	J	19	0	0	1	0
5	K	28	0	0	0	0
5	L	43	0	0	2	0
All	All	6656	0	6087	84	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 84 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:H:625[A]:ARG:HG2	4:H:625[A]:ARG:HH11	1.14	1.10
3:G:473:ASN:HA	4:H:625[A]:ARG:HH21	1.18	1.07
2:J:244:GLN:HB2	5:J:260:HOH:O	1.81	0.79
1:I:83:LYS:HE2	5:I:96:HOH:O	1.84	0.76
4:H:625[A]:ARG:HG2	4:H:625[A]:ARG:NH1	1.95	0.75

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	59/61 (97%)	59 (100%)	0	0	100	100
1	E	56/61 (92%)	56 (100%)	0	0	100	100
1	I	56/61 (92%)	56 (100%)	0	0	100	100
2	B	67/69 (97%)	67 (100%)	0	0	100	100
2	F	66/69 (96%)	66 (100%)	0	0	100	100
2	J	62/69 (90%)	61 (98%)	1 (2%)	0	100	100
3	C	69/70 (99%)	69 (100%)	0	0	100	100
3	G	69/70 (99%)	69 (100%)	0	0	100	100
3	K	64/70 (91%)	64 (100%)	0	0	100	100
4	D	62/64 (97%)	62 (100%)	0	0	100	100
4	H	62/64 (97%)	62 (100%)	0	0	100	100
4	L	60/64 (94%)	58 (97%)	1 (2%)	1 (2%)	10	1
All	All	752/792 (95%)	749 (100%)	2 (0%)	1 (0%)	53	30

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
4	L	648	ALA

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	48/48 (100%)	48 (100%)	0	100	100
1	E	46/48 (96%)	46 (100%)	0	100	100
1	I	46/48 (96%)	46 (100%)	0	100	100
2	B	60/60 (100%)	60 (100%)	0	100	100
2	F	59/60 (98%)	59 (100%)	0	100	100
2	J	57/60 (95%)	56 (98%)	1 (2%)	62	38
3	C	60/59 (102%)	57 (95%)	3 (5%)	27	7
3	G	60/59 (102%)	60 (100%)	0	100	100
3	K	56/59 (95%)	55 (98%)	1 (2%)	62	38
4	D	59/59 (100%)	59 (100%)	0	100	100
4	H	60/59 (102%)	58 (97%)	2 (3%)	41	15
4	L	58/59 (98%)	56 (97%)	2 (3%)	40	15
All	All	669/678 (99%)	660 (99%)	9 (1%)	76	53

5 of 9 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
4	H	625[B]	ARG
4	L	623	GLN
3	K	486	ASN
3	C	473	ASN
2	J	241	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 17 such sidechains are listed below:

Mol	Chain	Res	Type
1	E	33	GLN
2	F	215	ASN
2	J	224	GLN
4	D	641	HIS
3	K	478	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	61/61 (100%)	0.21	1 (1%) 72 72	27, 36, 54, 58	0
1	E	58/61 (95%)	-0.23	0 100 100	25, 33, 52, 69	0
1	I	58/61 (95%)	0.99	15 (25%) 0 0	28, 55, 109, 138	0
2	B	69/69 (100%)	-0.13	2 (2%) 51 49	26, 36, 55, 70	0
2	F	68/69 (98%)	-0.12	2 (2%) 51 49	24, 32, 55, 77	0
2	J	64/69 (92%)	0.77	11 (17%) 1 1	34, 59, 101, 122	0
3	C	70/70 (100%)	-0.11	1 (1%) 75 75	24, 33, 51, 67	0
3	G	69/70 (98%)	0.09	3 (4%) 35 32	23, 32, 60, 92	0
3	K	65/70 (92%)	0.52	8 (12%) 4 3	31, 48, 85, 110	0
4	D	64/64 (100%)	0.03	0 100 100	20, 34, 47, 51	0
4	H	63/64 (98%)	-0.15	0 100 100	25, 34, 47, 65	0
4	L	62/64 (96%)	0.44	7 (11%) 5 4	27, 43, 90, 100	0
All	All	771/792 (97%)	0.18	50 (6%) 19 17	20, 38, 79, 138	0

The worst 5 of 50 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	G	499	LEU	7.4
1	I	86	TRP	6.0
3	K	496	LEU	5.5
2	J	254	LYS	5.3
3	K	497	LYS	5.2

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

### 6.5 Other polymers [i](#)

There are no such residues in this entry.