



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 11, 2018 – 12:50 am GMT

PDB ID : 6B5K
Title : Mycobacterium tuberculosis RmlA in complex with Mg/dTTP
Authors : Brown, H.A.; Holden, H.A.
Deposited on : 2017-09-29
Resolution : 1.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.3 (157068), CSD as539be (2018)
Xtriage (Phenix) : 1.13
EDS : trunk30967
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)
Refmac : 5.8.0158
CCP4 : 7.0 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk30967


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	111664	2957 (1.60-1.60)
Clashscore	122126	3202 (1.60-1.60)
Ramachandran outliers	120053	3117 (1.60-1.60)
Sidechain outliers	120020	3116 (1.60-1.60)
RSRZ outliers	108989	2883 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	296	<div> <div>4%</div> <div> <div></div> <div>85%</div> <div>8%</div> <div>6%</div> </div> </div>
1	B	296	<div> <div>8%</div> <div> <div></div> <div>81%</div> <div>9%</div> <div>9%</div> </div> </div>

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 4682 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Glucose-1-phosphate thymidyltransferase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	277	Total	C	N	O	S	0	4	0
			2145	1380	358	401	6			
1	B	270	Total	C	N	O	S	0	0	0
			2072	1331	348	387	6			

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	289	LEU	-	expression tag	UNP P9WH13
A	290	GLU	-	expression tag	UNP P9WH13
A	291	HIS	-	expression tag	UNP P9WH13
A	292	HIS	-	expression tag	UNP P9WH13
A	293	HIS	-	expression tag	UNP P9WH13
A	294	HIS	-	expression tag	UNP P9WH13
A	295	HIS	-	expression tag	UNP P9WH13
A	296	HIS	-	expression tag	UNP P9WH13
B	289	LEU	-	expression tag	UNP P9WH13
B	290	GLU	-	expression tag	UNP P9WH13
B	291	HIS	-	expression tag	UNP P9WH13
B	292	HIS	-	expression tag	UNP P9WH13
B	293	HIS	-	expression tag	UNP P9WH13
B	294	HIS	-	expression tag	UNP P9WH13
B	295	HIS	-	expression tag	UNP P9WH13
B	296	HIS	-	expression tag	UNP P9WH13

- Molecule 2 is THYMIDINE-5'-TRIPHOSPHATE (three-letter code: TTP) (formula: C₁₀H₁₇N₂O₁₄P₃).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			29	10	2	14	3		
2	B	1	Total	C	N	O	P	0	0
			29	10	2	14	3		

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	2	Total	Mg	0	0
			2	2		
3	A	2	Total	Mg	0	0
			2	2		

- Molecule 4 is THYMIDINE-5'-DIPHOSPHATE (three-letter code: TYD) (formula: C₁₀H₁₆N₂O₁₁P₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	P	0	0
			25	10	2	11	2		
4	B	1	Total	C	N	O	P	0	0
			25	10	2	11	2		

- Molecule 5 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: $C_2H_6O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			4	2	2		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	231	Total 231	O 231	0	0
6	B	118	Total 118	O 118	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Glucose-1-phosphate thymidyltransferase



- Molecule 1: Glucose-1-phosphate thymidyltransferase



4 Data and refinement statistics

Property	Value	Source
Space group	I 4	Depositor
Cell constants a, b, c, α , β , γ	96.40Å 96.40Å 151.92Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 1.60 25.37 – 1.60	Depositor EDS
% Data completeness (in resolution range)	98.4 (50.00-1.60) 98.5 (25.37-1.60)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.05	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.27 (at 1.60Å)	Xtriage
Refinement program	REFMAC 5.8.0158	Depositor
R, R_{free}	0.195 , 0.227 0.203 , 0.232	Depositor DCC
R_{free} test set	4414 reflections (4.94%)	wwPDB-VP
Wilson B-factor (Å ²)	20.0	Xtriage
Anisotropy	0.037	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 44.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.43$, $\langle L^2 \rangle = 0.26$	Xtriage
Estimated twinning fraction	0.116 for h,-k,-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	4682	wwPDB-VP
Average B, all atoms (Å ²)	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.97% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: TYD, MG, TTP, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.76	0/2202	0.87	3/2992 (0.1%)
1	B	0.50	0/2114	0.71	2/2871 (0.1%)
All	All	0.65	0/4316	0.80	5/5863 (0.1%)

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	67	ASP	CB-CG-OD1	6.14	123.83	118.30
1	A	2	ARG	NE-CZ-NH1	-5.87	117.37	120.30
1	B	237	ARG	NE-CZ-NH1	-5.56	117.52	120.30
1	A	100	ASP	CB-CG-OD1	5.26	123.03	118.30
1	B	237	ARG	NE-CZ-NH2	5.16	122.88	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2145	0	2129	13	0
1	B	2072	0	2052	19	0
2	A	29	0	13	0	0
2	B	29	0	13	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	2	0	0	0	0
3	B	2	0	0	0	0
4	A	25	0	13	0	0
4	B	25	0	13	0	0
5	A	4	0	6	0	0
6	A	231	0	0	4	0
6	B	118	0	0	1	0
All	All	4682	0	4239	31	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (31) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:21:ILE:HD13	1:A:26:LEU:HD23	1.45	0.97
1:B:192:GLY:N	6:B:401:HOH:O	2.12	0.83
6:A:452:HOH:O	1:B:221:LEU:HD21	1.80	0.80
6:A:452:HOH:O	1:B:221:LEU:CD2	2.39	0.70
1:B:206:GLN:OE1	1:B:208:ARG:NH2	2.24	0.70
1:B:257:MET:HG2	1:B:259:TRP:CZ2	2.30	0.66
1:B:39:THR:CG2	1:B:253:VAL:HG21	2.30	0.62
1:B:99:ALA:C	1:B:177:ASN:ND2	2.56	0.59
1:B:39:THR:HG23	1:B:253:VAL:HG21	1.86	0.58
1:A:21:ILE:HD13	1:A:26:LEU:CD2	2.26	0.56
1:A:145:VAL:HG22	1:A:169:VAL:CG2	2.38	0.54
1:B:5:ILE:HD12	1:B:51:LEU:HD11	1.91	0.53
1:A:39[A]:THR:CG2	1:A:253:VAL:HG21	2.39	0.52
1:A:242:ARG:NH1	6:A:401:HOH:O	2.27	0.51
1:B:116:LEU:HD21	1:B:214:LEU:HD22	1.94	0.50
1:A:145:VAL:HG22	1:A:169:VAL:HG22	1.94	0.49
1:B:21:ILE:HD13	1:B:26:LEU:HD23	1.94	0.49
1:B:268:ARG:CZ	1:B:268:ARG:HA	2.45	0.47
1:B:285:LEU:O	1:B:286:GLU:C	2.53	0.47
1:A:145:VAL:O	1:A:165:SER:OG	2.22	0.46
1:A:242:ARG:HE	1:B:221:LEU:HD21	1.80	0.46
1:A:276:GLY:HA3	6:A:407:HOH:O	2.15	0.45
1:B:50:GLN:HE22	1:B:93:GLY:HA2	1.80	0.45
1:B:50:GLN:HE22	1:B:93:GLY:CA	2.29	0.45
1:A:126:ILE:HG23	1:A:126:ILE:O	2.17	0.44
1:B:198:GLU:O	1:B:202:VAL:HG23	2.17	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:156:LEU:O	1:A:157:GLU:HG2	2.18	0.43
1:B:257:MET:HG2	1:B:259:TRP:CE2	2.55	0.42
1:A:263:GLU:O	1:A:267:GLN:HG2	2.20	0.41
1:A:272:LEU:HD13	1:A:277:TYR:CD2	2.56	0.40
1:B:264:GLN:O	1:B:268:ARG:HG2	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	275/296 (93%)	269 (98%)	5 (2%)	1 (0%)	36	16
1	B	262/296 (88%)	256 (98%)	5 (2%)	1 (0%)	36	16
All	All	537/592 (91%)	525 (98%)	10 (2%)	2 (0%)	36	16

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	29	TYR
1	B	29	TYR

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	221/237 (93%)	215 (97%)	6 (3%)	48	21
1	B	212/237 (90%)	205 (97%)	7 (3%)	41	15
All	All	433/474 (91%)	420 (97%)	13 (3%)	44	18

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	109	ASN
1	A	145	VAL
1	A	154	LEU
1	A	155	SER
1	A	208	ARG
1	A	257	MET
1	B	21	ILE
1	B	109	ASN
1	B	119	SER
1	B	145	VAL
1	B	197	THR
1	B	208	ARG
1	B	257	MET

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	166	ASN
1	B	50	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry

Of 9 ligands modelled in this entry, 4 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	TTP	A	301	3	22,30,30	1.00	0	27,47,47	2.13	1 (3%)
4	TYD	A	304	-	24,26,26	2.27	5 (20%)	30,40,40	1.92	8 (26%)
5	EDO	A	305	-	3,3,3	0.61	0	2,2,2	0.13	0
2	TTP	B	303	3	22,30,30	0.96	1 (4%)	27,47,47	1.96	5 (18%)
4	TYD	B	304	-	24,26,26	2.50	4 (16%)	30,40,40	1.93	7 (23%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	TTP	A	301	3	-	0/18/34/34	0/2/2/2
4	TYD	A	304	-	-	0/16/28/28	0/2/2/2
5	EDO	A	305	-	-	0/1/1/1	0/0/0/0
2	TTP	B	303	3	-	0/18/34/34	0/2/2/2
4	TYD	B	304	-	-	0/16/28/28	0/2/2/2

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	304	TYD	C6-N1	-10.03	1.33	1.46
4	A	304	TYD	C6-N1	-8.54	1.35	1.46
4	B	304	TYD	C6-C5	-4.31	1.38	1.51
4	A	304	TYD	C6-C5	-3.86	1.39	1.51
4	B	304	TYD	C2-N3	-2.29	1.33	1.38
4	A	304	TYD	C2-N3	-2.02	1.34	1.38
4	A	304	TYD	PB-O3A	2.02	1.63	1.60
2	B	303	TTP	PG-O3B	2.99	1.64	1.60
4	B	304	TYD	C2-N1	3.44	1.40	1.35

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	304	TYD	C2-N1	4.15	1.41	1.35

All (21) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	304	TYD	C4-N3-C2	-5.25	120.44	126.86
4	B	304	TYD	C4-N3-C2	-4.90	120.87	126.86
4	A	304	TYD	O2-C2-N1	-4.69	117.34	123.12
4	B	304	TYD	O2-C2-N1	-3.66	118.60	123.12
4	A	304	TYD	C2'-C1'-N1	-2.71	112.30	115.61
2	B	303	TTP	O4'-C1'-N1	-2.54	103.51	107.78
4	B	304	TYD	O3A-PB-O1B	-2.41	96.96	111.48
2	B	303	TTP	C5-C6-N1	-2.02	119.96	122.15
4	A	304	TYD	C6-C5-C4	2.02	117.31	111.53
4	A	304	TYD	O2A-PA-O1A	2.02	122.42	112.14
4	A	304	TYD	O3B-PB-O2B	2.15	116.10	107.59
4	B	304	TYD	C6-C5-C4	2.44	118.54	111.53
2	B	303	TTP	O3G-PG-O2G	2.48	117.40	107.59
2	B	303	TTP	O2G-PG-O1G	2.49	120.30	110.60
4	B	304	TYD	O4'-C1'-N1	2.95	112.17	108.41
4	B	304	TYD	C5M-C5-C6	3.33	119.59	112.41
4	A	304	TYD	N3-C2-N1	3.46	120.15	116.73
4	A	304	TYD	C5M-C5-C6	3.95	120.92	112.41
4	B	304	TYD	N3-C2-N1	5.03	121.69	116.73
2	B	303	TTP	C4-N3-C2	7.42	121.46	115.14
2	A	301	TTP	C4-N3-C2	9.53	123.26	115.14

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	277/296 (93%)	-0.00	11 (3%) 38 36	12, 21, 48, 71	0
1	B	270/296 (91%)	0.46	24 (8%) 9 8	18, 38, 67, 81	0
All	All	547/592 (92%)	0.23	35 (6%) 19 18	12, 28, 62, 81	0

All (35) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	163	PRO	4.5
1	A	147	PHE	4.2
1	B	207	GLY	4.1
1	B	190	ALA	4.1
1	A	166	ASN	4.1
1	B	186	LEU	3.9
1	A	135	TRP	3.8
1	A	154	LEU	3.8
1	B	285	LEU	3.7
1	B	187	LYS	3.6
1	B	181	GLU	3.5
1	B	203	TYR	3.2
1	B	197	THR	3.1
1	B	135	TRP	3.1
1	B	267	GLN	3.0
1	B	205	ASN	2.9
1	A	141	ALA	2.9
1	B	91	VAL	2.8
1	B	194	TYR	2.7
1	B	270	ARG	2.6
1	A	142	TYR	2.5
1	B	273	VAL	2.5
1	B	126	ILE	2.5
1	B	155	SER	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	99	ALA	2.4
1	B	165	SER	2.3
1	A	146	GLU	2.2
1	B	198	GLU	2.2
1	B	210	ALA	2.2
1	A	136	VAL	2.1
1	B	189	SER	2.1
1	A	139	PRO	2.1
1	B	204	LEU	2.1
1	A	165	SER	2.1
1	B	82	GLN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	TYD	A	304	25/25	0.90	0.17	28,37,46,51	0
5	EDO	A	305	4/4	0.93	0.07	32,32,34,35	0
4	TYD	B	304	25/25	0.95	0.09	30,37,44,50	0
3	MG	B	302	1/1	0.96	0.07	24,24,24,24	0
2	TTP	B	303	29/29	0.97	0.10	23,26,32,34	0
3	MG	B	301	1/1	0.97	0.04	28,28,28,28	0
3	MG	A	302	1/1	0.99	0.07	13,13,13,13	0
2	TTP	A	301	29/29	0.99	0.08	11,13,14,17	0
3	MG	A	303	1/1	0.99	0.05	15,15,15,15	0

6.5 Other polymers [i](#)

There are no such residues in this entry.