



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2018 – 11:00 pm GMT

PDB ID : 2B7R  
Title : Structure of E378D mutant flavocytochrome c3  
Authors : Pankhurst, K.L.; Mowat, C.G.; Rothery, E.L.; Miles, C.S.; Walkinshaw, M.D.;  
Reid, G.A.; Chapman, S.K.  
Deposited on : 2005-10-05  
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.3 (157068), CSD as539be (2018)  
Xtriage (Phenix) : 1.13  
EDS : trunk30967  
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)  
Refmac : 5.8.0158  
CCP4 : 7.0 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk30967


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	111664	3793 (1.70-1.70)
Clashscore	122126	4167 (1.70-1.70)
Ramachandran outliers	120053	4100 (1.70-1.70)
Sidechain outliers	120020	4100 (1.70-1.70)
RSRZ outliers	108989	3718 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	571	<div> <div style="width: 94%;"></div> <div style="width: 5%;"></div> <div style="width: 1%;"></div> </div> <div>94% 5% ..</div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	FAD	A	1805	X	-	-	-

## 2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 5494 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Fumarate reductase flavoprotein subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	568	Total	C	N	O	S	0	0	0
			4176	2592	736	823	25			


There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	378	ASP	GLU	ENGINEERED	UNP Q02469

- Molecule 2 is SODIUM ION (three-letter code: NA) (formula: Na).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Na	0	0
			1	1		

- Molecule 3 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula:  $C_{34}H_{32}FeN_4O_4$ ).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
3	A	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
3	A	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
3	A	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
3	A	1	Total 43	C 34	Fe 1	N 4	O 4	0	0

- Molecule 4 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula:  $\text{C}_{27}\text{H}_{33}\text{N}_9\text{O}_{15}\text{P}_2$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		

- Molecule 5 is FUMARIC ACID (three-letter code: FUM) (formula:  $C_4H_4O_4$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			8	4	4		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1084	Total	O	0	0
			1084	1084		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Fumarate reductase flavoprotein subunit



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	45.96Å 92.88Å 79.49Å 90.00° 91.17° 90.00°	Depositor
Resolution (Å)	20.00 – 1.70 19.72 – 1.70	Depositor EDS
% Data completeness (in resolution range)	98.2 (20.00-1.70) 98.2 (19.72-1.70)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.43 (at 1.69Å)	Xtriage
Refinement program	REFMAC 5.2.0003	Depositor
R, $R_{free}$	0.156 , 0.180 0.155 , 0.179	Depositor DCC
$R_{free}$ test set	3629 reflections (4.94%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	14.5	Xtriage
Anisotropy	0.060	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.40 , 72.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.024 for h,-k,-l	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	5494	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	18.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.60% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: HEM, FUM, FAD, NA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.69	0/4246	0.74	6/5746 (0.1%)

There are no bond length outliers.

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	2	ASP	CB-CG-OD2	6.62	124.26	118.30
1	A	378	ASP	CB-CG-OD2	6.06	123.75	118.30
1	A	552	ASP	CB-CG-OD1	5.78	123.50	118.30
1	A	128	ASP	CB-CG-OD2	5.45	123.20	118.30
1	A	422	ASP	CB-CG-OD2	5.17	122.95	118.30
1	A	27	ASP	CB-CG-OD2	5.15	122.93	118.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4176	0	4074	41	0
2	A	1	0	0	0	0
3	A	172	0	120	20	0
4	A	53	0	29	8	0
5	A	8	0	2	1	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	A	1084	0	0	3	0
All	All	5494	0	4225	41	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (41) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:14:CYS:SG	3:A:801:HEM:HAB	1.62	1.38
1:A:82:CYS:SG	3:A:804:HEM:HAB	1.61	1.32
1:A:71:CYS:SG	3:A:803:HEM:HAC	1.91	1.07
1:A:375:MET:CE	4:A:1805:FAD:H6	1.86	1.05
1:A:375:MET:HE2	4:A:1805:FAD:H6	1.35	1.02
1:A:14:CYS:HG	3:A:801:HEM:CAB	1.58	1.01
1:A:17:CYS:SG	3:A:801:HEM:HAC	1.99	1.01
1:A:229:ASP:H	1:A:256:HIS:HE1	1.14	0.92
1:A:85:CYS:SG	3:A:804:HEM:HAC	2.16	0.86
1:A:39:CYS:HG	3:A:802:HEM:CAC	1.91	0.83
1:A:169:ALA:O	6:A:2598:HOH:O	1.96	0.82
1:A:375:MET:CE	4:A:1805:FAD:C6	2.56	0.82
1:A:204:ASN:H	1:A:204:ASN:HD22	1.28	0.81
1:A:229:ASP:H	1:A:256:HIS:CE1	2.03	0.75
1:A:378:ASP:OD1	1:A:381:ARG:NH2	2.25	0.68
1:A:82:CYS:SG	3:A:804:HEM:CBB	2.79	0.68
1:A:200:GLY:HA3	1:A:204:ASN:HD21	1.60	0.66
1:A:82:CYS:SG	3:A:804:HEM:C3B	2.91	0.62
1:A:375:MET:HE1	4:A:1805:FAD:H6	1.78	0.60
1:A:204:ASN:ND2	1:A:204:ASN:H	1.98	0.59
1:A:375:MET:HE2	4:A:1805:FAD:C6	2.20	0.59
1:A:17:CYS:SG	3:A:801:HEM:C3C	2.96	0.58
1:A:71:CYS:SG	3:A:803:HEM:CBC	2.91	0.57
1:A:85:CYS:SG	3:A:804:HEM:C3C	3.02	0.53
1:A:177:THR:OG1	1:A:245:HIS:HE1	1.91	0.52
1:A:71:CYS:SG	3:A:803:HEM:C3C	3.00	0.52
1:A:402:ARG:HH22	5:A:1806:FUM:C4	2.24	0.51
1:A:14:CYS:SG	3:A:801:HEM:CBB	2.91	0.50
1:A:82:CYS:CB	3:A:804:HEM:HAB	2.39	0.49
1:A:85:CYS:SG	3:A:804:HEM:CBC	2.97	0.49
1:A:375:MET:HE1	4:A:1805:FAD:C6	2.38	0.49
1:A:229:ASP:N	1:A:256:HIS:HE1	1.96	0.48

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:77:LYS:HG2	6:A:2151:HOH:O	2.13	0.47
1:A:17:CYS:SG	3:A:801:HEM:CBC	2.97	0.46
1:A:39:CYS:SG	3:A:802:HEM:C3C	3.07	0.46
1:A:566:LYS:HG3	6:A:2235:HOH:O	2.17	0.45
1:A:14:CYS:SG	3:A:801:HEM:C3B	2.99	0.44
1:A:39:CYS:HG	3:A:802:HEM:CBC	2.30	0.44
1:A:168:ALA:HA	4:A:1805:FAD:N5	2.35	0.42
1:A:375:MET:SD	4:A:1805:FAD:C6	3.07	0.42
1:A:204:ASN:ND2	1:A:204:ASN:N	2.69	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	566/571 (99%)	552 (98%)	14 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	429/445 (96%)	421 (98%)	8 (2%)	60	43

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	56	ASN
1	A	77	LYS
1	A	204	ASN
1	A	335	SER
1	A	337	ASN
1	A	401	THR
1	A	490	ASN
1	A	559	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	56	ASN
1	A	116	GLN
1	A	201	GLN
1	A	204	ASN
1	A	245	HIS
1	A	256	HIS
1	A	269	ASN
1	A	490	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 7 ligands modelled in this entry, 1 is monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	FAD	A	1805	-	51,58,58	2.58	19 (37%)	57,89,89	2.65	18 (31%)
5	FUM	A	1806	-	1,7,7	0.68	0	0,8,8	0.00	-
3	HEM	A	801	1	27,50,50	2.10	7 (25%)	17,82,82	1.78	3 (17%)
3	HEM	A	802	1	27,50,50	2.13	6 (22%)	17,82,82	1.46	3 (17%)
3	HEM	A	803	1	27,50,50	2.14	7 (25%)	17,82,82	2.07	5 (29%)
3	HEM	A	804	1	27,50,50	2.20	7 (25%)	17,82,82	1.37	2 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	FAD	A	1805	-	1/1/9/9	0/28/50/50	0/6/6/6
5	FUM	A	1806	-	-	0/0/5/5	0/0/0/0
3	HEM	A	801	1	-	0/6/54/54	0/0/8/8
3	HEM	A	802	1	-	0/6/54/54	0/0/8/8
3	HEM	A	803	1	-	0/6/54/54	0/0/8/8
3	HEM	A	804	1	-	0/6/54/54	0/0/8/8

All (46) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	1805	FAD	C10-N1	-4.28	1.27	1.33
3	A	801	HEM	C3B-C2B	-3.93	1.34	1.40
3	A	802	HEM	C3B-C2B	-3.70	1.35	1.40
3	A	803	HEM	C3B-C2B	-3.68	1.35	1.40
3	A	804	HEM	C3B-C2B	-3.64	1.35	1.40
3	A	803	HEM	C3C-C2C	-3.59	1.35	1.40
3	A	804	HEM	C3C-C2C	-3.53	1.35	1.40
3	A	801	HEM	C3C-C2C	-3.22	1.35	1.40
3	A	802	HEM	C3C-C2C	-3.12	1.36	1.40
4	A	1805	FAD	O4-C4	-2.57	1.18	1.24
4	A	1805	FAD	O4B-C1B	-2.22	1.38	1.41
4	A	1805	FAD	C5'-C4'	-2.05	1.48	1.51

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	801	HEM	CMA-C3A	2.04	1.55	1.51
3	A	802	HEM	CAA-C2A	2.06	1.55	1.52
4	A	1805	FAD	C4A-N3A	2.06	1.38	1.35
3	A	804	HEM	C1D-ND	2.12	1.40	1.36
3	A	803	HEM	C1D-ND	2.13	1.40	1.36
3	A	804	HEM	CMA-C3A	2.16	1.56	1.51
4	A	1805	FAD	C8-C7	2.25	1.46	1.40
4	A	1805	FAD	C4X-C10	2.28	1.44	1.41
4	A	1805	FAD	O3B-C3B	2.39	1.48	1.43
3	A	801	HEM	CAA-C2A	2.40	1.56	1.52
4	A	1805	FAD	O3'-C3'	2.60	1.49	1.43
3	A	803	HEM	CAA-C2A	2.98	1.57	1.52
4	A	1805	FAD	C2-N3	2.99	1.44	1.38
4	A	1805	FAD	C2-N1	3.04	1.44	1.38
3	A	801	HEM	C3B-CAB	3.63	1.55	1.47
3	A	804	HEM	C3B-CAB	3.64	1.55	1.47
4	A	1805	FAD	C4-N3	3.72	1.39	1.33
3	A	803	HEM	C3B-CAB	3.74	1.55	1.47
4	A	1805	FAD	O2'-C2'	3.75	1.51	1.43
3	A	802	HEM	C3B-CAB	3.79	1.55	1.47
4	A	1805	FAD	C4'-C3'	4.02	1.61	1.53
3	A	803	HEM	C3C-CAC	4.02	1.55	1.47
3	A	801	HEM	C3C-CAC	4.17	1.56	1.47
3	A	802	HEM	C3C-CAC	4.50	1.56	1.47
3	A	804	HEM	C3C-CAC	4.72	1.57	1.47
4	A	1805	FAD	C2A-N3A	4.74	1.39	1.32
3	A	804	HEM	C3D-C2D	5.07	1.52	1.37
3	A	801	HEM	C3D-C2D	5.07	1.52	1.37
4	A	1805	FAD	C6-C5X	5.12	1.49	1.41
3	A	803	HEM	C3D-C2D	5.14	1.52	1.37
3	A	802	HEM	C3D-C2D	5.32	1.53	1.37
4	A	1805	FAD	C4-C4X	6.02	1.52	1.41
4	A	1805	FAD	C9A-N10	7.11	1.47	1.38
4	A	1805	FAD	C5X-N5	7.26	1.46	1.35

All (31) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	1805	FAD	C4X-C4-N3	-6.31	114.50	123.47
4	A	1805	FAD	O3B-C3B-C2B	-6.10	92.29	111.83
4	A	1805	FAD	N3A-C2A-N1A	-4.85	124.72	128.86
4	A	1805	FAD	O4'-C4'-C3'	-4.29	98.51	109.10

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	803	HEM	CBD-CAD-C3D	-4.28	104.30	112.47
3	A	803	HEM	CMA-C3A-C4A	-3.68	122.80	128.46
3	A	801	HEM	CBD-CAD-C3D	-3.46	105.86	112.47
3	A	801	HEM	CMA-C3A-C4A	-3.38	123.26	128.46
4	A	1805	FAD	O5'-C5'-C4'	-3.36	100.39	109.36
4	A	1805	FAD	C9A-C5X-N5	-3.22	117.75	122.32
4	A	1805	FAD	C8M-C8-C9	-2.83	113.39	120.36
4	A	1805	FAD	C4-C4X-C10	-2.53	118.06	119.95
3	A	802	HEM	CMA-C3A-C4A	-2.50	124.63	128.46
3	A	804	HEM	CBA-CAA-C2A	-2.49	107.73	112.48
4	A	1805	FAD	C4A-C5A-N7A	-2.38	107.11	109.41
3	A	801	HEM	C3B-C4B-NB	-2.36	106.16	109.21
3	A	803	HEM	C3C-C4C-NC	-2.25	106.70	110.94
3	A	803	HEM	C1D-C2D-C3D	-2.21	105.45	107.00
3	A	802	HEM	CBA-CAA-C2A	-2.05	108.56	112.48
3	A	802	HEM	C4A-C3A-C2A	2.01	108.39	107.00
3	A	804	HEM	CMB-C2B-C3B	2.11	128.72	124.88
4	A	1805	FAD	C7M-C7-C8	2.34	125.60	120.72
4	A	1805	FAD	C6-C5X-C9A	2.52	122.39	119.01
4	A	1805	FAD	C9-C8-C7	3.17	125.49	119.94
3	A	803	HEM	C4A-C3A-C2A	3.26	109.27	107.00
4	A	1805	FAD	O2'-C2'-C1'	3.31	117.78	109.61
4	A	1805	FAD	C1B-N9A-C4A	3.35	132.43	126.64
4	A	1805	FAD	O4'-C4'-C5'	3.73	118.35	110.02
4	A	1805	FAD	C4X-N5-C5X	4.32	121.29	116.76
4	A	1805	FAD	O2B-C2B-C3B	5.76	130.28	111.83
4	A	1805	FAD	C4-N3-C2	9.27	123.03	115.14

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
4	A	1805	FAD	C2B

There are no torsion outliers.

There are no ring outliers.

6 monomers are involved in 29 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	1805	FAD	8	0
5	A	1806	FUM	1	0
3	A	801	HEM	7	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	802	HEM	3	0
3	A	803	HEM	3	0
3	A	804	HEM	7	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	568/571 (99%)	-0.26	5 (0%) 84 87	6, 13, 22, 29	0

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	474	SER	2.5
1	A	11	ASN	2.1
1	A	110	LYS	2.0
1	A	403	ASP	2.0
1	A	91	ASN	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	HEM	A	802	43/43	0.95	0.12	12,16,24,29	0
3	HEM	A	801	43/43	0.95	0.12	10,14,18,23	0

*Continued on next page...*



*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	HEM	A	803	43/43	0.96	0.10	9,13,25,32	0
4	FAD	A	1805	53/53	0.97	0.07	6,8,11,12	0
3	HEM	A	804	43/43	0.98	0.08	7,10,13,14	0
5	FUM	A	1806	8/8	0.99	0.07	10,11,13,13	0
2	NA	A	1810	1/1	0.99	0.05	8,8,8,8	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.