



# wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 11, 2018 – 06:07 am GMT

PDB ID : 6B8Q  
Title : Crystal Structure of the Mg<sup>2+</sup>/CaM:Kv7.5 (KCNQ5) AB domain complex  
Authors : Chang, A.; Abderemane-Ali, F.; Minor, D.L.  
Deposited on : 2017-10-09  
Resolution : 2.60 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : trunk30967  
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)  
Refmac : 5.8.0158  
CCP4 : 7.0 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk30967


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	111664	2767 (2.60-2.60)
Clashscore	122126	3110 (2.60-2.60)
Ramachandran outliers	120053	3062 (2.60-2.60)
Sidechain outliers	120020	3062 (2.60-2.60)
RSRZ outliers	108989	2706 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	75	
1	C	75	
1	E	75	
1	G	75	
2	B	149	
2	D	149	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	F	149	<div><div><div></div><div></div><div></div></div><div><div>3%</div><div>72%</div><div>24%</div><div>• •</div></div></div>
2	H	149	<div><div><div></div><div></div><div></div></div><div><div>%</div><div>85%</div><div>13%</div><div>• •</div></div></div>

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 6648 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Potassium voltage-gated channel subfamily KQT member 5.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	62	Total	C	N	O	S	0	0	0
			511	330	96	83	2			
1	C	60	Total	C	N	O	S	0	0	0
			494	318	93	81	2			
1	E	60	Total	C	N	O	S	0	0	0
			491	318	90	81	2			
1	G	66	Total	C	N	O	S	0	0	0
			550	356	103	89	2			

There are 28 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	356	GLY	-	expression tag	UNP Q9NR82
A	357	HIS	-	expression tag	UNP Q9NR82
A	358	MET	-	expression tag	UNP Q9NR82
A	359	ALA	-	expression tag	UNP Q9NR82
A	360	SER	-	expression tag	UNP Q9NR82
A	395	LYS	-	linker	UNP Q9NR82
A	396	LEU	VAL	linker	UNP Q9NR82
C	356	GLY	-	expression tag	UNP Q9NR82
C	357	HIS	-	expression tag	UNP Q9NR82
C	358	MET	-	expression tag	UNP Q9NR82
C	359	ALA	-	expression tag	UNP Q9NR82
C	360	SER	-	expression tag	UNP Q9NR82
C	395	LYS	-	linker	UNP Q9NR82
C	396	LEU	VAL	linker	UNP Q9NR82
E	356	GLY	-	expression tag	UNP Q9NR82
E	357	HIS	-	expression tag	UNP Q9NR82
E	358	MET	-	expression tag	UNP Q9NR82
E	359	ALA	-	expression tag	UNP Q9NR82
E	360	SER	-	expression tag	UNP Q9NR82
E	395	LYS	-	linker	UNP Q9NR82
E	396	LEU	VAL	linker	UNP Q9NR82

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
G	356	GLY	-	expression tag	UNP Q9NR82
G	357	HIS	-	expression tag	UNP Q9NR82
G	358	MET	-	expression tag	UNP Q9NR82
G	359	ALA	-	expression tag	UNP Q9NR82
G	360	SER	-	expression tag	UNP Q9NR82
G	395	LYS	-	linker	UNP Q9NR82
G	396	LEU	VAL	linker	UNP Q9NR82

- Molecule 2 is a protein called Calmodulin-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	146	Total	C	N	O	S	0	0	0
			1151	705	185	252	9			
2	D	143	Total	C	N	O	S	0	0	0
			1126	690	181	246	9			
2	F	144	Total	C	N	O	S	0	0	0
			1134	696	182	247	9			
2	H	146	Total	C	N	O	S	0	0	0
			1151	705	185	252	9			

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	H	4	Total	Mg	0	0
			4	4		
3	B	3	Total	Mg	0	0
			3	3		
3	D	4	Total	Mg	0	0
			4	4		
3	F	4	Total	Mg	0	0
			4	4		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	3	Total	O	0	0
			3	3		
4	B	2	Total	O	0	0
			2	2		
4	C	2	Total	O	0	0
			2	2		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	D	3	Total 3	O 3	0	0
4	E	1	Total 1	O 1	0	0
4	F	2	Total 2	O 2	0	0
4	G	4	Total 4	O 4	0	0
4	H	8	Total 8	O 8	0	0

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Potassium voltage-gated channel subfamily KQT member 5




- Molecule 1: Potassium voltage-gated channel subfamily KQT member 5




- Molecule 1: Potassium voltage-gated channel subfamily KQT member 5




- Molecule 1: Potassium voltage-gated channel subfamily KQT member 5




- Molecule 2: Calmodulin-1




- Molecule 2: Calmodulin-1







● Molecule 2: Calmodulin-1



● Molecule 2: Calmodulin-1





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	70.82Å 116.84Å 119.81Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	45.74 – 2.60 45.74 – 2.60	Depositor EDS
% Data completeness (in resolution range)	100.0 (45.74-2.60) 100.0 (45.74-2.60)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.39 (at 2.61Å)	Xtriage
Refinement program	PHENIX 1.11.1_2575	Depositor
R, $R_{free}$	0.222 , 0.270 0.222 , 0.269	Depositor DCC
$R_{free}$ test set	1556 reflections (4.97%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	60.5	Xtriage
Anisotropy	0.132	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 48.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.018 for -h,l,k	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	6648	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	69.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.30% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.26	0/521	0.41	0/700
1	C	0.23	0/504	0.37	0/678
1	E	0.25	0/501	0.36	0/675
1	G	0.25	0/562	0.37	0/754
2	B	0.35	0/1163	0.44	0/1562
2	D	0.26	0/1138	0.45	0/1528
2	F	0.27	0/1146	0.48	0/1539
2	H	0.26	0/1163	0.42	0/1562
All	All	0.28	0/6698	0.43	0/8998

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	511	0	553	14	0
1	C	494	0	529	9	0
1	E	491	0	527	10	0
1	G	550	0	588	9	0
2	B	1151	0	1075	13	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	D	1126	0	1052	18	0
2	F	1134	0	1063	36	0
2	H	1151	0	1075	14	0
3	B	3	0	0	0	0
3	D	4	0	0	0	0
3	F	4	0	0	0	0
3	H	4	0	0	0	0
4	A	3	0	0	0	0
4	B	2	0	0	0	0
4	C	2	0	0	0	0
4	D	3	0	0	0	0
4	E	1	0	0	0	0
4	F	2	0	0	1	0
4	G	4	0	0	0	0
4	H	8	0	0	0	0
All	All	6648	0	6462	105	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 105 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:F:86:ARG:NH2	2:F:139:GLU:HG2	1.78	0.97
2:F:86:ARG:CZ	2:F:139:GLU:HG2	2.00	0.90
1:E:392:THR:HG23	1:E:526:ARG:HH12	1.40	0.84
1:E:535:ARG:HG3	1:E:538:LYS:HE3	1.63	0.81
2:D:14:GLU:OE1	2:D:14:GLU:N	2.14	0.80

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	60/75 (80%)	58 (97%)	2 (3%)	0	100	100
1	C	58/75 (77%)	55 (95%)	3 (5%)	0	100	100
1	E	58/75 (77%)	51 (88%)	7 (12%)	0	100	100
1	G	64/75 (85%)	60 (94%)	4 (6%)	0	100	100
2	B	144/149 (97%)	141 (98%)	3 (2%)	0	100	100
2	D	141/149 (95%)	133 (94%)	8 (6%)	0	100	100
2	F	142/149 (95%)	137 (96%)	5 (4%)	0	100	100
2	H	144/149 (97%)	140 (97%)	4 (3%)	0	100	100
All	All	811/896 (90%)	775 (96%)	36 (4%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	55/66 (83%)	55 (100%)	0	100	100
1	C	53/66 (80%)	52 (98%)	1 (2%)	60	82
1	E	53/66 (80%)	52 (98%)	1 (2%)	60	82
1	G	59/66 (89%)	56 (95%)	3 (5%)	26	50
2	B	125/127 (98%)	124 (99%)	1 (1%)	83	93
2	D	122/127 (96%)	119 (98%)	3 (2%)	50	76
2	F	123/127 (97%)	119 (97%)	4 (3%)	41	68
2	H	125/127 (98%)	123 (98%)	2 (2%)	65	85
All	All	715/772 (93%)	700 (98%)	15 (2%)	56	80

5 of 15 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	F	86	ARG
2	F	95	ASP

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	G	535	ARG
1	E	395	LYS
1	G	396	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

Of 15 ligands modelled in this entry, 15 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2	OWAB(Å <sup>2</sup> )	Q < 0.9
1	A	62/75 (82%)	0.24	2 (3%) 47 40	41, 59, 94, 112	0
1	C	60/75 (80%)	-0.04	1 (1%) 70 65	48, 63, 92, 96	0
1	E	60/75 (80%)	0.26	3 (5%) 29 22	53, 68, 106, 113	0
1	G	66/75 (88%)	0.09	2 (3%) 50 43	37, 55, 92, 105	0
2	B	146/149 (97%)	-0.07	0 100 100	43, 61, 78, 92	0
2	D	143/149 (95%)	0.06	3 (2%) 63 58	46, 72, 92, 122	0
2	F	144/149 (96%)	0.39	5 (3%) 44 36	49, 86, 112, 122	0
2	H	146/149 (97%)	0.02	1 (0%) 87 85	41, 61, 87, 100	0
All	All	827/896 (92%)	0.11	17 (2%) 63 58	37, 67, 102, 122	0

The worst 5 of 17 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	396	LEU	3.4
2	F	83	GLU	2.8
1	E	535	ARG	2.6
1	E	537	PHE	2.6
1	A	391	ALA	2.4

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	MG	H	203	1/1	0.73	0.23	76,76,76,76	0
3	MG	B	203	1/1	0.90	0.18	56,56,56,56	0
3	MG	H	204	1/1	0.91	0.14	61,61,61,61	0
3	MG	D	203	1/1	0.93	0.14	75,75,75,75	0
3	MG	F	202	1/1	0.93	0.16	56,56,56,56	0
3	MG	F	203	1/1	0.93	0.18	97,97,97,97	0
3	MG	F	204	1/1	0.94	0.17	55,55,55,55	0
3	MG	D	202	1/1	0.95	0.17	43,43,43,43	0
3	MG	D	201	1/1	0.96	0.13	44,44,44,44	0
3	MG	B	201	1/1	0.97	0.11	48,48,48,48	0
3	MG	B	202	1/1	0.97	0.27	41,41,41,41	0
3	MG	F	201	1/1	0.97	0.09	44,44,44,44	0
3	MG	H	202	1/1	0.98	0.29	31,31,31,31	0
3	MG	H	201	1/1	0.98	0.19	42,42,42,42	0
3	MG	D	204	1/1	0.99	0.17	41,41,41,41	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.