



# wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 8, 2018 – 07:42 pm GMT

PDB ID : 6B9T  
Title : Crystal structure of MPnS with substrate 2-hydroxyethylphosphonate (2-HEP) and Fe(II) bound  
Authors : Born, D.A.; Drennan, C.L.  
Deposited on : 2017-10-11  
Resolution : 2.35 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.3 (157068), CSD as539be (2018)  
Xtriage (Phenix) : 1.13  
EDS : trunk30967  
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)  
Refmac : 5.8.0158  
CCP4 : 7.0 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk30967


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	111664	1015 (2.36-2.36)
Clashscore	122126	1081 (2.36-2.36)
Ramachandran outliers	120053	1066 (2.36-2.36)
Sidechain outliers	120020	1067 (2.36-2.36)
RSRZ outliers	108989	1002 (2.36-2.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	457	<div> <div></div> <div>90%</div> <div>8%</div> <div></div> </div>
1	B	457	<div> <div></div> <div>88%</div> <div>9%</div> <div></div> </div>
1	C	457	<div> <div></div> <div>89%</div> <div>9%</div> <div></div> </div>
1	D	457	<div> <div></div> <div>91%</div> <div>8%</div> <div></div> </div>
1	E	457	<div> <div></div> <div>89%</div> <div>8%</div> <div></div> </div>
1	F	457	<div> <div></div> <div>92%</div> <div>6%</div> <div></div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	G	457	<div><div></div><div>3%</div><div>87%</div><div>10%</div><div></div></div>
1	H	457	<div><div></div><div>5%</div><div>86%</div><div>10%</div><div></div></div>

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 29377 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Methylphosphonate synthase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	447	Total	C	N	O	S	0	0	0
			3551	2250	591	697	13			
1	B	446	Total	C	N	O	S	0	0	0
			3533	2240	589	691	13			
1	C	447	Total	C	N	O	S	0	0	0
			3552	2252	595	692	13			
1	D	453	Total	C	N	O	S	0	0	0
			3614	2294	604	703	13			
1	E	444	Total	C	N	O	S	0	0	0
			3527	2238	589	687	13			
1	F	447	Total	C	N	O	S	0	0	0
			3532	2234	590	695	13			
1	G	446	Total	C	N	O	S	0	0	0
			3509	2222	584	690	13			
1	H	442	Total	C	N	O	S	0	0	0
			3489	2212	580	684	13			

- Molecule 2 is FE (III) ION (three-letter code: FE) (formula: Fe).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	G	1	Total	Fe	0	0
			1	1		
2	D	1	Total	Fe	0	0
			1	1		
2	E	1	Total	Fe	0	0
			1	1		
2	H	1	Total	Fe	0	0
			1	1		
2	B	2	Total	Fe	0	0
			2	2		
2	C	1	Total	Fe	0	0
			1	1		

*Continued on next page...*

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Fe	0	0
			1	1		
2	F	1	Total	Fe	0	0
			1	1		

- Molecule 3 is (2-hydroxyethyl)phosphonic acid (three-letter code: 2HE) (formula:  $C_2H_7O_4P$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	O	P	0	0
			7	2	4	1		
3	B	1	Total	C	O	P	0	0
			7	2	4	1		
3	C	1	Total	C	O	P	0	0
			7	2	4	1		
3	D	1	Total	C	O	P	0	0
			7	2	4	1		
3	E	1	Total	C	O	P	0	0
			7	2	4	1		
3	F	1	Total	C	O	P	0	0
			7	2	4	1		
3	G	1	Total	C	O	P	0	0
			7	2	4	1		
3	H	1	Total	C	O	P	0	0
			7	2	4	1		

- Molecule 4 is FORMIC ACID (three-letter code: FMT) (formula:  $CH_2O_2$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	O	0	0
			3	1	2		
4	C	1	Total	C	O	0	0
			3	1	2		
4	D	1	Total	C	O	0	0
			3	1	2		
4	E	1	Total	C	O	0	0
			3	1	2		
4	F	1	Total	C	O	0	0
			3	1	2		
4	G	1	Total	C	O	0	0
			3	1	2		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	122	Total	O	0	0
			122	122		
5	B	125	Total	O	0	0
			125	125		
5	C	167	Total	O	0	0
			167	167		
5	D	162	Total	O	0	0
			162	162		
5	E	143	Total	O	0	0
			143	143		
5	F	112	Total	O	0	0
			112	112		

*Continued on next page...*


*Continued from previous page...*

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	G	48	Total 48	O 48	0	0
5	H	108	Total 108	O 108	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Methylphosphonate synthase




- Molecule 1: Methylphosphonate synthase



- Molecule 1: Methylphosphonate synthase



- Molecule 1: Methylphosphonate synthase









• Molecule 1: Methylphosphonate synthase

Chain E: 89% 8%




• Molecule 1: Methylphosphonate synthase

Chain F: 92% 6%




• Molecule 1: Methylphosphonate synthase

Chain G: 3% 87% 10%



• Molecule 1: Methylphosphonate synthase

Chain H: 5% 86% 10%



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	70.85Å 351.25Å 76.87Å 90.00° 103.36° 90.00°	Depositor
Resolution (Å)	46.10 – 2.35 74.79 – 2.35	Depositor EDS
% Data completeness (in resolution range)	89.9 (46.10-2.35) 89.9 (74.79-2.35)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.07 (at 2.34Å)	Xtriage
Refinement program	PHENIX (1.10_2155: ???)	Depositor
R, $R_{free}$	0.190 , 0.238 0.191 , 0.238	Depositor DCC
$R_{free}$ test set	6643 reflections (4.90%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	41.1	Xtriage
Anisotropy	0.473	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.26 , 25.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.46$ , $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	29377	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	46.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.02% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FMT, 2HE, FE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.27	0/3634	0.47	0/4926
1	B	0.27	0/3616	0.48	0/4904
1	C	0.28	0/3635	0.48	0/4924
1	D	0.28	0/3699	0.48	0/5008
1	E	0.27	0/3610	0.48	0/4895
1	F	0.27	0/3615	0.46	0/4906
1	G	0.26	0/3592	0.45	0/4878
1	H	0.27	0/3571	0.46	0/4846
All	All	0.27	0/28972	0.47	0/39287

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3551	0	3397	22	0
1	B	3533	0	3373	32	0
1	C	3552	0	3415	29	0
1	D	3614	0	3495	30	0
1	E	3527	0	3378	27	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	F	3532	0	3351	19	0
1	G	3509	0	3312	33	0
1	H	3489	0	3314	32	0
2	A	1	0	0	0	0
2	B	2	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
2	E	1	0	0	0	0
2	F	1	0	0	0	0
2	G	1	0	0	0	0
2	H	1	0	0	0	0
3	A	7	0	4	0	0
3	B	7	0	4	0	0
3	C	7	0	4	1	0
3	D	7	0	4	0	0
3	E	7	0	4	0	0
3	F	7	0	4	0	0
3	G	7	0	4	0	0
3	H	7	0	4	1	0
4	B	3	0	1	0	0
4	C	3	0	1	1	0
4	D	3	0	1	1	0
4	E	3	0	1	0	0
4	F	3	0	1	0	0
4	G	3	0	1	0	0
5	A	122	0	0	0	0
5	B	125	0	0	5	0
5	C	167	0	0	4	0
5	D	162	0	0	1	0
5	E	143	0	0	4	0
5	F	112	0	0	1	0
5	G	48	0	0	0	0
5	H	108	0	0	3	0
All	All	29377	0	27073	190	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 190 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:G:224:LEU:HD21	1:H:402:PRO:HD2	1.63	0.80
1:B:367:GLN:HG2	1:B:418:ARG:HG2	1.67	0.77
1:H:367:GLN:HG2	1:H:418:ARG:HG2	1.67	0.73
1:E:172:HIS:ND1	5:E:601:HOH:O	2.22	0.73
1:C:224:LEU:HD21	1:D:402:PRO:HD2	1.73	0.71

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	445/457 (97%)	439 (99%)	6 (1%)	0	100	100
1	B	444/457 (97%)	434 (98%)	8 (2%)	2 (0%)	31	34
1	C	445/457 (97%)	436 (98%)	6 (1%)	3 (1%)	24	25
1	D	451/457 (99%)	443 (98%)	8 (2%)	0	100	100
1	E	442/457 (97%)	432 (98%)	8 (2%)	2 (0%)	31	34
1	F	445/457 (97%)	436 (98%)	9 (2%)	0	100	100
1	G	444/457 (97%)	433 (98%)	9 (2%)	2 (0%)	31	34
1	H	440/457 (96%)	431 (98%)	7 (2%)	2 (0%)	31	34
All	All	3556/3656 (97%)	3484 (98%)	61 (2%)	11 (0%)	43	50

5 of 11 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	197	GLY
1	H	197	GLY
1	C	197	GLY
1	C	354	SER
1	E	197	GLY

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	389/409 (95%)	384 (99%)	5 (1%)	71	81
1	B	385/409 (94%)	382 (99%)	3 (1%)	83	90
1	C	389/409 (95%)	386 (99%)	3 (1%)	83	90
1	D	399/409 (98%)	395 (99%)	4 (1%)	78	86
1	E	385/409 (94%)	382 (99%)	3 (1%)	83	90
1	F	385/409 (94%)	383 (100%)	2 (0%)	90	94
1	G	379/409 (93%)	376 (99%)	3 (1%)	83	90
1	H	379/409 (93%)	376 (99%)	3 (1%)	83	90
All	All	3090/3272 (94%)	3064 (99%)	26 (1%)	83	90

5 of 26 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	115	MET
1	D	351	VAL
1	H	182	MET
1	D	182	MET
1	D	255	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	380	ASN
1	B	389	GLN
1	E	218	GLN
1	G	239	HIS
1	G	329	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 23 ligands modelled in this entry, 9 are monoatomic - leaving 14 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	2HE	A	502	2	6,6,6	2.86	4 (66%)	7,8,8	1.23	1 (14%)
3	2HE	B	503	2	6,6,6	1.97	3 (50%)	7,8,8	1.74	1 (14%)
4	FMT	B	504	-	0,2,2	0.00	-	0,1,1	0.00	-
3	2HE	C	502	2	6,6,6	2.86	4 (66%)	7,8,8	1.20	1 (14%)
4	FMT	C	503	-	0,2,2	0.00	-	0,1,1	0.00	-
3	2HE	D	502	2	6,6,6	2.84	4 (66%)	7,8,8	1.42	0
4	FMT	D	503	-	0,2,2	0.00	-	0,1,1	0.00	-
3	2HE	E	502	2	6,6,6	1.99	3 (50%)	7,8,8	1.42	1 (14%)
4	FMT	E	503	-	0,2,2	0.00	-	0,1,1	0.00	-
3	2HE	F	502	2	6,6,6	2.05	3 (50%)	7,8,8	1.40	1 (14%)
4	FMT	F	503	-	0,2,2	0.00	-	0,1,1	0.00	-
3	2HE	G	502	2	6,6,6	2.95	4 (66%)	7,8,8	1.59	1 (14%)
4	FMT	G	503	-	0,2,2	0.00	-	0,1,1	0.00	-
3	2HE	H	502	2	6,6,6	2.88	4 (66%)	7,8,8	1.35	1 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	2HE	A	502	2	-	0/4/4/4	0/0/0/0
3	2HE	B	503	2	-	0/4/4/4	0/0/0/0
4	FMT	B	504	-	-	0/0/0/0	0/0/0/0
3	2HE	C	502	2	-	0/4/4/4	0/0/0/0
4	FMT	C	503	-	-	0/0/0/0	0/0/0/0
3	2HE	D	502	2	-	0/4/4/4	0/0/0/0
4	FMT	D	503	-	-	0/0/0/0	0/0/0/0
3	2HE	E	502	2	-	0/4/4/4	0/0/0/0
4	FMT	E	503	-	-	0/0/0/0	0/0/0/0
3	2HE	F	502	2	-	0/4/4/4	0/0/0/0
4	FMT	F	503	-	-	0/0/0/0	0/0/0/0
3	2HE	G	502	2	-	0/4/4/4	0/0/0/0
4	FMT	G	503	-	-	0/0/0/0	0/0/0/0
3	2HE	H	502	2	-	0/4/4/4	0/0/0/0

The worst 5 of 29 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	H	502	2HE	P-O2	-2.79	1.48	1.54
3	A	502	2HE	P-O3	-2.64	1.48	1.54
3	G	502	2HE	P-O2	-2.60	1.48	1.54
3	D	502	2HE	P-O3	-2.58	1.48	1.54
3	C	502	2HE	P-O2	-2.54	1.48	1.54

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	503	2HE	P-CA-CB	-4.23	109.74	114.89
3	G	502	2HE	P-CA-CB	-3.56	110.56	114.89
3	E	502	2HE	P-CA-CB	-3.03	111.20	114.89
3	H	502	2HE	P-CA-CB	-2.88	111.38	114.89
3	F	502	2HE	P-CA-CB	-2.68	111.62	114.89

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	502	2HE	1	0
4	C	503	FMT	1	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	D	503	FMT	1	0
3	H	502	2HE	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	447/457 (97%)	-0.27	4 (0%) 84 90	33, 45, 70, 83	0
1	B	446/457 (97%)	-0.28	1 (0%) 94 97	33, 43, 58, 69	0
1	C	447/457 (97%)	-0.35	0 100 100	30, 40, 52, 63	0
1	D	453/457 (99%)	-0.34	0 100 100	28, 39, 55, 64	0
1	E	444/457 (97%)	-0.31	1 (0%) 94 97	31, 41, 59, 67	0
1	F	447/457 (97%)	-0.27	0 100 100	33, 44, 59, 72	0
1	G	446/457 (97%)	0.10	14 (3%) 49 61	34, 58, 96, 104	0
1	H	442/457 (96%)	0.04	21 (4%) 30 44	34, 49, 105, 113	0
All	All	3572/3656 (97%)	-0.21	41 (1%) 80 88	28, 44, 74, 113	0

The worst 5 of 41 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	H	39	GLY	6.2
1	G	39	GLY	5.7
1	H	55	ILE	4.5
1	H	45	ILE	4.3
1	H	34	ALA	3.9

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	2HE	G	502	7/7	0.92	0.14	56,62,68,69	0
4	FMT	G	503	3/3	0.96	0.13	38,38,41,43	0
4	FMT	F	503	3/3	0.97	0.11	39,39,40,42	0
3	2HE	F	502	7/7	0.97	0.12	39,42,44,45	0
3	2HE	H	502	7/7	0.98	0.13	46,50,53,54	0
3	2HE	D	502	7/7	0.98	0.12	31,33,36,37	0
4	FMT	E	503	3/3	0.98	0.09	34,34,34,37	0
3	2HE	A	502	7/7	0.98	0.12	38,39,42,45	0
4	FMT	C	503	3/3	0.98	0.11	29,29,34,36	0
4	FMT	B	504	3/3	0.98	0.13	35,35,36,40	0
4	FMT	D	503	3/3	0.98	0.15	35,35,37,41	0
3	2HE	E	502	7/7	0.98	0.12	33,34,39,39	0
3	2HE	C	502	7/7	0.98	0.14	36,38,42,43	0
3	2HE	B	503	7/7	0.98	0.11	38,39,43,44	0
2	FE	G	501	1/1	0.98	0.09	56,56,56,56	0
2	FE	F	501	1/1	0.99	0.07	42,42,42,42	0
2	FE	D	501	1/1	0.99	0.11	30,30,30,30	0
2	FE	B	502	1/1	0.99	0.07	60,60,60,60	0
2	FE	C	501	1/1	0.99	0.14	38,38,38,38	0
2	FE	E	501	1/1	0.99	0.08	34,34,34,34	0
2	FE	H	501	1/1	0.99	0.13	48,48,48,48	0
2	FE	A	501	1/1	1.00	0.07	43,43,43,43	0
2	FE	B	501	1/1	1.00	0.10	40,40,40,40	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.