



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 9, 2018 – 07:22 am GMT

PDB ID : 2C12
Title : Crystal Structure of Nitroalkane Oxidase in Complex with Spermine, a Competitive Inhibitor
Authors : Nagpal, A.; Valley, M.P.; Fitzpatrick, P.F.; Orville, A.M.
Deposited on : 2005-09-10
Resolution : 2.07 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.3 (157068), CSD as539be (2018)
Xtriage (Phenix) : 1.13
EDS : trunk30967
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)
Refmac : 5.8.0158
CCP4 : 7.0 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk30967


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.07 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	111664	2250 (2.08-2.04)
Clashscore	122126	2358 (2.08-2.04)
Ramachandran outliers	120053	2339 (2.08-2.04)
Sidechain outliers	120020	2339 (2.08-2.04)
RSRZ outliers	108989	2211 (2.08-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	439	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 1%, orange 1%, yellow 13%, green 82%, grey 4%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> % 82% 13% • • </div> </div>
1	B	439	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, green 79%, yellow 16%, orange 5%, grey 0%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> 79% 16% • • </div> </div>
1	C	439	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 3%, orange 1%, yellow 14%, green 81%, grey 2%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> 3% 81% 14% • • </div> </div>
1	D	439	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 1%, orange 1%, yellow 14%, green 82%, grey 2%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> % 82% 14% • • </div> </div>
1	E	439	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 2%, orange 1%, yellow 11%, green 85%, grey 2%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> 2% 85% 11% • • </div> </div>
1	F	439	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 3%, orange 1%, yellow 13%, green 83%, grey 2%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> 3% 83% 13% • • </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	SPM	C	1434	-	-	X	-
5	PE4	B	1436	-	-	X	-
5	PE4	F	1435	-	-	X	-

2 Entry composition [i](#)


There are 6 unique types of molecules in this entry. The entry contains 21487 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called NITROALKANE OXIDASE.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	430	Total	C	N	O	S	0	1	0
			3309	2096	567	625	21			
1	B	430	Total	C	N	O	S	0	1	0
			3309	2096	567	625	21			
1	C	430	Total	C	N	O	S	0	1	0
			3309	2096	567	625	21			
1	D	430	Total	C	N	O	S	0	3	0
			3313	2097	568	627	21			
1	E	430	Total	C	N	O	S	0	1	0
			3309	2096	567	625	21			
1	F	430	Total	C	N	O	S	0	1	0
			3309	2096	567	625	21			

- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula: C₂₇H₃₃N₉O₁₅P₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	B	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	C	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	D	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	E	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	F	1	Total	C	N	O	P	0	0
			53	27	9	15	2		

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	C	1	Total	C	O	0	0
			6	3	3		
3	C	1	Total	C	O	0	0
			6	3	3		
3	C	1	Total	C	O	0	0
			6	3	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	C	1	Total	C	O	0	0
			6	3	3		
3	D	1	Total	C	O	0	0
			6	3	3		
3	D	1	Total	C	O	0	0
			6	3	3		
3	D	1	Total	C	O	0	0
			6	3	3		
3	D	1	Total	C	O	0	0
			6	3	3		
3	E	1	Total	C	O	0	0
			6	3	3		
3	E	1	Total	C	O	0	0
			6	3	3		
3	E	1	Total	C	O	0	0
			6	3	3		
3	F	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is SPERMINE (three-letter code: SPM) (formula: $C_{10}H_{26}N_4$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	N	0	0
			14	10	4		
4	C	1	Total	C	N	0	0
			14	10	4		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	D	1	Total	C	N	0	0
			14	10	4		
4	F	1	Total	C	N	0	0
			14	10	4		

- Molecule 5 is 2-{2-[2-(2-{2-[2-(2-ETHOXY-ETHOXY)-ETHOXY]-ETHOXY}-ETHOXY)-ETHOXY]-ETHOXY}-ETHANOL (three-letter code: PE4) (formula: C₁₆H₃₄O₈).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	B	1	Total	C	O	0	0
			24	16	8		
5	F	1	Total	C	O	0	0
			24	16	8		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	195	Total	O	0	0
			195	195		
6	B	207	Total	O	0	0
			207	207		
6	C	164	Total	O	0	0
			164	164		
6	D	159	Total	O	0	0
			159	159		
6	E	187	Total	O	0	0
			187	187		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	F	205	Total 205	O 205	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: NITROALKANE OXIDASE



• Molecule 1: NITROALKANE OXIDASE






• Molecule 1: NITROALKANE OXIDASE











● Molecule 1: NITROALKANE OXIDASE



● Molecule 1: NITROALKANE OXIDASE



● Molecule 1: NITROALKANE OXIDASE



4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	103.39Å 103.39Å 485.13Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	50.00 – 2.07 49.24 – 2.07	Depositor EDS
% Data completeness (in resolution range)	97.8 (50.00-2.07) 97.8 (49.24-2.07)	Depositor EDS
R_{merge}	0.14	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	7.71 (at 2.07Å)	Xtriage
Refinement program	REFMAC 5.1.24	Depositor
R, R_{free}	0.188 , 0.225 0.199 , 0.236	Depositor DCC
R_{free} test set	9048 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	25.1	Xtriage
Anisotropy	0.303	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 40.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.021 for -h,-k,l	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	21487	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.13% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PE4, GOL, SPM, FAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.55	1/3384 (0.0%)	0.79	13/4592 (0.3%)
1	B	0.53	0/3384	0.80	11/4592 (0.2%)
1	C	0.52	0/3384	0.77	9/4592 (0.2%)
1	D	0.52	0/3398	0.78	15/4611 (0.3%)
1	E	0.53	0/3384	0.75	7/4592 (0.2%)
1	F	0.54	1/3384 (0.0%)	0.80	11/4592 (0.2%)
All	All	0.53	2/20318 (0.0%)	0.78	66/27571 (0.2%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	379	MET	SD-CE	-9.42	1.25	1.77
1	F	95	VAL	CB-CG1	-5.23	1.41	1.52

The worst 5 of 66 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	F	331	ARG	NE-CZ-NH2	-10.63	114.99	120.30
1	D	410	ARG	NE-CZ-NH1	9.35	124.97	120.30
1	F	331	ARG	NE-CZ-NH1	9.31	124.95	120.30
1	D	410	ARG	NE-CZ-NH2	-9.11	115.75	120.30
1	A	331	ARG	NE-CZ-NH2	-8.07	116.27	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3309	0	3320	45	0
1	B	3309	0	3320	58	0
1	C	3309	0	3320	45	0
1	D	3313	0	3323	34	0
1	E	3309	0	3320	28	0
1	F	3309	0	3320	36	0
2	A	53	0	31	6	0
2	B	53	0	31	8	0
2	C	53	0	31	4	0
2	D	53	0	31	3	0
2	E	53	0	31	3	0
2	F	53	0	31	5	0
3	A	6	0	8	0	0
3	B	12	0	16	0	0
3	C	24	0	32	2	0
3	D	24	0	32	2	0
3	E	18	0	24	1	0
3	F	6	0	8	3	0
4	B	14	0	26	4	0
4	C	14	0	26	9	0
4	D	14	0	26	0	0
4	F	14	0	26	5	0
5	B	24	0	34	17	0
5	F	24	0	34	9	0
6	A	195	0	0	4	0
6	B	207	0	0	14	0
6	C	164	0	0	7	0
6	D	159	0	0	5	0
6	E	187	0	0	5	0
6	F	205	0	0	14	0
All	All	21487	0	20401	261	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 261 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:B:1436:PE4:H141	6:B:2206:HOH:O	1.29	1.27
6:A:2191:HOH:O	1:B:310:ILE:HG21	1.35	1.23
1:C:141:ASN:HD21	2:C:1433:FAD:H61A	1.19	0.91
1:A:379:MET:HE2	1:B:401:PHE:HA	1.56	0.88
1:A:10:GLN:HG3	1:A:75:HIS:CE1	2.09	0.87

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	429/439 (98%)	423 (99%)	4 (1%)	2 (0%)	31	20
1	B	429/439 (98%)	421 (98%)	5 (1%)	3 (1%)	24	13
1	C	429/439 (98%)	423 (99%)	4 (1%)	2 (0%)	31	20
1	D	431/439 (98%)	425 (99%)	5 (1%)	1 (0%)	49	41
1	E	429/439 (98%)	423 (99%)	4 (1%)	2 (0%)	31	20
1	F	429/439 (98%)	422 (98%)	6 (1%)	1 (0%)	49	41
All	All	2576/2634 (98%)	2537 (98%)	28 (1%)	11 (0%)	36	26

5 of 11 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	169	TRP
1	B	169	TRP
1	B	430	GLY
1	C	169	TRP
1	D	169	TRP

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	356/364 (98%)	329 (92%)	27 (8%)	14	6
1	B	356/364 (98%)	328 (92%)	28 (8%)	13	6
1	C	356/364 (98%)	331 (93%)	25 (7%)	16	8
1	D	358/364 (98%)	335 (94%)	23 (6%)	19	10
1	E	356/364 (98%)	332 (93%)	24 (7%)	18	9
1	F	356/364 (98%)	333 (94%)	23 (6%)	19	10
All	All	2138/2184 (98%)	1988 (93%)	150 (7%)	17	8

5 of 150 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	164	SER
1	D	45	THR
1	F	192	ASP
1	C	192	ASP
1	C	365	VAL

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 80 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	198	GLN
1	D	43	GLN
1	F	113	GLN
1	C	256	ASN
1	C	314	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

27 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	FAD	A	1432	-	51,58,58	1.62	10 (19%)	57,89,89	2.18	12 (21%)
3	GOL	A	1433	-	5,5,5	0.41	0	5,5,5	0.30	0
2	FAD	B	1432	-	51,58,58	1.81	10 (19%)	57,89,89	2.05	12 (21%)
4	SPM	B	1433	-	13,13,13	0.51	0	12,12,12	1.17	1 (8%)
3	GOL	B	1434	-	5,5,5	0.38	0	5,5,5	0.31	0
3	GOL	B	1435	-	5,5,5	0.43	0	5,5,5	0.48	0
5	PE4	B	1436	-	23,23,23	0.62	0	22,22,22	0.98	2 (9%)
3	GOL	C	1432	-	5,5,5	0.39	0	5,5,5	0.64	0
2	FAD	C	1433	-	51,58,58	1.78	10 (19%)	57,89,89	2.13	16 (28%)
4	SPM	C	1434	-	13,13,13	0.32	0	12,12,12	1.18	1 (8%)
3	GOL	C	1435	-	5,5,5	0.36	0	5,5,5	0.34	0
3	GOL	C	1436	-	5,5,5	0.47	0	5,5,5	0.77	0
3	GOL	C	1437	-	5,5,5	0.58	0	5,5,5	1.19	1 (20%)
3	GOL	D	1432	-	5,5,5	0.42	0	5,5,5	0.54	0
2	FAD	D	1433	-	51,58,58	1.70	9 (17%)	57,89,89	2.19	13 (22%)
4	SPM	D	1434	-	13,13,13	0.34	0	12,12,12	0.74	0
3	GOL	D	1435	-	5,5,5	0.57	0	5,5,5	0.35	0
3	GOL	D	1436	-	5,5,5	0.44	0	5,5,5	0.33	0
3	GOL	D	1437	-	5,5,5	0.45	0	5,5,5	0.50	0
2	FAD	E	1432	-	51,58,58	1.97	9 (17%)	57,89,89	2.26	12 (21%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	GOL	E	1433	-	5,5,5	0.31	0	5,5,5	0.43	0
3	GOL	E	1434	-	5,5,5	0.46	0	5,5,5	0.89	0
3	GOL	E	1435	-	5,5,5	0.46	0	5,5,5	0.16	0
2	FAD	F	1432	-	51,58,58	1.66	9 (17%)	57,89,89	2.29	11 (19%)
4	SPM	F	1433	-	13,13,13	0.40	0	12,12,12	0.87	0
3	GOL	F	1434	-	5,5,5	0.48	0	5,5,5	0.42	0
5	PE4	F	1435	-	23,23,23	0.83	0	22,22,22	1.07	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FAD	A	1432	-	-	0/28/50/50	0/6/6/6
3	GOL	A	1433	-	-	0/4/4/4	0/0/0/0
2	FAD	B	1432	-	-	0/28/50/50	0/6/6/6
4	SPM	B	1433	-	-	0/11/11/11	0/0/0/0
3	GOL	B	1434	-	-	0/4/4/4	0/0/0/0
3	GOL	B	1435	-	-	0/4/4/4	0/0/0/0
5	PE4	B	1436	-	-	0/21/21/21	0/0/0/0
3	GOL	C	1432	-	-	0/4/4/4	0/0/0/0
2	FAD	C	1433	-	-	0/28/50/50	0/6/6/6
4	SPM	C	1434	-	-	0/11/11/11	0/0/0/0
3	GOL	C	1435	-	-	0/4/4/4	0/0/0/0
3	GOL	C	1436	-	-	0/4/4/4	0/0/0/0
3	GOL	C	1437	-	-	0/4/4/4	0/0/0/0
3	GOL	D	1432	-	-	0/4/4/4	0/0/0/0
2	FAD	D	1433	-	-	0/28/50/50	0/6/6/6
4	SPM	D	1434	-	-	0/11/11/11	0/0/0/0
3	GOL	D	1435	-	-	0/4/4/4	0/0/0/0
3	GOL	D	1436	-	-	0/4/4/4	0/0/0/0
3	GOL	D	1437	-	-	0/4/4/4	0/0/0/0
2	FAD	E	1432	-	-	0/28/50/50	0/6/6/6
3	GOL	E	1433	-	-	0/4/4/4	0/0/0/0
3	GOL	E	1434	-	-	0/4/4/4	0/0/0/0
3	GOL	E	1435	-	-	0/4/4/4	0/0/0/0
2	FAD	F	1432	-	-	0/28/50/50	0/6/6/6
4	SPM	F	1433	-	-	0/11/11/11	0/0/0/0
3	GOL	F	1434	-	-	0/4/4/4	0/0/0/0
5	PE4	F	1435	-	-	0/21/21/21	0/0/0/0

The worst 5 of 57 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	E	1432	FAD	C4'-C3'	-5.74	1.42	1.53
2	B	1432	FAD	C4'-C3'	-5.35	1.43	1.53
2	C	1433	FAD	C4'-C3'	-5.32	1.43	1.53
2	A	1432	FAD	C4'-C3'	-5.23	1.43	1.53
2	D	1433	FAD	C4'-C3'	-5.18	1.43	1.53

The worst 5 of 81 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	F	1432	FAD	N3A-C2A-N1A	-8.30	121.76	128.86
2	D	1433	FAD	N3A-C2A-N1A	-8.09	121.94	128.86
2	A	1432	FAD	N3A-C2A-N1A	-7.74	122.24	128.86
2	B	1432	FAD	N3A-C2A-N1A	-7.15	122.74	128.86
2	C	1433	FAD	N3A-C2A-N1A	-7.13	122.76	128.86

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

17 monomers are involved in 80 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1432	FAD	6	0
2	B	1432	FAD	8	0
4	B	1433	SPM	4	0
5	B	1436	PE4	17	0
2	C	1433	FAD	4	0
4	C	1434	SPM	9	0
3	C	1436	GOL	1	0
3	C	1437	GOL	1	0
2	D	1433	FAD	3	0
3	D	1436	GOL	1	0
3	D	1437	GOL	1	0
2	E	1432	FAD	3	0
3	E	1435	GOL	1	0
2	F	1432	FAD	5	0
4	F	1433	SPM	5	0
3	F	1434	GOL	3	0
5	F	1435	PE4	9	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	430/439 (97%)	-0.19	5 (1%) 79 80	17, 25, 36, 45	0
1	B	430/439 (97%)	-0.24	2 (0%) 90 92	16, 24, 36, 49	0
1	C	430/439 (97%)	-0.11	13 (3%) 50 53	18, 27, 42, 54	0
1	D	430/439 (97%)	-0.07	4 (0%) 84 85	18, 27, 42, 49	0
1	E	430/439 (97%)	-0.23	8 (1%) 66 69	16, 24, 38, 46	0
1	F	430/439 (97%)	-0.04	11 (2%) 56 60	16, 25, 39, 49	0
All	All	2580/2634 (97%)	-0.15	43 (1%) 70 72	16, 25, 39, 54	0

The worst 5 of 43 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	195	SER	4.8
1	F	151	GLN	4.5
1	C	196	LYS	4.5
1	F	431	SER	4.3
1	E	192	ASP	3.9

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	GOL	E	1435	6/6	0.40	0.39	58,59,59,59	0
3	GOL	A	1433	6/6	0.43	0.24	57,57,58,59	0
3	GOL	C	1437	6/6	0.59	0.33	42,44,45,45	0
5	PE4	F	1435	24/24	0.65	0.20	49,53,55,57	0
4	SPM	C	1434	14/14	0.65	0.27	45,46,47,49	0
3	GOL	B	1434	6/6	0.67	0.28	45,46,48,48	0
3	GOL	E	1434	6/6	0.67	0.34	45,48,49,50	0
3	GOL	B	1435	6/6	0.67	0.28	53,55,55,55	0
5	PE4	B	1436	24/24	0.69	0.20	43,47,56,60	0
3	GOL	F	1434	6/6	0.69	0.32	47,51,53,54	0
3	GOL	E	1433	6/6	0.74	0.30	46,47,48,48	0
3	GOL	D	1432	6/6	0.77	0.35	52,52,54,55	0
3	GOL	C	1432	6/6	0.78	0.25	52,54,54,55	0
4	SPM	F	1433	14/14	0.78	0.18	33,43,48,49	0
3	GOL	D	1435	6/6	0.83	0.18	45,46,48,49	0
3	GOL	D	1437	6/6	0.84	0.22	43,45,46,47	0
3	GOL	C	1435	6/6	0.84	0.34	48,49,50,50	0
3	GOL	D	1436	6/6	0.86	0.20	48,49,49,50	0
4	SPM	D	1434	14/14	0.86	0.20	40,52,56,56	0
3	GOL	C	1436	6/6	0.88	0.12	39,41,42,42	0
4	SPM	B	1433	14/14	0.89	0.15	34,42,50,51	0
2	FAD	A	1432	53/53	0.96	0.09	16,20,22,22	0
2	FAD	F	1432	53/53	0.96	0.08	13,17,22,24	0
2	FAD	D	1433	53/53	0.96	0.08	16,21,23,26	0
2	FAD	B	1432	53/53	0.97	0.09	15,19,22,24	0
2	FAD	C	1433	53/53	0.97	0.10	15,19,23,25	0
2	FAD	E	1432	53/53	0.97	0.09	15,18,21,22	0

6.5 Other polymers [i](#)

There are no such residues in this entry.