



Full wwPDB X-ray Structure Validation Report ⓘ

May 14, 2018 – 02:34 PM EDT

PDB ID : 6C1V
Title : MBD2 in complex with double-stranded DNA
Authors : Lei, M.; Tempel, W.; Arrowsmith, C.H.; Bountra, C.; Edwards, A.M.; Min, J.; Structural Genomics Consortium (SGC)
Deposited on : 2018-01-05
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	rb-20031172
Percentile statistics	:	20171227.v01 (using entries in the PDB archive December 27th 2017)
Refmac	:	5.8.0158
CCP4	:	7.0 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20031172


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	111664	4477 (2.30-2.30)
Clashscore	122126	5072 (2.30-2.30)
Ramachandran outliers	120053	5022 (2.30-2.30)
Sidechain outliers	120020	5021 (2.30-2.30)
RSRZ outliers	108989	4374 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	79	<div> <div>4%</div> <div> <div></div> <div>77%</div> <div>9%</div> <div>14%</div> </div> </div>
1	B	79	<div> <div>9%</div> <div> <div></div> <div>80%</div> <div>5%</div> <div>14%</div> </div> </div>
1	E	79	<div> <div>6%</div> <div> <div></div> <div>80%</div> <div>6%</div> <div>14%</div> </div> </div>
1	F	79	<div> <div>8%</div> <div> <div></div> <div>84%</div> <div>•</div> <div>14%</div> </div> </div>
2	C	12	<div> <div></div> <div> <div></div> <div>92%</div> <div>8%</div> </div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	G	12	 92% 8%
3	D	12	 92% 8%
3	H	12	 92% 8%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	UNX	A	309	-	-	-	X

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 3061 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Methyl-CpG-binding domain protein 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	68	Total	C	N	O	S	0	0	0
			514	329	88	93	4			
1	B	68	Total	C	N	O	S	0	0	1
			495	317	87	89	2			
1	E	68	Total	C	N	O	S	0	0	1
			513	331	86	92	4			
1	F	68	Total	C	N	O	S	0	0	1
			502	325	85	89	3			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	142	GLY	-	expression tag	UNP Q9UBB5
B	142	GLY	-	expression tag	UNP Q9UBB5
E	142	GLY	-	expression tag	UNP Q9UBB5
F	142	GLY	-	expression tag	UNP Q9UBB5

- Molecule 2 is a DNA chain called 12-mer DNA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	12	Total	C	N	O	P	0	0	0
			249	118	50	70	11			
2	G	12	Total	C	N	O	P	0	0	0
			249	118	50	70	11			

- Molecule 3 is a DNA chain called 12-mer DNA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	D	12	Total	C	N	O	P	0	0	0
			237	114	42	70	11			
3	H	12	Total	C	N	O	P	0	0	0
			237	114	42	70	11			

- Molecule 4 is UNKNOWN ATOM OR ION (three-letter code: UNX) (formula: X).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	G	4	Total X 4 4	0	0
4	D	1	Total X 1 1	0	0
4	E	7	Total X 7 7	0	0
4	H	1	Total X 1 1	0	0
4	B	6	Total X 6 6	0	0
4	C	4	Total X 4 4	0	0
4	A	9	Total X 9 9	0	0
4	F	4	Total X 4 4	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	2	Total O 2 2	0	0
5	B	5	Total O 5 5	0	0
5	C	1	Total O 1 1	0	0
5	D	6	Total O 6 6	0	0
5	E	2	Total O 2 2	0	0
5	F	6	Total O 6 6	0	0
5	G	2	Total O 2 2	0	0
5	H	5	Total O 5 5	0	0



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: Methyl-CpG-binding domain protein 2





- Molecule 1: Methyl-CpG-binding domain protein 2




- Molecule 1: Methyl-CpG-binding domain protein 2



- Molecule 1: Methyl-CpG-binding domain protein 2



- Molecule 2: 12-mer DNA



- Molecule 2: 12-mer DNA






- Molecule 3: 12-mer DNA

Chain D:
92% 8%

A horizontal validation bar for Chain D. It is 92% green and 8% orange.



- Molecule 3: 12-mer DNA

Chain H:
92% 8%

A horizontal validation bar for Chain H. It is 92% green and 8% yellow.



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	37.01Å 40.20Å 105.16Å 84.50° 85.48° 62.71°	Depositor
Resolution (Å)	34.80 – 2.30 34.45 – 2.00	Depositor EDS
% Data completeness (in resolution range)	96.1 (34.80-2.30) 95.0 (34.45-2.00)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.19 (at 2.00Å)	Xtriage
Refinement program	REFMAC 5.8.0158	Depositor
R, R_{free}	0.233 , 0.268 0.243 , 0.273	Depositor DCC
R_{free} test set	1336 reflections (3.88%)	wwPDB-VP
Wilson B-factor (Å ²)	42.7	Xtriage
Anisotropy	0.203	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.28 , 38.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3061	wwPDB-VP
Average B, all atoms (Å ²)	51.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 14.10% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: UNX

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.88	0/527	0.97	4/709 (0.6%)
1	B	0.88	0/506	0.91	2/680 (0.3%)
1	E	0.88	0/526	0.98	2/707 (0.3%)
1	F	0.97	0/515	0.86	0/694
2	C	0.79	0/280	1.03	1/432 (0.2%)
2	G	0.87	1/280 (0.4%)	0.99	0/432
3	D	0.75	0/264	1.14	1/404 (0.2%)
3	H	0.78	0/264	1.13	1/404 (0.2%)
All	All	0.87	1/3162 (0.0%)	0.99	11/4462 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
3	D	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	G	11	DG	O3'-P	-5.60	1.54	1.61

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	8	DT	O5'-P-OP1	-7.50	98.95	105.70
1	A	195	ARG	NE-CZ-NH2	-6.20	117.20	120.30
3	H	5	DA	O5'-P-OP2	-6.08	100.22	105.70
1	A	149	ARG	NE-CZ-NH1	5.96	123.28	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	166	ARG	NE-CZ-NH2	-5.87	117.36	120.30
3	D	9	DT	O5'-P-OP2	-5.72	100.55	105.70
1	E	176	ASP	CB-CG-OD2	5.71	123.44	118.30
1	A	149	ARG	NE-CZ-NH2	-5.48	117.56	120.30
1	E	166	ARG	NE-CZ-NH2	-5.31	117.65	120.30
1	A	166	ARG	NE-CZ-NH1	5.24	122.92	120.30
1	B	149	ARG	NE-CZ-NH2	-5.13	117.73	120.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
3	D	9	DT	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	514	0	473	2	0
1	B	495	0	448	2	0
1	E	513	0	474	1	0
1	F	502	0	465	1	0
2	C	249	0	136	0	0
2	G	249	0	136	0	0
3	D	237	0	136	0	0
3	H	237	0	136	0	0
4	A	9	0	0	0	0
4	B	6	0	0	0	0
4	C	4	0	0	0	0
4	D	1	0	0	0	0
4	E	7	0	0	0	0
4	F	4	0	0	0	0
4	G	4	0	0	0	0
4	H	1	0	0	0	0
5	A	2	0	0	0	0
5	B	5	0	0	0	0
5	C	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	D	6	0	0	0	0
5	E	2	0	0	0	0
5	F	6	0	0	0	0
5	G	2	0	0	0	0
5	H	5	0	0	0	0
All	All	3061	0	2404	6	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (6) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:190:LYS:HB3	1:A:191:PRO:HD3	1.84	0.59
1:A:207:ASP:OD2	1:A:210:THR:HG23	2.10	0.52
1:F:166:ARG:HD3	1:F:171:SER:OG	2.10	0.51
1:E:190:LYS:HB3	1:E:191:PRO:HD3	1.96	0.48
1:B:166:ARG:HD3	1:B:171:SER:OG	2.16	0.46
1:B:181:SER:HB2	1:B:182:PRO:HD2	2.03	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	66/79 (84%)	65 (98%)	1 (2%)	0	100	100
1	B	66/79 (84%)	61 (92%)	5 (8%)	0	100	100
1	E	66/79 (84%)	65 (98%)	1 (2%)	0	100	100
1	F	66/79 (84%)	62 (94%)	4 (6%)	0	100	100
All	All	264/316 (84%)	253 (96%)	11 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	48/67 (72%)	48 (100%)	0	100	100
1	B	43/67 (64%)	43 (100%)	0	100	100
1	E	48/67 (72%)	47 (98%)	1 (2%)	56	73
1	F	46/67 (69%)	46 (100%)	0	100	100
All	All	185/268 (69%)	184 (100%)	1 (0%)	90	96

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	209	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry

Of 36 ligands modelled in this entry, 36 are unknown - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	68/79 (86%)	0.15	3 (4%) 34 41	37, 55, 68, 95	0
1	B	68/79 (86%)	0.55	7 (10%) 6 9	37, 62, 85, 90	0
1	E	68/79 (86%)	0.19	5 (7%) 14 20	36, 53, 68, 93	0
1	F	68/79 (86%)	0.32	6 (8%) 10 13	32, 56, 79, 89	0
2	C	12/12 (100%)	-0.22	0 100 100	34, 40, 55, 57	0
2	G	12/12 (100%)	-0.31	0 100 100	34, 39, 59, 63	0
3	D	12/12 (100%)	-0.35	0 100 100	35, 42, 56, 61	0
3	H	12/12 (100%)	-0.46	0 100 100	33, 40, 55, 57	0
All	All	320/364 (87%)	0.21	21 (6%) 18 24	32, 54, 83, 95	0

All (21) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	153	PRO	8.7
1	B	198	GLY	4.6
1	B	152	CYS	4.3
1	F	198	GLY	3.6
1	B	182	PRO	3.6
1	F	199	ASN	3.6
1	A	157	PRO	3.3
1	B	156	PRO	3.1
1	F	182	PRO	2.9
1	B	158	GLY	2.8
1	F	202	ASP	2.7
1	E	157	PRO	2.6
1	E	214	MET	2.6
1	E	215	PRO	2.5
1	E	213	MET	2.4
1	A	215	PRO	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	151	ASP	2.4
1	F	151	ASP	2.4
1	A	198	GLY	2.2
1	E	207	ASP	2.1
1	F	152	CYS	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	UNX	B	302	1/1	0.56	0.22	57,57,57,57	0
4	UNX	D	101	1/1	0.69	0.17	48,48,48,48	0
4	UNX	E	303	1/1	0.71	0.27	33,33,33,33	0
4	UNX	F	303	1/1	0.74	0.21	36,36,36,36	0
4	UNX	A	309	1/1	0.74	0.47	43,43,43,43	0
4	UNX	E	301	1/1	0.76	0.21	48,48,48,48	0
4	UNX	C	101	1/1	0.78	0.31	28,28,28,28	0
4	UNX	A	301	1/1	0.81	0.29	46,46,46,46	0
4	UNX	G	103	1/1	0.83	0.31	30,30,30,30	0
4	UNX	A	304	1/1	0.85	0.30	31,31,31,31	0
4	UNX	G	102	1/1	0.85	0.18	39,39,39,39	0
4	UNX	B	303	1/1	0.86	0.24	36,36,36,36	0
4	UNX	E	304	1/1	0.86	0.57	45,45,45,45	0
4	UNX	B	305	1/1	0.86	0.42	49,49,49,49	0
4	UNX	A	305	1/1	0.87	0.23	37,37,37,37	0
4	UNX	G	104	1/1	0.87	0.39	37,37,37,37	0
4	UNX	F	301	1/1	0.87	0.21	41,41,41,41	0
4	UNX	A	306	1/1	0.88	0.23	26,26,26,26	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	UNX	A	308	1/1	0.88	0.44	30,30,30,30	0
4	UNX	E	302	1/1	0.88	0.33	36,36,36,36	0
4	UNX	B	301	1/1	0.88	0.27	43,43,43,43	0
4	UNX	E	307	1/1	0.89	0.26	25,25,25,25	0
4	UNX	G	101	1/1	0.91	0.28	27,27,27,27	0
4	UNX	B	304	1/1	0.91	0.35	29,29,29,29	0
4	UNX	A	303	1/1	0.92	0.20	39,39,39,39	0
4	UNX	C	104	1/1	0.93	0.26	33,33,33,33	0
4	UNX	A	302	1/1	0.93	0.56	45,45,45,45	0
4	UNX	H	101	1/1	0.94	0.28	22,22,22,22	0
4	UNX	A	307	1/1	0.94	0.41	49,49,49,49	0
4	UNX	F	302	1/1	0.94	0.19	42,42,42,42	0
4	UNX	F	304	1/1	0.94	0.25	29,29,29,29	0
4	UNX	E	306	1/1	0.95	0.42	42,42,42,42	0
4	UNX	B	306	1/1	0.96	0.34	21,21,21,21	0
4	UNX	C	103	1/1	0.96	0.23	26,26,26,26	0
4	UNX	C	102	1/1	0.97	0.45	33,33,33,33	0
4	UNX	E	305	1/1	0.97	0.25	28,28,28,28	0

6.5 Other polymers ⓘ

There are no such residues in this entry.