



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2018 – 12:02 pm GMT

PDB ID : 1C2R
Title : MOLECULAR STRUCTURE OF CYTOCHROME C2 ISOLATED FROM RHODOBACTER CAPSULATUS DETERMINED AT 2.5 ANGSTROMS RESOLUTION
Authors : Benning, M.M.; Wesenberg, G.; Caffrey, M.S.; Bartsch, R.G.; Meyer, T.E.; Cusanovich, M.A.; Rayment, I.; Holden, H.M.
Deposited on : 1991-03-19
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7.3 (157068), CSD as539be (2018)
Xtriage (Phenix)	:	1.13
EDS	:	trunk30967
Percentile statistics	:	20171227.v01 (using entries in the PDB archive December 27th 2017)
Refmac	:	5.8.0158
CCP4	:	7.0 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	trunk30967


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	122126	4827 (2.50-2.50)
Ramachandran outliers	120053	4735 (2.50-2.50)
Sidechain outliers	120020	4737 (2.50-2.50)
RSRZ outliers	108989	4058 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	116	
1	B	116	

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 1906 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CYTOCHROME C2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	116	Total	C	N	O	S	0	0	0
			864	552	141	168	3			
1	B	116	Total	C	N	O	S	0	0	0
			864	552	141	168	3			

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
2	B	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	48	Total 48	O 48	0	0
3	B	44	Total 44	O 44	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: CYTOCHROME C2



• Molecule 1: CYTOCHROME C2



4 Data and refinement statistics

Property	Value	Source
Space group	H 3 2	Depositor
Cell constants a, b, c, α , β , γ	100.03Å 100.03Å 162.10Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	(Not available) – 2.50 25.47 – 2.50	Depositor EDS
% Data completeness (in resolution range)	(Not available) ((Not available)-2.50) 86.4 (25.47-2.50)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.99 (at 2.50Å)	Xtriage
Refinement program	TNT	Depositor
R, R_{free}	0.168 , (Not available) 0.153 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	31.2	Xtriage
Anisotropy	0.090	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 114.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	1906	wwPDB-VP
Average B, all atoms (Å ²)	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.51% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: HEM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.14	4/880 (0.5%)	1.44	12/1181 (1.0%)
1	B	1.11	6/880 (0.7%)	1.52	20/1181 (1.7%)
All	All	1.12	10/1760 (0.6%)	1.48	32/2362 (1.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	1	0

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	85	GLU	CD-OE2	6.28	1.32	1.25
1	A	85	GLU	CD-OE2	6.27	1.32	1.25
1	A	26	GLU	CD-OE2	6.09	1.32	1.25
1	B	9	GLU	CD-OE2	5.77	1.32	1.25
1	B	7	GLU	CD-OE2	5.53	1.31	1.25
1	A	7	GLU	CD-OE2	5.49	1.31	1.25
1	B	105	GLU	CD-OE2	5.21	1.31	1.25
1	A	50	GLU	CD-OE2	5.15	1.31	1.25
1	B	50	GLU	CD-OE2	5.09	1.31	1.25
1	B	70	GLU	CB-CG	5.07	1.61	1.52

All (32) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	78	ASP	CB-CG-OD2	-8.04	111.06	118.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	5	LYS	CB-CA-C	7.68	125.77	110.40
1	B	88	ASP	CB-CG-OD2	-7.31	111.72	118.30
1	B	106	ASP	CB-CG-OD2	-7.09	111.92	118.30
1	B	78	ASP	CB-CG-OD1	7.08	124.67	118.30
1	A	71	ASP	CB-CG-OD2	-6.93	112.06	118.30
1	A	55	ASP	CB-CG-OD2	-6.85	112.14	118.30
1	B	51	PHE	CB-CG-CD1	-6.40	116.32	120.80
1	B	106	ASP	CB-CG-OD1	6.30	123.97	118.30
1	A	89	ASP	CB-CG-OD2	-6.21	112.71	118.30
1	B	43	ARG	NE-CZ-NH1	6.19	123.40	120.30
1	B	2	ASP	CB-CG-OD2	-6.11	112.80	118.30
1	B	55	ASP	CB-CG-OD2	-6.09	112.81	118.30
1	B	101	ALA	N-CA-CB	-5.99	101.72	110.10
1	A	88	ASP	CB-CG-OD2	-5.98	112.91	118.30
1	B	14	LYS	CB-CA-C	5.88	122.16	110.40
1	B	8	LYS	N-CA-CB	5.86	121.16	110.60
1	A	89	ASP	CB-CG-OD1	5.67	123.41	118.30
1	A	102	LYS	N-CA-CB	-5.57	100.58	110.60
1	A	71	ASP	CB-CG-OD1	5.52	123.27	118.30
1	B	89	ASP	CB-CG-OD2	-5.52	113.33	118.30
1	B	2	ASP	CB-CG-OD1	5.45	123.21	118.30
1	A	23	ASP	CB-CG-OD2	-5.44	113.40	118.30
1	B	71	ASP	CB-CG-OD2	-5.43	113.41	118.30
1	B	88	ASP	CB-CG-OD1	5.43	123.18	118.30
1	A	5	LYS	N-CA-CB	5.40	120.32	110.60
1	A	88	ASP	CB-CG-OD1	5.32	123.09	118.30
1	A	106	ASP	CB-CG-OD2	-5.16	113.65	118.30
1	B	38	TYR	CB-CG-CD2	-5.11	117.94	121.00
1	B	22	PRO	N-CA-CB	5.07	109.39	103.30
1	B	12	LYS	CB-CA-C	5.06	120.53	110.40
1	B	89	ASP	CB-CG-OD1	5.02	122.82	118.30

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
1	A	5	LYS	CA

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	864	0	872	25	1
1	B	864	0	872	39	1
2	A	43	0	30	7	0
2	B	43	0	30	9	0
3	A	48	0	0	1	1
3	B	44	0	0	3	1
All	All	1906	0	1804	63	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 18.

All (63) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:16:CYS:SG	2:B:120:HEM:HAC	1.38	1.60
1:A:16:CYS:SG	2:A:120:HEM:HAC	1.63	1.38
1:B:16:CYS:SG	2:B:120:HEM:CAC	2.35	1.15
1:B:2:ASP:OD2	1:B:4:ALA:HB3	1.60	0.99
1:A:11:ASN:HD21	1:A:12:LYS:NZ	1.76	0.84
1:A:60:LEU:O	1:A:63:SER:OG	2.06	0.74
1:B:69:GLU:OE1	3:B:145:HOH:O	2.07	0.72
1:B:48:TYR:O	3:B:150:HOH:O	2.09	0.70
1:A:16:CYS:SG	2:A:120:HEM:C3C	2.85	0.68
1:A:20:ILE:HD13	1:A:26:GLU:HG3	1.76	0.67
1:B:16:CYS:CB	2:B:120:HEM:HAC	2.25	0.66
1:A:76:VAL:HG21	2:A:120:HEM:HMB2	1.77	0.66
1:A:50:GLU:HA	1:B:62:ALA:O	1.96	0.66
1:B:87:LEU:O	1:B:88:ASP:CB	2.46	0.63
1:A:11:ASN:HD21	1:A:12:LYS:HZ1	1.45	0.62
1:B:16:CYS:HA	1:B:31:ALA:HB3	1.84	0.60
1:B:16:CYS:SG	1:B:33:THR:HG21	2.42	0.60
1:B:96:MET:HG3	2:B:120:HEM:C4C	2.37	0.59
1:A:38:TYR:CZ	1:A:115:VAL:HG12	2.38	0.59
1:B:16:CYS:HB2	2:B:120:HEM:C3C	2.39	0.57
1:B:2:ASP:HB3	1:B:5:LYS:HB2	1.86	0.56

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:11:ASN:HD21	1:A:12:LYS:HZ3	1.51	0.56
1:B:96:MET:HG3	2:B:120:HEM:CHD	2.37	0.54
1:B:111:LEU:O	1:B:114:VAL:HG22	2.07	0.54
1:A:11:ASN:ND2	1:A:12:LYS:NZ	2.54	0.53
1:B:36:ASN:ND2	1:B:38:TYR:H	2.07	0.53
1:B:68:THR:HB	1:B:70:GLU:OE2	2.08	0.53
1:A:5:LYS:O	1:A:9:GLU:HG2	2.10	0.51
1:A:16:CYS:SG	2:A:120:HEM:CBC	2.93	0.50
1:A:100:LEU:HD23	3:A:133:HOH:O	2.11	0.49
1:B:17:HIS:HE1	1:B:35:PRO:HD2	1.76	0.49
1:B:44:THR:O	3:B:147:HOH:O	2.19	0.49
1:A:76:VAL:HG21	2:A:120:HEM:CMB	2.42	0.48
1:B:17:HIS:CE1	1:B:35:PRO:HD2	2.49	0.48
1:A:78:ASP:CG	1:A:99:LYS:HE2	2.34	0.48
1:B:16:CYS:SG	1:B:33:THR:CG2	3.01	0.48
1:B:73:ALA:CB	1:B:105:GLU:HG3	2.43	0.47
1:B:110:TYR:O	1:B:114:VAL:HG13	2.14	0.47
1:A:16:CYS:HA	1:A:31:ALA:HB3	1.96	0.47
1:B:73:ALA:HB1	1:B:105:GLU:HG3	1.97	0.47
1:B:86:LYS:HD3	1:B:86:LYS:HA	1.73	0.47
1:B:104:GLY:O	1:B:107:VAL:HG12	2.15	0.46
1:B:96:MET:HG3	2:B:120:HEM:C1D	2.51	0.46
1:B:9:GLU:OE1	1:B:12:LYS:HE3	2.16	0.46
1:A:46:GLY:HA2	1:A:53:TYR:CE2	2.51	0.45
1:A:68:THR:OG1	1:A:71:ASP:OD1	2.33	0.45
1:A:84:LYS:HG2	1:A:89:ASP:O	2.16	0.45
1:B:22:PRO:HD3	1:B:38:TYR:CE1	2.51	0.45
1:B:84:LYS:CG	1:B:92:ALA:HB3	2.47	0.45
1:B:3:ALA:O	1:B:7:GLU:N	2.42	0.44
1:B:20:ILE:H	1:B:36:ASN:HD21	1.66	0.44
1:B:84:LYS:HG2	1:B:92:ALA:HB3	2.00	0.43
1:A:16:CYS:CB	2:A:120:HEM:C3C	3.01	0.43
1:B:36:ASN:HD22	1:B:37:LEU:N	2.16	0.43
1:B:10:PHE:CZ	1:B:19:ILE:HD11	2.54	0.43
1:A:9:GLU:C	1:A:11:ASN:H	2.22	0.43
1:B:103:GLY:O	1:B:104:GLY:C	2.58	0.42
1:A:16:CYS:HB2	2:A:120:HEM:C3C	2.53	0.42
1:A:48:TYR:CG	1:A:49:PRO:HD2	2.54	0.42
1:B:16:CYS:CB	2:B:120:HEM:C3C	3.03	0.41
1:A:48:TYR:HA	1:A:49:PRO:HD3	1.93	0.40
1:B:13:CYS:HA	2:B:120:HEM:HHC	2.03	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:54:LYS:HA	1:B:54:LYS:HD2	1.84	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:5:LYS:NZ	1:B:114:VAL:O[5_565]	1.93	0.27
3:A:161:HOH:O	3:B:154:HOH:O[4_555]	2.00	0.20

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	114/116 (98%)	113 (99%)	1 (1%)	0	100	100
1	B	114/116 (98%)	106 (93%)	8 (7%)	0	100	100
All	All	228/232 (98%)	219 (96%)	9 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	86/86 (100%)	76 (88%)	10 (12%)	6	11

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	86/86 (100%)	79 (92%)	7 (8%)	13	25
All	All	172/172 (100%)	155 (90%)	17 (10%)	8	16

All (17) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	5	LYS
1	A	11	ASN
1	A	20	ILE
1	A	26	GLU
1	A	55	ASP
1	A	102	LYS
1	A	105	GLU
1	A	113	SER
1	A	114	VAL
1	A	116	LYS
1	B	2	ASP
1	B	12	LYS
1	B	14	LYS
1	B	36	ASN
1	B	54	LYS
1	B	63	SER
1	B	111	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	11	ASN
1	B	36	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	HEM	A	120	1	27,50,50	2.28	7 (25%)	17,82,82	2.22	6 (35%)
2	HEM	B	120	1	27,50,50	2.18	6 (22%)	17,82,82	2.80	8 (47%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEM	A	120	1	-	0/6/54/54	0/0/8/8
2	HEM	B	120	1	-	0/6/54/54	0/0/8/8

All (13) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	120	HEM	C3B-C2B	-6.82	1.30	1.40
2	B	120	HEM	C3C-C2C	-6.38	1.31	1.40
2	A	120	HEM	C3C-C2C	-5.84	1.32	1.40
2	B	120	HEM	C3B-C2B	-5.80	1.32	1.40
2	A	120	HEM	C1A-CHA	-3.01	1.32	1.40
2	A	120	HEM	C4B-NB	-2.30	1.31	1.36
2	B	120	HEM	C1A-CHA	-2.21	1.34	1.40
2	A	120	HEM	C4A-NA	2.20	1.40	1.36
2	B	120	HEM	C3C-CAC	2.53	1.52	1.47
2	A	120	HEM	CBB-CAB	2.81	1.48	1.29
2	B	120	HEM	CBC-CAC	3.06	1.49	1.29

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	120	HEM	CBC-CAC	3.12	1.50	1.29
2	B	120	HEM	CBB-CAB	3.19	1.50	1.29

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	120	HEM	CBD-CAD-C3D	-4.93	103.06	112.47
2	A	120	HEM	CBD-CAD-C3D	-3.75	105.31	112.47
2	B	120	HEM	CMD-C2D-C1D	-3.07	123.75	128.46
2	B	120	HEM	C4A-C3A-C2A	-2.54	105.23	107.00
2	B	120	HEM	CBA-CAA-C2A	-2.15	108.38	112.48
2	A	120	HEM	C4C-C3C-C2C	2.05	108.33	106.90
2	B	120	HEM	C3B-C4B-NB	2.17	112.02	109.21
2	A	120	HEM	CMB-C2B-C3B	2.76	129.90	124.88
2	A	120	HEM	CMC-C2C-C3C	2.83	130.03	124.88
2	B	120	HEM	CAA-CBA-CGA	3.70	118.99	112.66
2	A	120	HEM	CAA-CBA-CGA	3.83	119.21	112.66
2	B	120	HEM	CMB-C2B-C3B	3.97	132.10	124.88
2	A	120	HEM	CAD-CBD-CGD	3.99	119.48	112.66
2	B	120	HEM	CAD-CBD-CGD	6.41	123.61	112.66

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 16 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	120	HEM	7	0
2	B	120	HEM	9	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	116/116 (100%)	-0.81	2 (1%) 70 72	9, 24, 63, 100	0
1	B	116/116 (100%)	-0.81	0 100 100	9, 29, 65, 80	0
All	All	232/232 (100%)	-0.81	2 (0%) 84 85	9, 26, 65, 100	0

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	116	LYS	2.7
1	A	90	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	HEM	B	120	43/43	0.97	0.11	1,20,38,41	0
2	HEM	A	120	43/43	0.97	0.10	3,14,27,35	0

6.5 Other polymers [i](#)

There are no such residues in this entry.