



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 21, 2018 – 02:55 pm GMT

PDB ID : 1C4C
Title : BINDING OF EXOGENOUSLY ADDED CARBON MONOXIDE AT
THE ACTIVE SITE OF THE FE-ONLY HYDROGENASE (CPI) FROM
CLOSTRIDIUM PASTEURIANUM
Authors : Lemon, B.J.; Peters, J.W.
Deposited on : 1999-08-16
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.3 (157068), CSD as539be (2018)
Xtriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk30686


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	122078	3953 (2.40-2.40)
Ramachandran outliers	120005	3894 (2.40-2.40)
Sidechain outliers	119972	3895 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	574	 81% 17% .

2 Entry composition [i](#)


There are 5 unique types of molecules in this entry. The entry contains 4828 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PROTEIN (FE-ONLY HYDROGENASE).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	574	Total	C	N	O	S	0	0	0
			4461	2795	767	859	40			

- Molecule 2 is 2 IRON/2 SULFUR/6 CARBONYL/1 WATER INORGANIC CLUSTER (three-letter code: HC0) (formula: C₆H₇Fe₂O₇S₂).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	Fe	O	S	0	0
			17	6	2	7	2		

- Molecule 3 is IRON/SULFUR CLUSTER (three-letter code: SF4) (formula: Fe₄S₄).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	Fe	S	0	0
			8	4	4		
3	A	1	Total	Fe	S	0	0
			8	4	4		
3	A	1	Total	Fe	S	0	0
			8	4	4		
3	A	1	Total	Fe	S	0	0
			8	4	4		

- Molecule 4 is FE2/S2 (INORGANIC) CLUSTER (three-letter code: FES) (formula: Fe_2S_2).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	Fe	S	0	0
			4	2	2		

- Molecule 5 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	314	Total	O	0	0
			314	314		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: PROTEIN (FE-ONLY HYDROGENASE)



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	133.50Å 83.80Å 55.60Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 2.40	Depositor
% Data completeness (in resolution range)	97.7 (20.00-2.40)	Depositor
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	X-PLOR 3.857	Depositor
R, R_{free}	0.193 , 0.240	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	4828	wwPDB-VP
Average B, all atoms (Å ²)	30.0	wwPDB-VP

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SF4, FES, HC0

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.62	0/4534	1.15	18/6103 (0.3%)

There are no bond length outliers.

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	303	TRP	CD1-CG-CD2	8.32	112.95	106.30
1	A	303	TRP	CE2-CD2-CG	-7.99	100.91	107.30
1	A	548	LEU	CA-CB-CG	6.23	129.62	115.30
1	A	290	ASN	N-CA-C	6.18	127.68	111.00
1	A	373	ARG	NE-CZ-NH2	-6.16	117.22	120.30
1	A	303	TRP	CB-CG-CD1	-6.15	119.01	127.00
1	A	387	MET	CA-CB-CG	6.08	123.64	113.30
1	A	303	TRP	CG-CD2-CE3	6.04	139.34	133.90
1	A	86	ARG	NE-CZ-NH1	5.77	123.19	120.30
1	A	213	ARG	NE-CZ-NH1	5.75	123.17	120.30
1	A	52	GLU	CA-CB-CG	5.66	125.86	113.40
1	A	286	ARG	NE-CZ-NH1	5.49	123.05	120.30
1	A	373	ARG	NE-CZ-NH1	5.41	123.00	120.30
1	A	303	TRP	CG-CD1-NE1	-5.33	104.77	110.10
1	A	144	ARG	NE-CZ-NH1	5.25	122.92	120.30
1	A	364	ARG	NE-CZ-NH1	5.18	122.89	120.30
1	A	382	ARG	NE-CZ-NH1	5.10	122.85	120.30
1	A	105	GLU	CA-CB-CG	5.09	124.59	113.40

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4461	0	4430	50	0
2	A	17	0	0	3	0
3	A	32	0	0	1	0
4	A	4	0	0	1	0
5	A	314	0	0	8	0
All	All	4828	0	4430	52	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (52) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:472:ASN:HD21	1:A:501:GLY:H	1.27	0.83
1:A:512:VAL:HG11	1:A:517:LEU:HD13	1.71	0.72
1:A:467:ASN:H	1:A:492:HIS:HD2	1.38	0.72
1:A:167:LYS:HE3	1:A:180:GLU:HG2	1.84	0.60
1:A:45:LYS:HB3	4:A:585:FES:S1	2.42	0.59
1:A:43:ILE:HD11	5:A:615:HOH:O	2.05	0.56
1:A:392:LYS:HG3	5:A:620:HOH:O	2.05	0.56
1:A:137:SER:O	1:A:208:LYS:HG3	2.08	0.54
1:A:432:LYS:HE2	1:A:460:GLU:O	2.07	0.53
1:A:76:SER:HB3	1:A:79:VAL:HB	1.89	0.53
2:A:580:HC0:S2	2:A:580:HC0:O1	2.66	0.52
2:A:580:HC0:S1	2:A:580:HC0:O1	2.67	0.52
1:A:307:ALA:O	1:A:315:LEU:HD21	2.09	0.52
1:A:52:GLU:HG3	1:A:73:ASN:HB2	1.91	0.52
1:A:6:ILE:HG23	1:A:72:ILE:HB	1.92	0.51
1:A:7:ASN:HD21	1:A:74:THR:H	1.58	0.51
1:A:82:LYS:O	1:A:86:ARG:HG2	2.11	0.51
1:A:155:ASN:O	1:A:159:LYS:HG2	2.11	0.50
1:A:67:GLU:HB3	5:A:835:HOH:O	2.11	0.50
1:A:489:LYS:HG2	5:A:888:HOH:O	2.11	0.50
1:A:488:GLU:HB2	5:A:888:HOH:O	2.12	0.50
1:A:512:VAL:O	1:A:513:ASN:HB3	2.13	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:546:THR:O	1:A:550:LYS:HD3	2.13	0.49
1:A:294:PRO:O	1:A:317:ASN:HB3	2.13	0.49
1:A:472:ASN:ND2	1:A:501:GLY:H	2.02	0.48
1:A:556:PHE:HE1	1:A:567:ILE:HD11	1.77	0.48
1:A:46:CYS:SG	1:A:48:ILE:HG12	2.53	0.48
1:A:102:ASN:HD22	1:A:103:ARG:HG2	1.78	0.48
1:A:210:HIS:HD2	1:A:378:VAL:H	1.63	0.47
1:A:472:ASN:HD21	1:A:501:GLY:N	2.04	0.46
1:A:67:GLU:HB2	1:A:70:MET:HG3	1.98	0.46
1:A:231:PRO:HD3	1:A:269:ASN:ND2	2.31	0.46
1:A:467:ASN:H	1:A:492:HIS:CD2	2.26	0.46
1:A:107:CYS:O	1:A:108:GLU:HB3	2.17	0.45
1:A:355:CYS:HB2	3:A:581:SF4:S3	2.55	0.45
1:A:210:HIS:CD2	1:A:378:VAL:H	2.35	0.45
1:A:445:TYR:O	1:A:448:VAL:HG22	2.16	0.44
1:A:179:ALA:HB1	1:A:188:THR:HG21	2.01	0.43
1:A:388:ILE:HG23	1:A:393:ILE:HB	2.00	0.43
1:A:514:PRO:HA	1:A:517:LEU:HB3	2.01	0.43
1:A:23:ALA:HB1	1:A:28:ILE:O	2.18	0.43
1:A:333:LYS:HB2	5:A:657:HOH:O	2.19	0.43
1:A:277:MET:HG3	1:A:559:PRO:HG2	2.00	0.42
1:A:326:GLN:HE22	1:A:362:ALA:HA	1.83	0.42
1:A:503:CYS:CB	2:A:580:HC0:C4	2.97	0.42
1:A:392:LYS:HE2	5:A:620:HOH:O	2.20	0.42
1:A:355:CYS:SG	1:A:358:LYS:HG2	2.60	0.41
1:A:53:VAL:HG13	1:A:56:THR:OG1	2.20	0.41
1:A:21:LYS:HE2	1:A:38:ASN:HD21	1.86	0.41
1:A:432:LYS:HG3	1:A:461:ILE:HG12	2.02	0.41
1:A:137:SER:HB3	1:A:140:LEU:O	2.21	0.40
1:A:558:LYS:HE3	5:A:634:HOH:O	2.20	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	572/574 (100%)	533 (93%)	27 (5%)	12 (2%)	8	9

All (12) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	76	SER
1	A	120	ALA
1	A	121	SER
1	A	44	ASN
1	A	108	GLU
1	A	109	PHE
1	A	308	GLU
1	A	516	ASP
1	A	77	ASP
1	A	290	ASN
1	A	513	ASN
1	A	99	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	490/490 (100%)	465 (95%)	25 (5%)	26	42

All (25) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	24	ARG
1	A	39	CYS
1	A	40	ASN
1	A	44	ASN
1	A	52	GLU
1	A	75	ASN
1	A	77	ASP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	100	PRO
1	A	102	ASN
1	A	105	GLU
1	A	106	ASN
1	A	177	ILE
1	A	180	GLU
1	A	210	HIS
1	A	211	MET
1	A	215	LYS
1	A	316	ASN
1	A	345	LYS
1	A	465	LYS
1	A	513	ASN
1	A	518	GLU
1	A	519	LYS
1	A	537	LEU
1	A	545	ASN
1	A	568	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (16) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	7	ASN
1	A	38	ASN
1	A	102	ASN
1	A	195	GLN
1	A	210	HIS
1	A	269	ASN
1	A	306	GLN
1	A	316	ASN
1	A	325	GLN
1	A	326	GLN
1	A	462	ASN
1	A	472	ASN
1	A	492	HIS
1	A	511	HIS
1	A	533	GLN
1	A	545	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	HC0	A	580	1	1,19,19	0.78	0	0,52,52	0.00	-
3	SF4	A	581	1	0,12,12	0.00	-	0,24,24	0.00	-
3	SF4	A	582	1	0,12,12	0.00	-	0,24,24	0.00	-
3	SF4	A	583	1	0,12,12	0.00	-	0,24,24	0.00	-
3	SF4	A	584	1	0,12,12	0.00	-	0,24,24	0.00	-
4	FES	A	585	1	0,4,4	0.00	-	0,4,4	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HC0	A	580	1	-	0/0/105/105	0/0/3/3
3	SF4	A	581	1	-	0/0/48/48	0/6/5/5
3	SF4	A	582	1	-	0/0/48/48	0/6/5/5
3	SF4	A	583	1	-	0/0/48/48	0/6/5/5

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	SF4	A	584	1	-	0/0/48/48	0/6/5/5
4	FES	A	585	1	-	0/0/4/4	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	580	HC0	3	0
3	A	581	SF4	1	0
4	A	585	FES	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.