



# Full wwPDB X-ray Structure Validation Report ⓘ

Jun 11, 2018 – 06:51 PM EDT

PDB ID : 2C4A  
Title : Structure of Neuraminidase Subtype N9 Complexed with 30 MM Sialic Acid (NANA, NEU5AC), Crystal Soaked for 3 Hours at 291 K.  
Authors : Rudino-Pinera, E.; Tunnah, P.; Crennell, S.J.; Webster, R.G.; Laver, W.G.; Garman, E.F.  
Deposited on : 2005-10-17  
Resolution : 2.15 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.3 (157068), CSD as539be (2018)  
Xtriage (Phenix) : 1.13  
EDS : rb-20031172  
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)  
Refmac : 5.8.0158  
CCP4 : 7.0 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20031172


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*


The reported resolution of this entry is 2.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	111664	1287 (2.16-2.16)
Clashscore	122126	1390 (2.16-2.16)
Ramachandran outliers	120053	1368 (2.16-2.16)
Sidechain outliers	120020	1367 (2.16-2.16)
RSRZ outliers	108989	1262 (2.16-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	388	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	BMA	A	1476	-	-	X	-

## 2 Entry composition [i](#)


There are 7 unique types of molecules in this entry. The entry contains 3516 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called NEURAMINIDASE SUBTYPE N9.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	388	3067	1914	538	592	23	0	0	0

- Molecule 2 is O-SIALIC ACID (three-letter code: SIA) (formula:  $C_{11}H_{19}NO_9$ ).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
2	A	1	21	11	1	9	0	0
2	A	1	21	11	1	9	0	0

- Molecule 3 is N-ACETYL-D-GLUCOSAMINE (three-letter code: NAG) (formula:  $C_8H_{15}NO_6$ ).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	A	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 4 is BETA-D-MANNOSE (three-letter code: BMA) (formula:  $C_6H_{12}O_6$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			11	6	5		

- Molecule 5 is ALPHA-D-MANNOSE (three-letter code: MAN) (formula: C<sub>6</sub>H<sub>12</sub>O<sub>6</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			11	6	5		
5	A	1	Total	C	O	0	0
			11	6	5		
5	A	1	Total	C	O	0	0
			11	6	5		
5	A	1	Total	C	O	0	0
			11	6	5		
5	A	1	Total	C	O	0	0
			11	6	5		
5	A	1	Total	C	O	0	0
			11	6	5		

- Molecule 6 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1	Total	Ca	0	0
			1	1		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	259	Total 259	O 259	0	0

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: NEURAMINIDASE SUBTYPE N9



## 4 Data and refinement statistics

Property	Value	Source
Space group	I 4 3 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	181.34Å 181.34Å 181.34Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.10 – 2.15 38.66 – 2.15	Depositor EDS
% Data completeness (in resolution range)	99.9 (30.10-2.15) 99.9 (38.66-2.15)	Depositor EDS
$R_{merge}$	0.12	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	7.00 (at 2.16Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, $R_{free}$	0.158 , 0.205 0.159 , 0.205	Depositor DCC
$R_{free}$ test set	1401 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	13.6	Xtriage
Anisotropy	0.000	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 29.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	3516	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	12.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.89% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MAN, CA, BMA, NAG, SIA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.98	5/3150 (0.2%)	0.86	9/4290 (0.2%)

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	416	GLU	CD-OE1	-21.55	1.01	1.25
1	A	392	GLN	C-N	14.11	1.58	1.33
1	A	333	VAL	C-N	13.54	1.57	1.33
1	A	385	ASP	C-N	6.03	1.48	1.34
1	A	465	GLU	CG-CD	5.41	1.60	1.51

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	416	GLU	OE1-CD-OE2	16.27	142.82	123.30
1	A	127	ASP	CB-CG-OD2	-11.83	107.65	118.30
1	A	364	ARG	NE-CZ-NH1	9.41	125.00	120.30
1	A	416	GLU	CG-CD-OE2	-9.38	99.53	118.30
1	A	364	ARG	NE-CZ-NH2	-8.59	116.00	120.30
1	A	172	ARG	NE-CZ-NH2	-6.94	116.83	120.30
1	A	392	GLN	C-N-CA	-5.51	110.73	122.30
1	A	386	ASP	CB-CG-OD2	5.23	123.01	118.30
1	A	172	ARG	NE-CZ-NH1	5.07	122.83	120.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3067	0	2895	39	0
2	A	42	0	36	0	0
3	A	70	0	64	12	0
4	A	11	0	10	7	0
5	A	66	0	59	7	0
6	A	1	0	0	0	0
7	A	259	0	0	2	0
All	All	3516	0	3064	51	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (51) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:146:ASN:HD21	3:A:1475:NAG:C1	1.09	1.57
1:A:200:ASN:HD21	3:A:1471:NAG:C1	0.87	1.51
4:A:1476:BMA:C3	5:A:1477:MAN:C1	2.37	1.03
1:A:172:ARG:HD3	1:A:209:ARG:NH2	1.74	1.00
3:A:1472:NAG:HO4	4:A:1476:BMA:C1	1.64	0.92
1:A:168:THR:H	1:A:170:ASN:HD21	1.13	0.91
1:A:87:LEU:H	1:A:233:HIS:HD2	1.18	0.91
3:A:1473:NAG:HO4	3:A:1474:NAG:C1	1.86	0.86
4:A:1476:BMA:O3	5:A:1477:MAN:C2	2.26	0.84
1:A:172:ARG:HD3	1:A:209:ARG:HH22	1.43	0.81
1:A:200:ASN:HD21	3:A:1471:NAG:C2	1.95	0.78
1:A:168:THR:H	1:A:170:ASN:ND2	1.86	0.73
1:A:87:LEU:H	1:A:233:HIS:CD2	2.06	0.73
3:A:1472:NAG:C4	4:A:1476:BMA:C1	2.69	0.70
4:A:1476:BMA:H3	5:A:1477:MAN:C1	2.23	0.68
1:A:292:ARG:HE	1:A:294:ASN:HD22	1.44	0.65
1:A:172:ARG:CD	1:A:209:ARG:NH2	2.57	0.65
1:A:200:ASN:CG	3:A:1471:NAG:C1	2.64	0.63
5:A:1478:MAN:C2	5:A:1479:MAN:C1	2.78	0.60
1:A:422:PHE:CZ	1:A:446:MET:HG3	2.37	0.60

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:98:HIS:HE1	1:A:419:ARG:HH11	1.49	0.59
1:A:82:ARG:N	7:A:2001:HOH:O	2.36	0.59
1:A:200:ASN:ND2	3:A:1471:NAG:C2	2.59	0.58
1:A:98:HIS:CE1	1:A:419:ARG:HH11	2.22	0.57
1:A:200:ASN:ND2	3:A:1471:NAG:O5	2.34	0.56
1:A:465:GLU:CD	1:A:465:GLU:H	2.09	0.56
1:A:172:ARG:CD	1:A:209:ARG:HH22	2.17	0.56
5:A:1478:MAN:HO2	5:A:1479:MAN:C1	2.10	0.55
1:A:172:ARG:HD3	1:A:209:ARG:CZ	2.35	0.55
1:A:97:TRP:H	1:A:395:GLN:HE22	1.55	0.52
3:A:1473:NAG:C4	3:A:1474:NAG:C1	2.87	0.52
4:A:1476:BMA:C6	5:A:1480:MAN:C1	2.86	0.52
1:A:228:SER:HB3	1:A:350:LYS:HE2	1.92	0.51
3:A:1472:NAG:O4	4:A:1476:BMA:C2	2.55	0.50
1:A:367:SER:HB2	1:A:400:ASN:HD21	1.77	0.49
5:A:1480:MAN:O3	5:A:1482:MAN:C2	2.60	0.49
1:A:98:HIS:HD2	1:A:99:ILE:O	1.96	0.48
1:A:168:THR:OG1	1:A:170:ASN:ND2	2.46	0.48
1:A:318:CYS:O	1:A:335:GLY:HA3	2.14	0.48
1:A:204:VAL:HG22	1:A:206:TRP:NE1	2.30	0.47
1:A:330:ASP:CG	1:A:364:ARG:HH22	2.18	0.47
1:A:240:VAL:HG22	1:A:254:ILE:HD13	1.97	0.46
1:A:320:PRO:HD3	1:A:386:ASP:O	2.16	0.46
1:A:140:ILE:HD11	7:A:2223:HOH:O	2.16	0.45
1:A:292:ARG:HH21	1:A:294:ASN:ND2	2.15	0.45
1:A:364:ARG:HD2	1:A:375:GLU:OE2	2.16	0.45
1:A:121:TYR:CG	1:A:228:SER:HA	2.55	0.42
1:A:146:ASN:CG	3:A:1475:NAG:C1	2.79	0.42
1:A:174:GLU:OE1	1:A:209:ARG:NH2	2.53	0.41
1:A:386:ASP:OD1	1:A:386:ASP:N	2.53	0.41
1:A:292:ARG:HH21	1:A:294:ASN:HD21	1.69	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	386/388 (100%)	367 (95%)	19 (5%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	341/341 (100%)	333 (98%)	8 (2%)	53	56

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	82	ARG
1	A	170	ASN
1	A	204	VAL
1	A	221	ASN
1	A	254	ILE
1	A	364	ARG
1	A	416	GLU
1	A	465	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	95	ASN
1	A	98	HIS
1	A	146	ASN
1	A	170	ASN
1	A	200	ASN
1	A	216	ASN
1	A	221	ASN
1	A	233	HIS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	294	ASN
1	A	346	ASN
1	A	392	GLN
1	A	395	GLN
1	A	400	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

Of 15 ligands modelled in this entry, 1 is monoatomic - leaving 14 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	SIA	A	1469	-	18,21,21	1.32	2 (11%)	19,31,31	1.19	3 (15%)
2	SIA	A	1470	-	18,21,21	0.92	2 (11%)	19,31,31	1.04	2 (10%)
3	NAG	A	1471	1,3	14,14,15	0.60	0	17,19,21	2.01	6 (35%)
3	NAG	A	1472	3,4	14,14,15	0.38	0	17,19,21	1.39	2 (11%)
3	NAG	A	1473	1,3	14,14,15	0.43	0	17,19,21	1.48	3 (17%)
3	NAG	A	1474	3	14,14,15	0.52	0	17,19,21	1.78	3 (17%)
3	NAG	A	1475	1	14,14,15	0.34	0	17,19,21	0.90	2 (11%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	BMA	A	1476	3,5	11,11,12	0.26	0	15,15,17	1.45	4 (26%)
5	MAN	A	1477	5,4	11,11,12	0.50	0	15,15,17	1.88	5 (33%)
5	MAN	A	1478	5	11,11,12	0.55	0	15,15,17	1.74	4 (26%)
5	MAN	A	1479	5	11,11,12	0.72	0	15,15,17	1.56	2 (13%)
5	MAN	A	1480	5,4	11,11,12	0.48	0	15,15,17	1.56	2 (13%)
5	MAN	A	1481	5	11,11,12	0.67	0	15,15,17	2.04	5 (33%)
5	MAN	A	1482	5	11,11,12	0.49	0	15,15,17	2.12	3 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	SIA	A	1469	-	-	0/14/38/38	0/1/1/1
2	SIA	A	1470	-	-	0/14/38/38	0/1/1/1
3	NAG	A	1471	1,3	-	0/6/23/26	0/1/1/1
3	NAG	A	1472	3,4	-	0/6/23/26	0/1/1/1
3	NAG	A	1473	1,3	-	0/6/23/26	0/1/1/1
3	NAG	A	1474	3	-	0/6/23/26	0/1/1/1
3	NAG	A	1475	1	-	0/6/23/26	0/1/1/1
4	BMA	A	1476	3,5	-	0/2/19/22	0/1/1/1
5	MAN	A	1477	5,4	-	0/2/19/22	0/1/1/1
5	MAN	A	1478	5	-	0/2/19/22	0/1/1/1
5	MAN	A	1479	5	-	0/2/19/22	0/1/1/1
5	MAN	A	1480	5,4	-	0/2/19/22	0/1/1/1
5	MAN	A	1481	5	-	0/2/19/22	0/1/1/1
5	MAN	A	1482	5	-	0/2/19/22	0/1/1/1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1469	SIA	C3-C2	2.06	1.54	1.51
2	A	1470	SIA	O2-C2	2.17	1.42	1.39
2	A	1470	SIA	C3-C2	2.35	1.54	1.51
2	A	1469	SIA	O2-C2	4.51	1.44	1.39

All (46) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	1477	MAN	O2-C2-C1	-4.25	100.60	109.17
5	A	1480	MAN	C3-C4-C5	-3.71	103.60	110.24
3	A	1471	NAG	C4-C3-C2	-3.51	105.87	111.02
3	A	1471	NAG	O4-C4-C3	-3.46	102.27	110.34
4	A	1476	BMA	O3-C3-C2	-3.21	104.06	110.04
3	A	1473	NAG	C1-C2-N2	-3.00	105.36	110.49
3	A	1471	NAG	C8-C7-N2	-2.97	110.90	116.10
3	A	1474	NAG	O5-C1-C2	-2.78	107.68	111.52
3	A	1474	NAG	C4-C3-C2	-2.76	106.97	111.02
3	A	1471	NAG	C1-C2-N2	-2.75	105.78	110.49
3	A	1472	NAG	C4-C3-C2	-2.51	107.35	111.02
2	A	1470	SIA	C4-C5-N5	-2.49	105.28	110.40
3	A	1475	NAG	C1-C2-N2	-2.45	106.31	110.49
5	A	1478	MAN	O3-C3-C2	-2.31	105.74	110.04
2	A	1469	SIA	C6-C5-N5	-2.25	107.05	110.94
5	A	1477	MAN	C3-C4-C5	-2.18	106.34	110.24
4	A	1476	BMA	C6-C5-C4	-2.17	107.86	112.99
4	A	1476	BMA	C1-C2-C3	-2.15	106.94	109.66
3	A	1473	NAG	O4-C4-C3	-2.14	105.34	110.34
5	A	1477	MAN	O6-C6-C5	-2.13	103.86	111.29
2	A	1470	SIA	O2-C2-O6	-2.12	104.85	109.69
4	A	1476	BMA	O6-C6-C5	-2.02	104.23	111.29
5	A	1481	MAN	O2-C2-C3	2.03	114.16	110.19
3	A	1475	NAG	O5-C5-C6	2.04	110.38	107.15
2	A	1469	SIA	O6-C6-C7	2.11	110.62	107.33
2	A	1469	SIA	O2-C2-C3	2.15	112.22	109.33
5	A	1477	MAN	C1-C2-C3	2.15	112.38	109.66
3	A	1471	NAG	C3-C4-C5	2.26	114.29	110.24
5	A	1481	MAN	C1-C2-C3	2.28	112.54	109.66
5	A	1478	MAN	C1-C2-C3	2.56	112.90	109.66
5	A	1481	MAN	O5-C5-C6	2.62	111.29	107.15
3	A	1472	NAG	O5-C1-C2	2.66	115.19	111.52
3	A	1471	NAG	O5-C5-C6	2.80	111.58	107.15
5	A	1478	MAN	O5-C1-C2	2.87	115.25	110.78
3	A	1473	NAG	O5-C5-C6	3.27	112.32	107.15
5	A	1482	MAN	O5-C1-C2	3.55	116.31	110.78
5	A	1480	MAN	C1-O5-C5	3.64	117.19	112.19
5	A	1482	MAN	C1-C2-C3	3.64	114.26	109.66
5	A	1479	MAN	O5-C1-C2	3.77	116.66	110.78
5	A	1479	MAN	C1-O5-C5	3.93	117.59	112.19
5	A	1481	MAN	O5-C1-C2	3.94	116.92	110.78
5	A	1478	MAN	C1-O5-C5	4.14	117.88	112.19
5	A	1477	MAN	C1-O5-C5	4.29	118.09	112.19

Continued on next page...

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1474	NAG	C2-N2-C7	5.03	130.28	122.94
5	A	1481	MAN	C1-O5-C5	5.03	119.11	112.19
5	A	1482	MAN	C1-O5-C5	5.86	120.25	112.19

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

11 monomers are involved in 19 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1471	NAG	5	0
3	A	1472	NAG	3	0
3	A	1473	NAG	2	0
3	A	1474	NAG	2	0
3	A	1475	NAG	2	0
4	A	1476	BMA	7	0
5	A	1477	MAN	3	0
5	A	1478	MAN	2	0
5	A	1479	MAN	2	0
5	A	1480	MAN	2	0
5	A	1482	MAN	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	388/388 (100%)	-0.75	1 (0%) 93 95	6, 10, 17, 38	0

All (1) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	82	ARG	2.4

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	NAG	A	1474	14/15	0.77	0.26	46,49,51,52	0
5	MAN	A	1481	11/12	0.78	0.23	55,56,57,57	0
5	MAN	A	1482	11/12	0.86	0.23	35,41,41,41	0
3	NAG	A	1475	14/15	0.87	0.20	36,38,40,41	0
5	MAN	A	1477	11/12	0.91	0.10	15,18,21,26	0
5	MAN	A	1480	11/12	0.92	0.17	24,27,30,31	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	NAG	A	1473	14/15	0.94	0.14	21,23,25,29	0
3	NAG	A	1471	14/15	0.95	0.12	11,17,22,23	0
4	BMA	A	1476	11/12	0.96	0.09	13,16,19,21	0
3	NAG	A	1472	14/15	0.96	0.10	13,16,19,21	0
2	SIA	A	1469	21/21	0.96	0.08	8,12,14,19	0
5	MAN	A	1478	11/12	0.97	0.07	14,16,17,18	0
5	MAN	A	1479	11/12	0.97	0.07	13,14,16,17	0
2	SIA	A	1470	21/21	0.97	0.08	15,17,19,19	0
6	CA	A	1483	1/1	0.99	0.04	12,12,12,12	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.