



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2018 – 10:35 pm GMT

PDB ID : 2C7U
Title : Conflicting selective forces affect CD8 T-cell receptor contact sites in an HLA-A2 immunodominant HIV epitope.
Authors : Iversen, A.K.; Stewart-Jones, G.; Learn, G.H.; Christie, N.; Sylvester-Hviid, C.; Armitage, A.E.; Kaul, R.; Beattie, T.; Lee, J.K.; Li, Y.; Chotiyarnwong, P.; Dong, T.; Xu, X.; Luscher, M.A.; MacDonald, K.; Ullum, H.; Klarlund-Pedersen, B.; Skinhoj, P.; Fugger, J.L.; Buus, S.; Mullins, J.I.; Jones, E.Y.; van der Merwe, P.A.; McMichael, A.J.
Deposited on : 2005-11-29
Resolution : 2.38 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	trunk30967
Percentile statistics	:	20171227.v01 (using entries in the PDB archive December 27th 2017)
Refmac	:	5.8.0158
CCP4	:	7.0 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	trunk30967


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.38 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	122126	5405 (2.40-2.36)
Ramachandran outliers	120053	5324 (2.40-2.36)
Sidechain outliers	120020	5326 (2.40-2.36)
RSRZ outliers	108989	4741 (2.40-2.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	276	
1	D	276	
2	B	100	
2	E	100	
3	C	9	
3	F	9	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 6519 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HLA CLASS I HISTOCOMPATIBILITY ANTIGEN, A-2 ALPHA CHAIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	275	Total	C	N	O	S	0	0	0
			2246	1403	409	425	9			
1	D	275	Total	C	N	O	S	0	0	0
			2246	1403	409	425	9			

- Molecule 2 is a protein called BETA-2-MICROGLOBULIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	100	Total	C	N	O	S	0	0	0
			836	533	141	158	4			
2	E	100	Total	C	N	O	S	0	0	0
			836	533	141	158	4			

- Molecule 3 is a protein called GAG PROTEIN.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	C	9	Total	C	N	O	0	0	0
			68	46	10	12			
3	F	9	Total	C	N	O	0	0	0
			68	46	10	12			

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	69	Total	O	0	0
			69	69		
4	B	37	Total	O	0	0
			37	37		
4	C	4	Total	O	0	0
			4	4		
4	D	69	Total	O	0	0
			69	69		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	E	38	Total	O	0	0
			38	38		
4	F	2	Total	O	0	0
			2	2		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: HLA CLASS I HISTOCOMPATIBILITY ANTIGEN, A-2 ALPHA CHAIN




- Molecule 1: HLA CLASS I HISTOCOMPATIBILITY ANTIGEN, A-2 ALPHA CHAIN




- Molecule 2: BETA-2-MICROGLOBULIN




- Molecule 2: BETA-2-MICROGLOBULIN



- Molecule 3: GAG PROTEIN



• Molecule 3: GAG PROTEIN



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	63.16Å 93.85Å 81.50Å 90.00° 89.99° 90.00°	Depositor
Resolution (Å)	81.38 – 2.38 28.03 – 2.38	Depositor EDS
% Data completeness (in resolution range)	97.0 (81.38-2.38) 89.0 (28.03-2.38)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.00 (at 2.39Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.248 , 0.314 0.244 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	42.0	Xtriage
Anisotropy	0.357	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 49.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.478 for h,-k,-l	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	6519	wwPDB-VP
Average B, all atoms (Å ²)	55.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.99% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.81	4/2311 (0.2%)	0.65	1/3137 (0.0%)
1	D	0.70	5/2311 (0.2%)	0.68	1/3137 (0.0%)
2	B	0.66	0/859	0.67	0/1162
2	E	0.65	0/859	0.68	0/1162
3	C	0.58	0/68	0.73	0/92
3	F	0.58	0/68	0.77	0/92
All	All	0.73	9/6476 (0.1%)	0.67	2/8782 (0.0%)

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	58	GLU	CD-OE1	20.57	1.48	1.25
1	A	58	GLU	CD-OE2	17.96	1.45	1.25
1	D	219	ARG	CZ-NH2	11.45	1.48	1.33
1	D	219	ARG	CZ-NH1	10.52	1.46	1.33
1	D	219	ARG	CD-NE	7.92	1.59	1.46
1	D	220	ASP	CG-OD1	7.71	1.43	1.25
1	D	144	LYS	CD-CE	7.43	1.69	1.51
1	A	220	ASP	CG-OD2	7.11	1.41	1.25
1	A	220	ASP	CG-OD1	5.99	1.39	1.25

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	219	ARG	NE-CZ-NH2	-12.09	114.25	120.30
1	A	58	GLU	OE1-CD-OE2	5.99	130.49	123.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2246	0	2096	42	0
1	D	2246	0	2096	44	0
2	B	836	0	803	4	0
2	E	836	0	803	10	0
3	C	68	0	76	3	0
3	F	68	0	76	2	0
4	A	69	0	0	19	0
4	B	37	0	0	1	0
4	C	4	0	0	1	0
4	D	69	0	0	16	0
4	E	38	0	0	5	0
4	F	2	0	0	1	0
All	All	6519	0	5950	102	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (102) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:119:ASP:HB3	4:A:2041:HOH:O	1.60	1.00
1:A:82:ARG:HH21	1:A:89:GLU:HG3	1.29	0.93
1:D:67:VAL:HG22	4:D:2022:HOH:O	1.67	0.93
1:D:11:SER:HB2	1:D:74:HIS:HD2	1.33	0.92
1:A:6:ARG:HH11	1:A:6:ARG:HG3	1.35	0.90
1:D:54:GLN:HB2	4:D:2017:HOH:O	1.71	0.88
1:A:98:MET:HB2	4:A:2038:HOH:O	1.75	0.84
1:D:6:ARG:NH2	1:D:113:TYR:CD1	2.51	0.79
4:A:2041:HOH:O	2:B:0:MET:HB2	1.84	0.77
1:A:67:VAL:HG22	4:A:2021:HOH:O	1.83	0.77
2:E:48:LYS:O	2:E:48:LYS:HG3	1.84	0.76
1:D:11:SER:HB2	1:D:74:HIS:CD2	2.21	0.74
1:A:165:VAL:HG12	4:A:2039:HOH:O	1.89	0.71
2:E:97:ARG:HB2	4:E:2038:HOH:O	1.90	0.71
1:A:22:PHE:HE2	4:A:2021:HOH:O	1.75	0.70

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:252:GLY:O	4:D:2067:HOH:O	2.09	0.69
1:A:21:ARG:HB2	4:A:2001:HOH:O	1.94	0.68
1:A:68:LYS:HG3	4:A:2018:HOH:O	1.95	0.67
1:A:6:ARG:HH11	1:A:6:ARG:CG	2.07	0.66
1:A:263:HIS:CD2	1:A:265:GLY:H	2.14	0.66
1:A:74:HIS:HE1	1:A:97:ARG:HD2	1.60	0.66
1:D:263:HIS:CD2	1:D:265:GLY:H	2.14	0.66
1:D:19:GLU:HG3	1:D:20:PRO:HD2	1.78	0.65
2:E:44:GLU:HB2	4:E:2020:HOH:O	1.96	0.65
1:D:81:LEU:O	4:D:2028:HOH:O	2.14	0.65
1:D:106:ASP:O	1:D:107:TRP:HB2	2.01	0.60
1:D:22:PHE:HE2	4:D:2022:HOH:O	1.86	0.59
1:D:122:ASP:OD1	4:D:2041:HOH:O	2.16	0.58
1:D:255:GLN:HG3	4:D:2067:HOH:O	2.03	0.58
1:A:5:MET:HB2	1:A:168:LEU:HD13	1.84	0.58
1:A:48:ARG:HG2	4:A:2012:HOH:O	2.04	0.57
1:D:97:ARG:CZ	3:F:6:ILE:HD11	2.34	0.57
1:D:212:GLU:OE2	1:D:212:GLU:HA	2.05	0.57
1:A:82:ARG:NH2	1:A:89:GLU:HG3	2.10	0.57
1:D:77:ASP:OD2	1:D:116:TYR:OH	2.21	0.56
1:A:88:SER:HB3	4:A:2034:HOH:O	2.04	0.56
1:A:77:ASP:OD2	1:A:116:TYR:OH	2.23	0.56
1:D:48:ARG:HG2	4:D:2015:HOH:O	2.05	0.55
1:D:133:TRP:HB2	1:D:144:LYS:HG3	1.87	0.55
1:D:263:HIS:HD2	1:D:265:GLY:H	1.55	0.55
1:D:95:VAL:HG22	1:D:118:TYR:HD1	1.72	0.54
1:A:263:HIS:HD2	1:A:265:GLY:H	1.57	0.53
1:A:97:ARG:HH21	1:A:114:HIS:HE1	1.55	0.53
1:A:19:GLU:OE1	1:A:75:ARG:HD2	2.09	0.53
1:D:28:VAL:HG23	1:D:33:PHE:CE1	2.44	0.53
1:A:6:ARG:HD2	1:A:98:MET:SD	2.49	0.53
1:A:139:ALA:HB2	4:A:2048:HOH:O	2.09	0.53
1:A:73:THR:HG21	3:C:6:ILE:HD11	1.90	0.53
2:E:74:GLU:HB3	4:E:2027:HOH:O	2.09	0.52
1:A:87:GLN:HG2	4:A:2028:HOH:O	2.10	0.52
1:A:1:GLY:O	1:A:3:HIS:NE2	2.43	0.52
1:D:64:THR:O	1:D:67:VAL:HG12	2.10	0.51
1:A:111:ARG:HD3	4:A:2040:HOH:O	2.10	0.50
4:A:2038:HOH:O	2:B:56:PHE:HE1	1.94	0.49
1:A:6:ARG:NH1	1:A:6:ARG:HG3	2.12	0.49
1:A:93:HIS:HE1	4:A:2033:HOH:O	1.95	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:21:ARG:HD3	1:D:23:ILE:HD11	1.94	0.49
1:D:5:MET:HB2	1:D:168:LEU:HD13	1.94	0.49
1:A:93:HIS:HD2	1:A:119:ASP:OD2	1.96	0.49
1:A:24:ALA:O	1:A:35:ARG:HA	2.14	0.48
1:D:26:GLY:HA2	4:D:2007:HOH:O	2.14	0.48
1:A:1:GLY:O	1:A:3:HIS:CD2	2.67	0.47
1:A:43:GLN:N	4:A:2009:HOH:O	2.30	0.47
1:D:93:HIS:HE1	4:D:2032:HOH:O	1.97	0.47
1:D:26:GLY:HA3	4:D:2002:HOH:O	2.15	0.46
2:E:36:GLU:OE1	4:E:2015:HOH:O	2.20	0.46
1:D:139:ALA:HB2	4:D:2044:HOH:O	2.15	0.46
1:D:97:ARG:HH21	1:D:114:HIS:HE1	1.63	0.45
1:D:111:ARG:CG	1:D:112:GLY:N	2.79	0.45
3:F:7:ALA:HA	4:F:2002:HOH:O	2.16	0.45
1:D:97:ARG:HE	1:D:114:HIS:CE1	2.34	0.44
1:D:14:ARG:HB3	1:D:17:ARG:HB3	1.99	0.44
1:D:27:TYR:HD1	4:D:2007:HOH:O	1.99	0.44
1:D:111:ARG:HB3	4:D:2037:HOH:O	2.17	0.44
1:A:133:TRP:HB2	1:A:144:LYS:HG3	2.00	0.44
1:D:13:SER:HB3	1:D:78:LEU:HD22	2.00	0.44
3:C:8:VAL:HG23	4:C:2003:HOH:O	2.16	0.44
2:E:51:HIS:HA	2:E:65:LEU:O	2.18	0.44
1:A:206:LEU:HD11	4:B:2003:HOH:O	2.18	0.43
1:A:117:ALA:HB2	2:B:60:TRP:CE2	2.54	0.43
2:E:22:PHE:CZ	2:E:69:GLU:HG2	2.53	0.43
2:E:24:ASN:HB3	2:E:65:LEU:HD11	2.01	0.43
1:D:187:THR:HA	1:D:204:TRP:O	2.19	0.43
1:A:151:HIS:HD2	4:A:2051:HOH:O	2.01	0.42
2:E:48:LYS:HB3	4:E:2023:HOH:O	2.20	0.42
2:B:45:ARG:HG2	2:B:45:ARG:HH11	1.85	0.42
4:A:2017:HOH:O	3:C:2:LEU:HB2	2.20	0.42
1:A:253:GLN:NE2	1:A:256:ARG:HH11	2.18	0.42
2:E:7:ILE:HD12	2:E:91:LYS:HD3	2.02	0.42
1:A:111:ARG:HG2	1:A:112:GLY:N	2.35	0.41
1:D:113:TYR:CD1	1:D:113:TYR:N	2.89	0.41
1:D:191:HIS:CE1	1:D:199:ALA:HB1	2.55	0.41
1:D:201:LEU:HD12	1:D:249:VAL:HG11	2.02	0.41
1:D:125:ALA:HB2	4:D:2041:HOH:O	2.20	0.41
1:D:11:SER:HB3	1:D:78:LEU:HD11	2.03	0.41
1:D:189:MET:HE3	1:D:272:LEU:HB3	2.03	0.41
1:A:187:THR:HA	1:A:204:TRP:O	2.21	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:44:ARG:O	4:A:2010:HOH:O	2.22	0.40
1:A:106:ASP:O	1:A:107:TRP:HB2	2.21	0.40
1:A:123:TYR:HD2	1:A:124:ILE:HG22	1.86	0.40
1:D:88:SER:HB3	4:D:2033:HOH:O	2.19	0.40
1:D:95:VAL:HG22	1:D:118:TYR:CD1	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	273/276 (99%)	262 (96%)	11 (4%)	0	100	100
1	D	273/276 (99%)	257 (94%)	16 (6%)	0	100	100
2	B	98/100 (98%)	96 (98%)	2 (2%)	0	100	100
2	E	98/100 (98%)	94 (96%)	3 (3%)	1 (1%)	17	23
3	C	7/9 (78%)	6 (86%)	1 (14%)	0	100	100
3	F	7/9 (78%)	4 (57%)	3 (43%)	0	100	100
All	All	756/770 (98%)	719 (95%)	36 (5%)	1 (0%)	53	69

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	E	74	GLU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	231/232 (100%)	218 (94%)	13 (6%)	23	34
1	D	231/232 (100%)	214 (93%)	17 (7%)	15	21
2	B	95/95 (100%)	90 (95%)	5 (5%)	25	37
2	E	95/95 (100%)	89 (94%)	6 (6%)	20	28
3	C	8/8 (100%)	6 (75%)	2 (25%)	0	0
3	F	8/8 (100%)	7 (88%)	1 (12%)	5	5
All	All	668/670 (100%)	624 (93%)	44 (7%)	18	26

All (44) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	4	SER
1	A	6	ARG
1	A	34	VAL
1	A	35	ARG
1	A	67	VAL
1	A	87	GLN
1	A	195	SER
1	A	196	ASP
1	A	216	THR
1	A	225	THR
1	A	226	GLN
1	A	255	GLN
1	A	275	GLU
2	B	0	MET
2	B	9	VAL
2	B	44	GLU
2	B	47	GLU
2	B	70	PHE
3	C	1	SER
3	C	6	ILE
1	D	11	SER
1	D	21	ARG
1	D	34	VAL
1	D	35	ARG
1	D	45	MET
1	D	74	HIS
1	D	88	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	D	102	ASP
1	D	113	TYR
1	D	116	TYR
1	D	122	ASP
1	D	181	ARG
1	D	186	LYS
1	D	225	THR
1	D	226	GLN
1	D	255	GLN
1	D	268	LYS
2	E	0	MET
2	E	9	VAL
2	E	48	LYS
2	E	70	PHE
2	E	74	GLU
2	E	75	LYS
3	F	4	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (14) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	74	HIS
1	A	93	HIS
1	A	114	HIS
1	A	141	GLN
1	A	174	ASN
1	A	188	HIS
1	A	253	GLN
1	A	263	HIS
1	D	32	GLN
1	D	74	HIS
1	D	93	HIS
1	D	114	HIS
1	D	253	GLN
1	D	263	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å ²)	Q<0.9
1	A	275/276 (99%)	0.39	9 (3%)	46	49	42, 56, 61, 63	0
1	D	275/276 (99%)	0.46	9 (3%)	46	49	45, 55, 61, 64	0
2	B	100/100 (100%)	0.00	1 (1%)	82	83	47, 54, 61, 67	0
2	E	100/100 (100%)	0.08	1 (1%)	82	83	48, 54, 62, 67	0
3	C	9/9 (100%)	1.47	3 (33%)	0	0	65, 72, 75, 76	0
3	F	9/9 (100%)	1.26	2 (22%)	0	0	61, 73, 74, 74	0
All	All	768/770 (99%)	0.35	25 (3%)	46	49	42, 55, 62, 76	0

All (25) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	F	5	THR	5.6
3	C	5	THR	3.8
1	A	81	LEU	3.2
1	D	90	ALA	3.0
1	D	111	ARG	2.9
1	A	192	HIS	2.8
1	D	113	TYR	2.7
1	A	172	LEU	2.7
1	D	166	GLU	2.6
1	D	85	TYR	2.5
3	F	4	ASN	2.5
1	A	76	VAL	2.4
1	D	69	ALA	2.4
2	B	0	MET	2.4
1	D	118	TYR	2.4
1	A	85	TYR	2.3
1	D	192	HIS	2.3
1	A	180	GLN	2.3
3	C	7	ALA	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	266	LEU	2.2
2	E	0	MET	2.2
1	A	113	TYR	2.2
1	A	108	ARG	2.2
1	D	123	TYR	2.1
3	C	3	PHE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.