



Full wwPDB X-ray Structure Validation Report ⓘ

Nov 14, 2019 – 05:49 AM EST

PDB ID : 6C8M
Title : RNA-activated 2-AIpG monomer, 1.5h soaking
Authors : Zhang, W.; Szostak, J.W.
Deposited on : 2018-01-25
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.0 (224370), CSD as540be (2019)
Xtriage (Phenix) : 1.13
EDS : 2.4
buster-report : 1.1.7 (2018)
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)
Refmac : 5.8.0158
CCP4 : 7.0 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.4


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	111664	3481 (2.40-2.40)
Clashscore	122126	3956 (2.40-2.40)
RSRZ outliers	108989	3386 (2.40-2.40)
RNA backbone	2636	1023 (2.80-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	15	
2	B	14	

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 760 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a RNA chain called RNA (5'-R*(LCC)P*(LCC)P*(LCC)P*(LCG)P*AP*CP*UP*UP*AP*AP*GP*UP*CP*GP*G)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	15	Total	C	N	O	P	0	1	0
			322	149	56	103	14			


- Molecule 2 is a RNA chain called RNA (5'-R*(LCC)P*(LCC)P*(LCC)P*(LCG)P*AP*CP*UP*UP*AP*AP*GP*UP*CP*G)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	14	Total	C	N	O	P	0	0	0
			299	139	51	96	13			

- Molecule 3 is 5'-O-[(R)-(2-amino-1H-imidazol-1-yl)(hydroxy)phosphoryl]guanosine (three-letter code: EQ4) (formula: C₁₃H₁₇N₈O₇P).



- Molecule 4 is [2-azanyl-3-[[[(2 {R},3 {S},4 {R},5 {R})-5-(2-azanyl-6-oxidanylidene-1 {H}-purin-9-yl)-3,4-bis(oxidanyl)oxolan-2-yl]methoxy-oxidanyl-phosphoryl]imidazol-1-yl]-[[[(2 {R},3 {S},4 {R},5 {R})-5-(2-azanyl-6-oxidanylidene-1 {H}-purin-9-yl)-3,4-bis(oxidanyl)oxolan-2-yl]methoxy]phosphinic acid (three-letter code: EQ1) (formula: C₂₃H₃₀N₁₃O₁₄P₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
4	A	1	52	23	13	14	2	0	1
4	A	1	52	23	13	14	2	0	0

- Molecule 5 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	4	Total 4 O 4	0	0
5	B	2	Total 2 O 2	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: RNA (5'-R*(LCC)P*(LCC)P*(LCC)P*(LCG)P*AP*CP*UP*UP*AP*AP*GP*UP*CP*GP*G)-3')

Chain A: 



- Molecule 2: RNA (5'-R*(LCC)P*(LCC)P*(LCC)P*(LCG)P*AP*CP*UP*UP*AP*AP*GP*UP*CP*G)-3')

Chain B: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 3 2 1	Depositor
Cell constants a, b, c, α , β , γ	47.87Å 47.87Å 82.73Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	50.00 – 2.40 22.99 – 2.40	Depositor EDS
% Data completeness (in resolution range)	91.8 (50.00-2.40) 91.9 (22.99-2.40)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.35 (at 2.41Å)	Xtriage
Refinement program	REFMAC 5.8.0135	Depositor
R, R_{free}	0.240 , 0.272 0.251 , 0.277	Depositor DCC
R_{free} test set	219 reflections (5.18%)	wwPDB-VP
Wilson B-factor (Å ²)	65.3	Xtriage
Anisotropy	0.005	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.23 , 27.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.53$, $\langle L^2 \rangle = 0.38$	Xtriage
Estimated twinning fraction	0.450 for -h,-k,l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	760	wwPDB-VP
Average B, all atoms (Å ²)	67.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 10.10% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: EQ1, LCG, EQ4, LCC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.52	0/262	0.86	0/406
2	B	0.59	1/236 (0.4%)	0.86	0/365
All	All	0.56	1/498 (0.2%)	0.86	0/771

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	13	C	O3'-P	-5.76	1.54	1.61

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	322	0	169	8	0
2	B	299	0	158	2	0
3	A	29	0	0	6	0
4	A	104	0	0	0	0
5	A	4	0	0	0	0
5	B	2	0	0	0	0
All	All	760	0	327	11	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (11) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:15[B]:G:H3'	3:A:101[B]:EQ4:C1	1.97	0.94
1:A:15[B]:G:H2'	3:A:101[B]:EQ4:O4	1.82	0.77
3:A:101[B]:EQ4:C10	3:A:101[B]:EQ4:C1	2.63	0.76
1:A:15[B]:G:C3'	3:A:101[B]:EQ4:C1	2.74	0.62
2:B:10:A:H2'	2:B:11:G:O4'	2.03	0.59
1:A:7:U:H2'	1:A:8:U:C6	2.39	0.57
2:B:7:U:H2'	2:B:8:U:C6	2.39	0.57
1:A:15[B]:G:O3'	3:A:101[B]:EQ4:C2	2.58	0.52
1:A:15[B]:G:C3'	3:A:101[B]:EQ4:C2	2.96	0.42
1:A:12:U:H2'	1:A:13:C:O4'	2.19	0.42
1:A:2:LCC:H41	1:A:2:LCC:H5M1	1.60	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

There are no protein molecules in this entry.

5.3.2 Protein sidechains [i](#)

There are no protein molecules in this entry.

5.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
1	A	11/15 (73%)	2 (18%)	1 (9%)
2	B	10/14 (71%)	2 (20%)	1 (10%)
All	All	21/29 (72%)	4 (19%)	2 (9%)

All (4) RNA backbone outliers are listed below:

Mol	Chain	Res	Type
1	A	8	U
1	A	10	A
2	B	9	A
2	B	14	G

All (2) RNA pucker outliers are listed below:

Mol	Chain	Res	Type
1	A	3	LCC
2	B	3	LCC

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

8 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
1	LCC	A	1	1	16,21,25	2.55	7 (43%)	24,33,40	2.49	7 (29%)
1	LCC	A	2	1	16,24,25	2.85	6 (37%)	22,37,40	1.93	6 (27%)
1	LCC	A	3	1	16,24,25	2.96	6 (37%)	22,37,40	2.23	8 (36%)
1	LCG	A	4	1,2	19,27,28	3.18	9 (47%)	24,42,45	3.07	13 (54%)
2	LCC	B	1	2	16,21,25	3.01	5 (31%)	24,33,40	2.37	11 (45%)
2	LCC	B	2	2	16,24,25	2.67	7 (43%)	22,37,40	2.21	8 (36%)
2	LCC	B	3	2	16,24,25	2.40	8 (50%)	22,37,40	2.25	7 (31%)
2	LCG	B	4	1,2	19,27,28	3.14	8 (42%)	24,42,45	3.00	12 (50%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	LCC	A	1	1	-	0/3/32/36	0/4/3/3

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	LCC	A	2	1	-	0/4/35/36	0/4/3/3
1	LCC	A	3	1	-	1/4/35/36	0/4/3/3
1	LCG	A	4	1,2	-	0/4/35/36	0/5/4/4
2	LCC	B	1	2	-	0/3/32/36	0/4/3/3
2	LCC	B	2	2	-	0/4/35/36	0/4/3/3
2	LCC	B	3	2	-	0/4/35/36	0/4/3/3
2	LCG	B	4	1,2	-	0/4/35/36	0/5/4/4

All (56) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1	LCC	C4'-C3'	-9.31	1.42	1.53
1	A	3	LCC	C4'-C3'	-8.26	1.44	1.53
1	A	4	LCG	O2'-C6'	7.34	1.57	1.43
2	B	4	LCG	C4'-C3'	-7.05	1.45	1.53
2	B	4	LCG	O2'-C6'	6.93	1.57	1.43
1	A	4	LCG	C6'-C4'	5.95	1.64	1.53
1	A	2	LCC	C4'-C3'	-5.89	1.46	1.53
2	B	4	LCG	C6'-C4'	5.83	1.64	1.53
1	A	3	LCC	C5M-C5	-5.52	1.39	1.51
1	A	2	LCC	C5M-C5	-5.42	1.40	1.51
1	A	1	LCC	C4'-C3'	-5.21	1.47	1.53
1	A	4	LCG	C6-C5	-5.12	1.32	1.41
2	B	4	LCG	C6-C5	-4.68	1.33	1.41
2	B	2	LCC	C5M-C5	-4.65	1.41	1.51
1	A	1	LCC	O2'-C6'	4.64	1.52	1.43
2	B	2	LCC	O2'-C6'	4.60	1.52	1.43
1	A	4	LCG	C6-N1	4.50	1.40	1.33
2	B	3	LCC	C5M-C5	-4.46	1.42	1.51
2	B	2	LCC	C2'-C3'	-4.35	1.43	1.53
1	A	1	LCC	C5M-C5	-4.29	1.42	1.51
1	A	2	LCC	O2'-C6'	4.20	1.51	1.43
1	A	2	LCC	O4'-C4'	-4.15	1.40	1.45
2	B	3	LCC	C4'-C3'	-4.08	1.49	1.53
1	A	4	LCG	C2-N1	4.07	1.42	1.35
2	B	2	LCC	C4'-C3'	-3.98	1.49	1.53
2	B	1	LCC	O2'-C6'	3.95	1.51	1.43
2	B	3	LCC	O2'-C6'	3.86	1.51	1.43
1	A	2	LCC	C2'-C3'	-3.77	1.44	1.53
2	B	2	LCC	O3'-C3'	-3.59	1.35	1.42
1	A	3	LCC	C2'-C3'	-3.53	1.45	1.53
1	A	4	LCG	C4'-C3'	-3.46	1.49	1.53

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1	LCC	C5M-C5	-3.41	1.44	1.51
2	B	1	LCC	C6'-C4'	3.08	1.58	1.53
1	A	3	LCC	C5'-C4'	2.95	1.56	1.51
2	B	4	LCG	C6-N1	2.91	1.38	1.33
1	A	1	LCC	O4'-C4'	2.76	1.49	1.45
1	A	1	LCC	O2'-C2'	2.74	1.50	1.43
2	B	4	LCG	C5-C4	-2.74	1.34	1.40
2	B	1	LCC	C2'-C3'	-2.73	1.47	1.53
2	B	3	LCC	O4'-C4'	-2.68	1.42	1.45
2	B	4	LCG	O3'-C3'	2.68	1.48	1.42
2	B	3	LCC	C2'-C3'	-2.66	1.47	1.53
1	A	4	LCG	C5'-C4'	2.57	1.55	1.51
2	B	2	LCC	C2-N3	-2.53	1.33	1.38
1	A	1	LCC	C6'-C4'	2.48	1.57	1.53
2	B	3	LCC	C6'-C4'	2.47	1.57	1.53
2	B	3	LCC	O3'-C3'	-2.44	1.37	1.42
2	B	3	LCC	C5'-C4'	2.41	1.55	1.51
1	A	2	LCC	C2-N3	-2.41	1.33	1.38
2	B	2	LCC	O4'-C4'	-2.38	1.42	1.45
1	A	1	LCC	O3'-C3'	-2.33	1.38	1.42
1	A	3	LCC	O2'-C6'	2.32	1.48	1.43
1	A	4	LCG	C5-C4	-2.29	1.35	1.40
1	A	3	LCC	O4'-C1'	-2.28	1.39	1.42
2	B	4	LCG	C2-N1	2.26	1.39	1.35
1	A	4	LCG	O4'-C4'	-2.07	1.42	1.45

All (72) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	4	LCG	N3-C2-N1	-7.54	117.05	127.25
2	B	4	LCG	N3-C2-N1	-7.24	117.45	127.25
1	A	1	LCC	O4'-C4'-C5'	6.27	120.41	109.00
2	B	4	LCG	O4'-C1'-N9	-6.24	100.47	108.71
1	A	4	LCG	O4'-C1'-N9	-6.11	100.65	108.71
1	A	1	LCC	O3'-C3'-C2'	-5.86	94.16	112.65
1	A	3	LCC	O4'-C1'-N1	-5.40	101.58	108.71
2	B	2	LCC	C2-N3-C4	5.28	121.95	115.92
1	A	2	LCC	C5-C6-N1	-4.79	116.96	122.15
1	A	4	LCG	C1'-N9-C4	-4.70	118.52	126.64
2	B	3	LCC	O4'-C4'-C3'	-4.58	96.65	102.13
2	B	3	LCC	C5-C6-N1	-4.33	117.46	122.15
1	A	1	LCC	C2-N3-C4	4.29	120.82	115.92

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1	LCC	C2-N3-C4	4.28	120.80	115.92
1	A	1	LCC	O3'-C3'-C4'	4.27	123.35	112.82
2	B	1	LCC	O4'-C4'-C5'	4.25	116.73	109.00
2	B	1	LCC	O3'-C3'-C4'	4.23	123.27	112.82
2	B	2	LCC	O4'-C1'-N1	4.15	114.18	108.71
2	B	4	LCG	C6'-O2'-C2'	-4.10	100.14	107.69
2	B	1	LCC	O3'-C3'-C2'	-3.92	100.27	112.65
1	A	3	LCC	C5M-C5-C4	3.91	125.58	121.68
2	B	3	LCC	O4'-C1'-N1	-3.90	103.56	108.71
2	B	4	LCG	C2-N3-C4	3.86	119.77	115.36
1	A	2	LCC	O4'-C4'-C6'	3.80	114.18	105.97
1	A	4	LCG	C6-N1-C2	3.75	121.39	116.06
2	B	3	LCC	C5M-C5-C4	3.68	125.36	121.68
2	B	1	LCC	O5'-C5'-C4'	-3.57	104.44	111.73
2	B	4	LCG	C6-N1-C2	3.48	121.02	116.06
1	A	3	LCC	C5-C6-N1	-3.48	118.38	122.15
1	A	3	LCC	O4'-C4'-C5'	3.45	119.19	108.75
1	A	4	LCG	C6'-O2'-C2'	-3.43	101.38	107.69
2	B	4	LCG	N2-C2-N3	3.41	123.39	117.77
2	B	4	LCG	O4'-C4'-C3'	3.41	106.20	102.13
2	B	4	LCG	C1'-N9-C4	-3.31	120.92	126.64
1	A	4	LCG	O3'-C3'-C4'	-3.30	104.66	112.82
1	A	3	LCC	O2'-C6'-C4'	-3.30	97.37	103.81
1	A	4	LCG	C5-C6-N1	-3.28	118.90	123.47
2	B	3	LCC	O4'-C4'-C5'	3.26	118.63	108.75
1	A	4	LCG	O2'-C2'-C1'	3.17	115.51	105.63
1	A	3	LCC	N4-C4-N3	3.13	121.49	117.03
2	B	4	LCG	O3'-C3'-C4'	-3.12	105.11	112.82
2	B	2	LCC	C5-C6-N1	-3.11	118.78	122.15
1	A	2	LCC	C4'-O4'-C1'	-3.07	102.15	107.61
2	B	2	LCC	C4'-O4'-C1'	-3.05	102.19	107.61
2	B	4	LCG	C4'-O4'-C1'	-3.03	102.23	107.61
1	A	4	LCG	O4'-C4'-C3'	3.02	105.75	102.13
2	B	1	LCC	O4'-C1'-N1	-3.02	104.73	108.71
1	A	4	LCG	C2-N3-C4	3.00	118.78	115.36
1	A	4	LCG	N2-C2-N1	2.95	121.86	117.25
2	B	2	LCC	O2'-C6'-C4'	-2.92	98.11	103.81
2	B	4	LCG	C5-C6-N1	-2.89	119.45	123.47
1	A	1	LCC	O2'-C6'-C4'	-2.86	98.22	103.81
2	B	1	LCC	O2'-C6'-C4'	-2.78	98.38	103.81
1	A	1	LCC	C5M-C5-C4	2.77	124.44	121.68
1	A	3	LCC	O2'-C2'-C3'	-2.72	98.07	103.42

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	2	LCC	O4'-C4'-C6'	2.69	111.80	105.97
1	A	4	LCG	C4'-O4'-C1'	-2.66	102.89	107.61
1	A	2	LCC	C2-N3-C4	2.64	118.93	115.92
2	B	1	LCC	C5-C6-N1	-2.58	119.36	122.15
2	B	1	LCC	O2'-C2'-C3'	-2.53	98.44	103.42
2	B	4	LCG	O2'-C2'-C1'	2.50	113.42	105.63
2	B	1	LCC	C5M-C5-C4	2.45	124.12	121.68
2	B	3	LCC	O4'-C4'-C6'	2.41	111.17	105.97
2	B	3	LCC	O2'-C6'-C4'	-2.33	99.26	103.81
2	B	2	LCC	O2'-C2'-C1'	2.29	112.77	105.63
2	B	1	LCC	C5'-C4'-C3'	-2.25	111.25	117.07
2	B	2	LCC	O4'-C4'-C5'	2.20	115.42	108.75
1	A	2	LCC	N4-C4-N3	2.17	120.13	117.03
1	A	1	LCC	C5'-C4'-C3'	-2.05	111.78	117.07
1	A	2	LCC	C6'-O2'-C2'	-2.02	103.98	107.69
1	A	4	LCG	N2-C2-N3	2.01	121.08	117.77
1	A	3	LCC	C4'-O4'-C1'	-2.00	104.05	107.61

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	3	LCC	O4'-C4'-C5'-O5'

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	2	LCC	1	0

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	EQ4	A	101[B]	4	26,32,32	4.08	9 (34%)	23,49,49	2.70	4 (17%)
4	EQ1	A	102[A]	1,3	46,58,58	3.68	20 (43%)	52,91,91	2.72	20 (38%)
4	EQ1	A	103	-	46,58,58	3.45	16 (34%)	52,91,91	2.38	16 (30%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	EQ4	A	101[B]	4	-	3/5/32/32	0/4/4/4
4	EQ1	A	102[A]	1,3	-	5/12/77/77	0/7/7/7
4	EQ1	A	103	-	-	4/12/77/77	0/7/7/7

All (45) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	101[B]	EQ4	P1-O2	16.42	1.61	1.47
4	A	102[A]	EQ1	PG-O2G	13.07	1.62	1.46
4	A	103	EQ1	PG-O2G	12.94	1.61	1.46
4	A	102[A]	EQ1	C2C-N2C	-9.06	1.33	1.47
4	A	103	EQ1	C2C-N2C	-8.65	1.34	1.47
3	A	101[B]	EQ4	P1-O3	8.41	1.65	1.57
4	A	102[A]	EQ1	C3C-N1C	-8.39	1.34	1.47
4	A	103	EQ1	C3C-N1C	-8.11	1.35	1.47
4	A	102[A]	EQ1	C3C-C2C	-7.35	1.33	1.51
4	A	103	EQ1	C3C-C2C	-7.22	1.33	1.51
4	A	102[A]	EQ1	PA-N1C	6.63	1.76	1.63
4	A	103	EQ1	C6B-C5B	-4.70	1.33	1.41
4	A	102[A]	EQ1	C6B-N1B	4.61	1.41	1.33
4	A	102[A]	EQ1	PG-O5E	4.55	1.70	1.57
4	A	102[A]	EQ1	C6A-N1A	4.41	1.40	1.33
3	A	101[B]	EQ4	C9-C8	-4.40	1.33	1.41
4	A	103	EQ1	PA-N1C	4.31	1.72	1.63
4	A	103	EQ1	PA-O5D	4.14	1.69	1.57

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	102[A]	EQ1	PA-O5D	4.09	1.69	1.57
4	A	103	EQ1	O4E-C1E	4.01	1.46	1.41
4	A	102[A]	EQ1	C6A-C5A	-4.01	1.34	1.41
3	A	101[B]	EQ4	O4-C5	3.94	1.46	1.41
3	A	101[B]	EQ4	C9-N1	3.91	1.39	1.33
4	A	102[A]	EQ1	C6B-C5B	-3.85	1.34	1.41
4	A	103	EQ1	C6A-N1A	3.80	1.39	1.33
4	A	103	EQ1	C6A-C5A	-3.74	1.34	1.41
4	A	103	EQ1	C6B-N1B	3.72	1.39	1.33
4	A	102[A]	EQ1	PA-O1A	-3.61	1.46	1.56
4	A	102[A]	EQ1	PA-O2A	3.59	1.50	1.46
4	A	103	EQ1	PA-O1A	-3.59	1.46	1.56
4	A	103	EQ1	PG-O5E	3.40	1.67	1.57
3	A	101[B]	EQ4	P1-O1	-2.97	1.48	1.54
3	A	101[B]	EQ4	C11-N6	-2.90	1.34	1.39
4	A	102[A]	EQ1	PG-O1G	-2.74	1.49	1.56
4	A	102[A]	EQ1	C3D-C2D	-2.74	1.46	1.53
4	A	103	EQ1	PG-O1G	-2.73	1.49	1.56
4	A	102[A]	EQ1	C2B-N1B	2.67	1.40	1.35
3	A	101[B]	EQ4	C8-C7	-2.65	1.34	1.40
4	A	103	EQ1	C5A-C4A	-2.63	1.34	1.40
4	A	103	EQ1	O3E-C3E	2.58	1.49	1.43
4	A	102[A]	EQ1	C5A-C4A	-2.58	1.34	1.40
4	A	102[A]	EQ1	C2A-N1A	2.45	1.39	1.35
4	A	102[A]	EQ1	C5B-C4B	-2.41	1.35	1.40
3	A	101[B]	EQ4	C6-N1	2.39	1.39	1.35
4	A	102[A]	EQ1	O3D-C3D	2.04	1.47	1.43

All (40) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	101[B]	EQ4	O3-P1-O2	-9.37	103.14	115.31
4	A	102[A]	EQ1	O5D-PA-O2A	-6.94	95.04	115.66
4	A	102[A]	EQ1	N3B-C2B-N1B	-6.26	118.78	127.25
4	A	102[A]	EQ1	N3A-C2A-N1A	-6.11	118.99	127.25
4	A	103	EQ1	N3A-C2A-N1A	-6.06	119.05	127.25
3	A	101[B]	EQ4	N3-C6-N1	-5.60	119.67	127.25
4	A	102[A]	EQ1	C2A-N3A-C4A	5.36	121.48	115.36
4	A	103	EQ1	C2A-N3A-C4A	5.21	121.31	115.36
4	A	103	EQ1	O5D-PA-O2A	-5.07	100.59	115.66
4	A	103	EQ1	O5E-PG-O2G	-4.95	100.94	115.66
4	A	103	EQ1	O2A-PA-N1C	4.91	120.67	113.14

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	103	EQ1	N3B-C2B-N1B	-4.82	120.73	127.25
4	A	102[A]	EQ1	O5E-PG-O2G	-4.78	101.44	115.66
4	A	102[A]	EQ1	C1D-N9A-C4A	-4.69	118.53	126.64
4	A	102[A]	EQ1	C2B-N3B-C4B	4.68	120.70	115.36
3	A	101[B]	EQ4	C6-N3-C7	4.43	120.42	115.36
4	A	102[A]	EQ1	O2A-PA-N1C	4.37	119.85	113.14
4	A	102[A]	EQ1	O5D-PA-N1C	-4.09	93.71	105.90
4	A	102[A]	EQ1	O1A-PA-O2A	3.84	119.10	109.81
4	A	102[A]	EQ1	C5B-C6B-N1B	-3.44	118.68	123.47
4	A	103	EQ1	N2A-C2A-N3A	3.35	123.30	117.77
4	A	103	EQ1	C5B-C6B-N1B	-3.23	118.98	123.47
4	A	102[A]	EQ1	N2B-C2B-N1B	3.21	122.27	117.25
4	A	102[A]	EQ1	C2C-C3C-N1C	3.12	109.22	105.38
4	A	103	EQ1	PA-O5D-C5D	3.09	129.35	120.61
4	A	103	EQ1	C2B-N3B-C4B	3.01	118.79	115.36
4	A	103	EQ1	O1A-PA-O5D	-2.99	94.25	107.44
3	A	101[B]	EQ4	C8-C9-N1	-2.95	119.36	123.47
4	A	103	EQ1	O1A-PA-O2A	2.79	116.57	109.81
4	A	103	EQ1	C2C-C3C-N1C	2.78	108.80	105.38
4	A	103	EQ1	N2B-C2B-N3B	2.65	122.14	117.77
4	A	102[A]	EQ1	C4E-O4E-C1E	2.62	112.56	109.83
4	A	102[A]	EQ1	C6B-N1B-C2B	2.60	119.75	116.06
4	A	102[A]	EQ1	C5A-C6A-N1A	-2.56	119.92	123.47
4	A	103	EQ1	O2G-PG-N2C	-2.53	109.26	113.14
4	A	102[A]	EQ1	C1E-N9B-C4B	-2.42	122.45	126.64
4	A	102[A]	EQ1	O2E-C2E-C3E	-2.35	104.19	111.80
4	A	103	EQ1	C6B-N1B-C2B	2.29	119.32	116.06
4	A	102[A]	EQ1	O1A-PA-O5D	-2.23	97.63	107.44
4	A	102[A]	EQ1	C4B-C5B-N7B	-2.10	107.21	109.40

There are no chirality outliers.

All (12) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	101[B]	EQ4	C1-O3-P1-O2
4	A	102[A]	EQ1	C5E-O5E-PG-N2C
4	A	102[A]	EQ1	C5E-O5E-PG-O2G
4	A	103	EQ1	C5D-O5D-PA-N1C
4	A	103	EQ1	C5E-O5E-PG-N2C
4	A	103	EQ1	O4D-C4D-C5D-O5D
4	A	103	EQ1	C3D-C4D-C5D-O5D
4	A	102[A]	EQ1	C5E-O5E-PG-O1G

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
3	A	101[B]	EQ4	O3-C1-C2-O4
4	A	102[A]	EQ1	O4D-C4D-C5D-O5D
4	A	102[A]	EQ1	C3D-C4D-C5D-O5D
3	A	101[B]	EQ4	O3-C1-C2-C3


There are no ring outliers.

1 monomer is involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	101[B]	EQ4	6	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	11/15 (73%)	-1.18	0 100 100	50, 67, 84, 88	1 (9%)
2	B	10/14 (71%)	-1.21	0 100 100	51, 75, 89, 89	0
All	All	21/29 (72%)	-1.20	0 100 100	50, 71, 88, 89	1 (4%)

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	LCC	A	3	22/23	0.96	0.12	42,49,52,55	0
2	LCC	B	1	19/23	0.96	0.11	55,62,75,76	0
2	LCC	B	3	22/23	0.96	0.11	40,48,53,57	0
1	LCC	A	1	19/23	0.97	0.09	41,57,74,83	0
1	LCC	A	2	22/23	0.97	0.09	43,49,56,60	0
2	LCG	B	4	24/25	0.97	0.12	41,54,62,68	0
2	LCC	B	2	22/23	0.97	0.10	46,49,56,58	0
1	LCG	A	4	24/25	0.98	0.09	36,47,56,65	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands ⓘ




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	EQ1	A	103	52/52	0.93	0.13	51,74,128,142	0
4	EQ1	A	102[A]	52/52	0.95	0.12	51,68,135,150	52
3	EQ4	A	101[B]	29/29	0.96	0.14	57,66,76,78	29

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around EQ1 A 103:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around EQ1 A 102 (A):

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around EQ4 A 101 (B):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.