



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2018 – 06:47 am GMT

PDB ID : 5C9C  
Title : CRYSTAL STRUCTURE OF BRAF(V600E) IN COMPLEX WITH  
LY3009120 COMPND  
Authors : Edwards, T.; Abendroth, J.; Chun, L.  
Deposited on : 2015-06-26  
Resolution : 2.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : trunk30967  
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)  
Refmac : 5.8.0158  
CCP4 : 7.0 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk30967


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	111664	2449 (2.70-2.70)
Clashscore	122126	2756 (2.70-2.70)
Ramachandran outliers	120053	2716 (2.70-2.70)
Sidechain outliers	120020	2716 (2.70-2.70)
RSRZ outliers	108989	2376 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	307	<div> <div>2%</div> <div> <div></div> <div>70%</div> <div>12%</div> <div>•</div> <div>17%</div> </div> </div>
1	B	307	<div> <div>3%</div> <div> <div></div> <div>68%</div> <div>13%</div> <div>18%</div> </div> </div>

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 4153 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Serine/threonine-protein kinase B-raf.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	254	Total	C	N	O	S	0	1	0
			1997	1284	344	357	12			
1	B	251	Total	C	N	O	S	0	1	0
			1958	1261	334	350	13			

There are 26 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	419	MET	-	initiating methionine	UNP P15056
A	420	ASP	-	expression tag	UNP P15056
A	421	ARG	-	expression tag	UNP P15056
A	422	GLY	-	expression tag	UNP P15056
A	423	SER	-	expression tag	UNP P15056
A	424	HIS	-	expression tag	UNP P15056
A	425	HIS	-	expression tag	UNP P15056
A	426	HIS	-	expression tag	UNP P15056
A	427	HIS	-	expression tag	UNP P15056
A	428	HIS	-	expression tag	UNP P15056
A	429	HIS	-	expression tag	UNP P15056
A	430	GLY	-	expression tag	UNP P15056
A	599	GLU	VAL	engineered mutation	UNP P15056
B	419	MET	-	initiating methionine	UNP P15056
B	420	ASP	-	expression tag	UNP P15056
B	421	ARG	-	expression tag	UNP P15056
B	422	GLY	-	expression tag	UNP P15056
B	423	SER	-	expression tag	UNP P15056
B	424	HIS	-	expression tag	UNP P15056
B	425	HIS	-	expression tag	UNP P15056
B	426	HIS	-	expression tag	UNP P15056
B	427	HIS	-	expression tag	UNP P15056
B	428	HIS	-	expression tag	UNP P15056
B	429	HIS	-	expression tag	UNP P15056
B	430	GLY	-	expression tag	UNP P15056

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	599	GLU	VAL	engineered mutation	UNP P15056

- Molecule 2 is 1-(3,3-dimethylbutyl)-3-{2-fluoro-4-methyl-5-[7-methyl-2-(methylamino)pyridin-2-yl]phenyl}urea (three-letter code: 4Z5) (formula: C<sub>23</sub>H<sub>29</sub>FN<sub>6</sub>O).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	F	N	O	0	0
			31	23	1	6	1		
2	B	1	Total	C	F	N	O	0	0
			31	23	1	6	1		

- Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Cl	0	0
			1	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	80	Total	O	0	0
			80	80		
4	B	55	Total	O	0	0
			55	55		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Serine/threonine-protein kinase B-raf



- Molecule 1: Serine/threonine-protein kinase B-raf



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	93.88Å 93.88Å 165.55Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 2.70 19.62 – 2.70	Depositor EDS
% Data completeness (in resolution range)	98.9 (50.00-2.70) 98.9 (19.62-2.70)	Depositor EDS
$R_{merge}$	0.13	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.00 (at 2.71Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, $R_{free}$	0.194 , 0.238 0.195 , 0.198	Depositor DCC
$R_{free}$ test set	1065 reflections (5.14%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	33.6	Xtriage
Anisotropy	0.115	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 44.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.51$ , $\langle L^2 \rangle = 0.35$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	4153	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 33.86 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 7.4744e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 4Z5, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	1.05	2/2043 (0.1%)	0.79	0/2764
1	B	0.76	0/2003	0.76	1/2712 (0.0%)
All	All	0.92	2/4046 (0.0%)	0.78	1/5476 (0.0%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	632	TYR	CD2-CE2	-5.44	1.31	1.39
1	A	518	TYR	CD2-CE2	-5.34	1.31	1.39

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	494	LEU	CA-CB-CG	5.68	128.36	115.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1997	0	1990	20	0
1	B	1958	0	1946	24	0
2	A	31	0	29	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	31	0	29	0	0
3	B	1	0	0	1	0
4	A	80	0	0	0	0
4	B	55	0	0	2	0
All	All	4153	0	3994	43	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (43) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:505:ARG:NH2	3:B:802:CL:CL	2.29	1.03
1:B:464:SER:HB3	1:B:469:THR:HG23	1.54	0.90
1:B:530:TRP:HE1	1:B:532:GLU:HG2	1.55	0.71
1:A:585:GLU:HA	1:A:585:GLU:OE1	1.95	0.66
1:A:627:GLN:HG2	1:A:627:GLN:O	1.97	0.65
1:B:675:PRO:HG2	1:B:696:LEU:HD13	1.77	0.64
1:A:554:ASP:OD1	1:A:557:ARG:NH2	2.34	0.61
1:B:670:ARG:NH1	4:B:901:HOH:O	2.35	0.59
1:B:530:TRP:NE1	1:B:532:GLU:HG2	2.18	0.58
1:A:581:ILE:HG23	1:A:589:VAL:HG13	1.87	0.55
1:A:484:LEU:O	1:A:485:ASN:HB3	2.07	0.54
1:A:585:GLU:O	1:A:586:ASP:HB2	2.08	0.53
1:A:573:HIS:O	1:A:574:ARG:HB2	2.06	0.53
1:A:452:PRO:HB2	1:A:455:GLN:NE2	2.25	0.51
1:B:674:SER:HB2	1:B:675:PRO:CD	2.41	0.50
1:A:508:ARG:HD2	1:B:449:TRP:CZ3	2.46	0.50
1:B:715:LEU:O	1:B:715:LEU:HD12	2.12	0.49
1:B:501:VAL:O	1:B:505:ARG:HG2	2.12	0.48
1:A:584:HIS:O	1:A:585:GLU:HB2	2.13	0.48
1:B:580:ASN:HA	1:B:594:PHE:HD1	1.80	0.46
1:B:532:GLU:H	1:B:584:HIS:CE1	2.33	0.46
1:B:583:LEU:HA	1:B:583:LEU:HD23	1.75	0.45
1:A:531:CYS:HB3	1:A:582:PHE:CE2	2.52	0.45
1:A:619:MET:HG2	1:A:623:VAL:CG1	2.47	0.45
1:A:547:PHE:HB2	1:A:552:LEU:HD21	1.99	0.44
1:B:554:ASP:OD1	1:B:557:ARG:NH1	2.41	0.44
1:B:452:PRO:HG2	1:B:455:GLN:NE2	2.33	0.44
1:A:580:ASN:HA	1:A:594:PHE:HD1	1.83	0.44
1:A:574:ARG:HG2	1:A:632:TYR:CD2	2.53	0.43

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:520:THR:O	1:A:520:THR:HG23	2.18	0.43
1:A:629:LYS:HA	1:A:630:ASN:C	2.38	0.43
1:B:452:PRO:HG2	1:B:455:GLN:HE22	1.83	0.43
1:B:457:THR:HB	1:B:474:LYS:HB2	2.00	0.43
1:A:582:PHE:O	1:A:589:VAL:HA	2.19	0.43
1:B:663:GLN:O	1:B:667:MET:CG	2.68	0.41
1:A:482:LYS:O	1:A:525:ALA:HA	2.20	0.41
1:A:547:PHE:HB2	1:A:552:LEU:CD2	2.50	0.41
1:B:488:ALA:HA	1:B:489:PRO:HD2	1.93	0.41
1:B:674:SER:HB2	1:B:675:PRO:HD2	2.01	0.41
1:B:708:GLN:HB2	4:B:934:HOH:O	2.21	0.41
1:B:499:ASN:O	1:B:503:VAL:HG23	2.22	0.40
1:B:510:VAL:O	1:B:590:LYS:NZ	2.39	0.40
1:B:616:ILE:HA	1:B:619:MET:HG3	2.02	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	249/307 (81%)	240 (96%)	9 (4%)	0	100	100
1	B	246/307 (80%)	237 (96%)	9 (4%)	0	100	100
All	All	495/614 (81%)	477 (96%)	18 (4%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	213/271 (79%)	201 (94%)	12 (6%)	23	49
1	B	208/271 (77%)	200 (96%)	8 (4%)	36	66
All	All	421/542 (78%)	401 (95%)	20 (5%)	30	57

All (20) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	448	ASP
1	A	466	SER
1	A	494	LEU
1	A	520	THR
1	A	521	LYS
1	A	534	SER
1	A	577	LYS
1	A	587	LEU
1	A	615	SER
1	A	708[A]	GLN
1	A	708[B]	GLN
1	A	716	LEU
1	B	448	ASP
1	B	487	THR
1	B	520	THR
1	B	523	GLN
1	B	544	GLU
1	B	596	LEU
1	B	708	GLN
1	B	716	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	495	GLN
1	A	499	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 1 is monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
2	4Z5	A	900	-	33,33,33	1.31	3 (9%)	45,48,48	2.16	16 (35%)
2	4Z5	B	801	-	33,33,33	1.34	4 (12%)	45,48,48	1.97	13 (28%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	4Z5	A	900	-	-	0/17/17/17	0/3/3/3
2	4Z5	B	801	-	-	0/17/17/17	0/3/3/3

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	900	4Z5	C16-N22	-2.91	1.36	1.41
2	B	801	4Z5	C8-C9	-2.53	1.39	1.41
2	B	801	4Z5	C16-N22	-2.33	1.37	1.41
2	A	900	4Z5	C3-N13	2.46	1.36	1.34
2	B	801	4Z5	C3-N13	2.85	1.36	1.34
2	B	801	4Z5	C9-N10	4.37	1.36	1.32

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	900	4Z5	C9-N10	4.92	1.36	1.32

All (29) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	900	4Z5	N2-C3-N4	-6.79	119.95	126.63
2	B	801	4Z5	N2-C3-N4	-5.46	121.26	126.63
2	A	900	4Z5	C6-C5-N4	-3.98	118.97	122.47
2	B	801	4Z5	C6-C5-N4	-3.72	119.19	122.47
2	B	801	4Z5	C11-C9-C8	-3.28	119.96	122.74
2	B	801	4Z5	C12-C8-C9	-2.97	118.93	123.15
2	A	900	4Z5	C12-C8-C9	-2.77	119.22	123.15
2	B	801	4Z5	C8-C12-C19	-2.71	116.09	121.52
2	A	900	4Z5	C8-C9-N10	-2.38	120.12	122.08
2	A	900	4Z5	C8-C7-C6	-2.32	118.42	121.90
2	B	801	4Z5	C6-C5-N10	-2.32	120.43	122.47
2	A	900	4Z5	C14-N13-C3	-2.30	120.81	123.59
2	A	900	4Z5	C8-C12-C19	-2.28	116.94	121.52
2	B	801	4Z5	C6-C1-N2	-2.18	119.76	123.97
2	A	900	4Z5	C6-C1-N2	-2.08	119.94	123.97
2	B	801	4Z5	C1-N2-C3	2.25	119.48	115.84
2	A	900	4Z5	C15-C16-C17	2.31	119.52	117.87
2	A	900	4Z5	C9-N10-C5	2.45	120.22	117.68
2	A	900	4Z5	C25-N24-C23	2.49	126.53	121.74
2	B	801	4Z5	C7-C8-C12	2.55	122.71	117.01
2	A	900	4Z5	N13-C3-N4	2.57	119.43	116.96
2	B	801	4Z5	C15-C16-C17	2.61	119.73	117.87
2	A	900	4Z5	C7-C8-C9	3.00	120.10	117.92
2	A	900	4Z5	C1-N2-C3	3.00	120.70	115.84
2	B	801	4Z5	N13-C3-N4	3.28	120.11	116.96
2	B	801	4Z5	C9-N10-C5	3.41	121.21	117.68
2	A	900	4Z5	N13-C3-N2	3.75	120.62	117.00
2	B	801	4Z5	C3-N4-C5	4.70	120.47	115.11
2	A	900	4Z5	C3-N4-C5	5.52	121.41	115.11

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	254/307 (82%)	-0.44	7 (2%) 53 53	10, 23, 47, 57	0
1	B	251/307 (81%)	-0.42	8 (3%) 47 47	11, 23, 47, 60	0
All	All	505/614 (82%)	-0.43	15 (2%) 50 50	10, 23, 47, 60	0

All (15) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	487	THR	4.1
1	B	630	ASN	3.5
1	B	626	MET	3.3
1	A	628	ASP	3.0
1	B	466	SER	3.0
1	A	627	GLN	3.0
1	B	719	SER	3.0
1	A	629	LYS	2.8
1	B	596	LEU	2.8
1	A	630	ASN	2.7
1	A	467	PHE	2.4
1	B	465	GLY	2.3
1	B	595	GLY	2.3
1	A	718	ARG	2.2
1	A	598	THR	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	CL	B	802	1/1	0.97	0.10	50,50,50,50	0
2	4Z5	B	801	31/31	0.98	0.10	12,15,18,19	0
2	4Z5	A	900	31/31	0.98	0.10	6,11,15,16	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.