



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 17, 2018 – 05:59 am GMT

PDB ID : 1CBM
Title : THE 1.8 ANGSTROM STRUCTURE OF CARBONMONOXY-BETA4
HEMOGLOBIN: ANALYSIS OF A HOMOTETRAMER WITH THE R QUATERNARY
STRUCTURE OF LIGANDED ALPHA2BETA2 HEMOGLOBIN
Authors : Borgstahl, G.E.O.; Arnone, A.
Deposited on : 1993-02-18
Resolution : 1.74 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7.3 (157068), CSD as539be (2018)
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
Percentile statistics	:	20171227.v01 (using entries in the PDB archive December 27th 2017)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	trunk30686


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.74 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	122078	3201 (1.76-1.72)
Ramachandran outliers	120005	3169 (1.76-1.72)
Sidechain outliers	119972	3169 (1.76-1.72)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	146	 73% 21% 7%
1	B	146	 73% 24% ..
1	C	146	 73% 23% ..
1	D	146	 70% 25% 5% .

2 Entry composition [i](#)


There are 5 unique types of molecules in this entry. The entry contains 4953 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HEMOGLOBIN BETA 4 (CARBONMONOXY).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	146	Total	C	N	O	S	0	6	0
			1144	739	198	203	4			
1	B	146	Total	C	N	O	S	0	11	0
			1162	750	199	209	4			
1	C	146	Total	C	N	O	S	0	8	0
			1153	744	200	205	4			
1	D	146	Total	C	N	O	S	0	10	0
			1161	748	200	209	4			

- Molecule 2 is SULFATE ION (three-letter code: SO₄) (formula: O₄S).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	C	1	Total	O	S	0	0
			5	4	1		
2	D	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
3	B	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
3	C	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
3	D	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		

- Molecule 4 is CARBON MONOXIDE (three-letter code: CMO) (formula: CO).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			2	1	1		
4	B	1	Total	C	O	0	0
			2	1	1		
4	C	1	Total	C	O	0	0
			2	1	1		
4	D	1	Total	C	O	0	0
			2	1	1		

- Molecule 5 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	31	Total	O	0	0
			31	31		
5	B	34	Total	O	0	0
			34	34		
5	C	46	Total	O	0	0
			46	46		
5	D	22	Total	O	0	0
			22	22		

3 Residue-property plots


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


• Molecule 1: HEMOGLOBIN BETA 4 (CARBONMONOXY)




• Molecule 1: HEMOGLOBIN BETA 4 (CARBONMONOXY)



• Molecule 1: HEMOGLOBIN BETA 4 (CARBONMONOXY)



• Molecule 1: HEMOGLOBIN BETA 4 (CARBONMONOXY)



H117	F118	G119	K120	E121	Y130	A140	L141	A142	H143	K144	Y145	H146
------	------	------	------	------	------	------	------	------	------	------	------	------

4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	63.30 Å 82.40 Å 53.70 Å 90.00° 90.10° 90.00°	Depositor
Resolution (Å)	10.00 – 1.74	Depositor
% Data completeness (in resolution range)	(Not available) (10.00-1.74)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	PROLSQ, TNT	Depositor
R, R_{free}	0.177 , (Not available)	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	4953	wwPDB-VP
Average B, all atoms (Å ²)	30.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CMO, HEM, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.88	0/1205	1.47	13/1637 (0.8%)
1	B	0.89	0/1249	1.42	10/1695 (0.6%)
1	C	0.93	0/1225	1.64	20/1664 (1.2%)
1	D	0.92	2/1243 (0.2%)	1.52	15/1688 (0.9%)
All	All	0.91	2/4922 (0.0%)	1.52	58/6684 (0.9%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	112[A]	CYS	CB-SG	-5.22	1.73	1.81
1	D	112[B]	CYS	CB-SG	-5.22	1.73	1.81

The worst 5 of 58 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	30	ARG	NE-CZ-NH1	18.99	129.80	120.30
1	C	30	ARG	NE-CZ-NH2	-12.19	114.21	120.30
1	A	30	ARG	NE-CZ-NH1	11.29	125.95	120.30
1	D	104	ARG	NE-CZ-NH1	11.11	125.85	120.30
1	C	104	ARG	NE-CZ-NH1	10.84	125.72	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1144	0	1145	19	0
1	B	1162	0	1160	27	0
1	C	1153	0	1150	22	0
1	D	1161	0	1151	28	0
2	A	5	0	0	0	0
2	B	5	0	0	0	0
2	C	5	0	0	0	0
2	D	5	0	0	0	0
3	A	43	0	30	0	0
3	B	43	0	30	4	0
3	C	43	0	30	1	0
3	D	43	0	30	0	0
4	A	2	0	0	0	0
4	B	2	0	0	0	0
4	C	2	0	0	0	0
4	D	2	0	0	0	0
5	A	31	0	0	1	0
5	B	34	0	0	2	0
5	C	46	0	0	0	0
5	D	22	0	0	0	0
All	All	4953	0	4726	94	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

The worst 5 of 94 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:90:GLU:HG3	1:B:144:LYS:HE2	1.63	0.81
1:A:82:LYS:HD2	1:A:140:ALA:HA	1.71	0.72
1:D:23:VAL:HG12	1:D:68:LEU:HD11	1.72	0.72
1:B:55:MET:HE3	5:B:214:HOH:O	1.92	0.69
1:D:14:LEU:HD21	1:D:121[B]:GLU:HG3	1.75	0.68

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	150/146 (103%)	146 (97%)	3 (2%)	1 (1%)	24	8
1	B	155/146 (106%)	151 (97%)	4 (3%)	0	100	100
1	C	152/146 (104%)	148 (97%)	4 (3%)	0	100	100
1	D	154/146 (106%)	150 (97%)	4 (3%)	0	100	100
All	All	611/584 (105%)	595 (97%)	15 (2%)	1 (0%)	49	30

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	119	GLY

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	124/118 (105%)	113 (91%)	11 (9%)	11	1
1	B	129/118 (109%)	123 (95%)	6 (5%)	29	8
1	C	126/118 (107%)	117 (93%)	9 (7%)	16	2
1	D	128/118 (108%)	114 (89%)	14 (11%)	7	1
All	All	507/472 (107%)	467 (92%)	40 (8%)	15	2

5 of 40 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	66	LYS
1	C	99[A]	ASP
1	D	99[B]	ASP
1	C	79	ASP
1	C	99[B]	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	117	HIS
1	B	131	GLN
1	C	131	GLN
1	C	139	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

12 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	HEM	A	147	1,4	27,50,50	1.80	7 (25%)	17,82,82	1.79	4 (23%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	CMO	A	148	3	0,1,1	0.00	-	0,0,0	0.00	-
2	SO4	A	150	-	4,4,4	0.75	0	6,6,6	0.24	0
3	HEM	B	147	1,4	27,50,50	1.93	6 (22%)	17,82,82	2.10	7 (41%)
4	CMO	B	148	3	0,1,1	0.00	-	0,0,0	0.00	-
2	SO4	B	151	-	4,4,4	0.77	0	6,6,6	0.33	0
3	HEM	C	147	1,4	27,50,50	1.82	5 (18%)	17,82,82	2.15	8 (47%)
4	CMO	C	148	3	0,1,1	0.00	-	0,0,0	0.00	-
2	SO4	C	153	-	4,4,4	0.74	0	6,6,6	0.12	0
3	HEM	D	147	1,4	27,50,50	2.00	7 (25%)	17,82,82	2.17	5 (29%)
4	CMO	D	148	3	0,1,1	0.00	-	0,0,0	0.00	-
2	SO4	D	152	-	4,4,4	0.75	0	6,6,6	0.09	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	HEM	A	147	1,4	-	0/6/54/54	0/0/8/8
4	CMO	A	148	3	-	0/0/0/0	0/0/0/0
2	SO4	A	150	-	-	0/0/0/0	0/0/0/0
3	HEM	B	147	1,4	-	0/6/54/54	0/0/8/8
4	CMO	B	148	3	-	0/0/0/0	0/0/0/0
2	SO4	B	151	-	-	0/0/0/0	0/0/0/0
3	HEM	C	147	1,4	-	0/6/54/54	0/0/8/8
4	CMO	C	148	3	-	0/0/0/0	0/0/0/0
2	SO4	C	153	-	-	0/0/0/0	0/0/0/0
3	HEM	D	147	1,4	-	0/6/54/54	0/0/8/8
4	CMO	D	148	3	-	0/0/0/0	0/0/0/0
2	SO4	D	152	-	-	0/0/0/0	0/0/0/0

The worst 5 of 25 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	D	147	HEM	C3C-C2C	-5.21	1.33	1.40
3	C	147	HEM	C3C-C2C	-4.85	1.33	1.40
3	B	147	HEM	C3B-C2B	-4.59	1.34	1.40
3	B	147	HEM	C3C-C2C	-4.16	1.34	1.40
3	A	147	HEM	C3C-C2C	-4.04	1.34	1.40

The worst 5 of 24 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	D	147	HEM	CMA-C3A-C4A	-5.36	120.22	128.46
3	C	147	HEM	CMD-C2D-C1D	-4.41	121.68	128.46
3	B	147	HEM	CMA-C3A-C4A	-4.15	122.09	128.46
3	A	147	HEM	CMA-C3A-C4A	-3.55	123.01	128.46
3	D	147	HEM	CMD-C2D-C1D	-3.30	123.39	128.46

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	147	HEM	4	0
3	C	147	HEM	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.