



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 9, 2018 – 12:28 am GMT

PDB ID : 2CDB
Title : Sulfolobus solfataricus Glucose Dehydrogenase 1 in complex with NADP and glucose
Authors : Milburn, C.C.; Lambie, H.J.; Theodossis, A.; Hough, D.W.; Danson, M.J.; Taylor, G.L.
Deposited on : 2006-01-23
Resolution : 1.60 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.3 (157068), CSD as539be (2018)
Xtriage (Phenix) : 1.13
EDS : trunk30967
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)
Refmac : 5.8.0158
CCP4 : 7.0 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk30967


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	111664	2957 (1.60-1.60)
Clashscore	122126	3202 (1.60-1.60)
Ramachandran outliers	120053	3117 (1.60-1.60)
Sidechain outliers	120020	3116 (1.60-1.60)
RSRZ outliers	108989	2883 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	366	<div> <div>4%</div> <div> <div></div> <div>84%</div> <div>13%</div> <div>..</div> </div> </div>
1	B	366	<div> <div>4%</div> <div> <div></div> <div>84%</div> <div>13%</div> <div>..</div> </div> </div>
1	C	366	<div> <div>5%</div> <div> <div></div> <div>84%</div> <div>13%</div> <div>..</div> </div> </div>
1	D	366	<div> <div>3%</div> <div> <div></div> <div>86%</div> <div>11%</div> <div>..</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit crite-

ria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	EDO	B	1367	-	-	X	-

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 13128 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called GLUCOSE 1-DEHYDROGENASE (DHG-1).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	360	Total	C	N	O	S	0	20	0
			2924	1870	494	545	15			
1	B	359	Total	C	N	O	S	0	18	0
			2899	1856	487	543	13			
1	C	358	Total	C	N	O	S	0	16	0
			2886	1847	491	535	13			
1	D	359	Total	C	N	O	S	0	12	0
			2877	1838	488	538	13			

There are 4 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	41	ALA	THR	engineered mutation	UNP O93715
B	41	ALA	THR	engineered mutation	UNP O93715
C	41	ALA	THR	engineered mutation	UNP O93715
D	41	ALA	THR	engineered mutation	UNP O93715

- Molecule 2 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			4	2	2		
2	B	1	Total	C	O	0	0
			4	2	2		
2	C	1	Total	C	O	0	0
			4	2	2		
2	D	1	Total	C	O	0	0
			4	2	2		

- Molecule 3 is BETA-D-GLUCOSE (three-letter code: BGC) (formula: $C_6H_{12}O_6$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total 13	C 6	O 7	0	1
3	B	1	Total 12	C 6	O 6	0	0
3	C	1	Total 12	C 6	O 6	0	0
3	D	1	Total 13	C 6	O 7	0	1

- Molecule 4 is NADP NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (three-letter code: NAP) (formula: $C_{21}H_{28}N_7O_{17}P_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
4	A	1	Total 48	C 21	N 7	O 17	P 3	0	0
4	B	1	Total 48	C 21	N 7	O 17	P 3	0	0
4	C	1	Total 48	C 21	N 7	O 17	P 3	0	0
4	D	1	Total 48	C 21	N 7	O 17	P 3	0	0

- Molecule 5 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	B	2	Total Zn 2 2	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	2	Total 2	Zn 2	0	0
5	D	2	Total 2	Zn 2	0	0
5	C	2	Total 2	Zn 2	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	320	Total 320	O 320	0	0
6	B	291	Total 291	O 291	0	0
6	C	323	Total 323	O 323	0	0
6	D	342	Total 342	O 342	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: GLUCOSE 1-DEHYDROGENASE (DHG-1)




• Molecule 1: GLUCOSE 1-DEHYDROGENASE (DHG-1)




• Molecule 1: GLUCOSE 1-DEHYDROGENASE (DHG-1)



• Molecule 1: GLUCOSE 1-DEHYDROGENASE (DHG-1)





4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	68.67Å 91.39Å 138.43Å 90.00° 90.03° 90.00°	Depositor
Resolution (Å)	138.68 – 1.60 38.29 – 1.60	Depositor EDS
% Data completeness (in resolution range)	96.4 (138.68-1.60) 96.3 (38.29-1.60)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.03 (at 1.60Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.190 , 0.218 0.193 , 0.222	Depositor DCC
R_{free} test set	10908 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	13.3	Xtriage
Anisotropy	0.531	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 32.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.41$, $\langle L^2 \rangle = 0.23$	Xtriage
Estimated twinning fraction	0.218 for h,-k,-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	13128	wwPDB-VP
Average B, all atoms (Å ²)	13.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 18.45% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, BGC, NAP, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.86	6/3035 (0.2%)	0.73	8/4099 (0.2%)
1	B	0.67	4/3012 (0.1%)	0.78	8/4069 (0.2%)
1	C	0.49	0/3010	0.62	0/4064
1	D	0.72	2/2963 (0.1%)	0.65	6/4001 (0.1%)
All	All	0.70	12/12020 (0.1%)	0.70	22/16233 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	1	0
1	B	1	1
1	D	0	1
All	All	2	2

The worst 5 of 12 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	255[A]	GLY	N-CA	21.21	1.77	1.46
1	A	255[B]	GLY	N-CA	21.21	1.77	1.46
1	D	254[A]	THR	CA-C	20.47	2.06	1.52
1	D	254[B]	THR	CA-C	20.47	2.06	1.52
1	A	346[A]	GLU	CG-CD	-16.51	1.27	1.51

The worst 5 of 22 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	254[A]	THR	CA-C-O	13.82	149.12	120.10

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	254[B]	THR	CA-C-O	13.82	149.12	120.10
1	B	254[A]	THR	CA-C-N	-10.69	94.82	116.20
1	B	254[B]	THR	CA-C-N	-10.69	94.82	116.20
1	A	254[A]	THR	CA-C-N	-9.86	96.48	116.20

All (2) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
1	A	254[A]	THR	CA
1	B	254[A]	THR	CA

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	254[A]	THR	Peptide
1	D	255[B]	GLY	Mainchain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2924	0	2978	52	0
1	B	2899	0	2951	55	0
1	C	2886	0	2946	49	0
1	D	2877	0	2918	43	0
2	A	4	0	6	2	0
2	B	4	0	6	5	0
2	C	4	0	6	0	0
2	D	4	0	6	0	0
3	A	13	0	6	0	0
3	B	12	0	12	0	0
3	C	12	0	12	0	0
3	D	13	0	6	0	0
4	A	48	0	25	1	0
4	B	48	0	25	0	0
4	C	48	0	25	4	0
4	D	48	0	25	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	A	2	0	0	0	0
5	B	2	0	0	0	0
5	C	2	0	0	0	0
5	D	2	0	0	0	0
6	A	320	0	0	10	0
6	B	291	0	0	6	0
6	C	323	0	0	6	0
6	D	342	0	0	8	0
All	All	13128	0	11953	191	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 191 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:254[A]:THR:C	1:B:254[A]:THR:CA	1.75	1.52
1:A:255[A]:GLY:CA	1:A:255[A]:GLY:N	1.77	1.46
1:A:254[B]:THR:C	1:A:254[B]:THR:CA	1.86	1.43
1:D:254[A]:THR:C	1:D:254[A]:THR:CA	2.06	1.24
1:B:254[B]:THR:C	1:B:254[B]:THR:CA	2.11	1.19

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	376/366 (103%)	368 (98%)	7 (2%)	1 (0%)	43	22
1	B	373/366 (102%)	361 (97%)	11 (3%)	1 (0%)	43	22
1	C	370/366 (101%)	360 (97%)	9 (2%)	1 (0%)	43	22

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	D	367/366 (100%)	358 (98%)	8 (2%)	1 (0%)	43	22
All	All	1486/1464 (102%)	1447 (97%)	35 (2%)	4 (0%)	43	22

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	307	ASN
1	B	307	ASN
1	D	307	ASN
1	C	307	ASN

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	327/317 (103%)	317 (97%)	10 (3%)	43	17
1	B	325/317 (102%)	319 (98%)	6 (2%)	62	38
1	C	326/317 (103%)	318 (98%)	8 (2%)	50	24
1	D	319/317 (101%)	316 (99%)	3 (1%)	81	68
All	All	1297/1268 (102%)	1270 (98%)	27 (2%)	60	30

5 of 27 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	140	LYS
1	B	354[B]	LYS
1	D	133	LYS
1	B	346	GLU
1	A	232	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 22 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	259	ASN
1	B	297	HIS
1	D	297	HIS
1	B	263	ASN
1	B	293	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 22 ligands modelled in this entry, 8 are monoatomic - leaving 14 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	EDO	A	1367	-	3,3,3	0.53	0	2,2,2	0.30	0
3	BGC	A	1368[A]	-	12,12,12	0.60	0	17,17,17	0.89	1 (5%)
3	BGC	A	1368[B]	-	12,12,12	0.59	0	17,17,17	0.86	1 (5%)
4	NAP	A	1369	-	44,52,52	1.09	4 (9%)	53,80,80	1.78	10 (18%)
2	EDO	B	1367	-	3,3,3	0.51	0	2,2,2	0.39	0
3	BGC	B	1368	-	12,12,12	0.60	0	17,17,17	0.72	0
4	NAP	B	1369	-	44,52,52	1.16	6 (13%)	53,80,80	1.66	5 (9%)
2	EDO	C	1367	-	3,3,3	0.48	0	2,2,2	0.25	0
3	BGC	C	1368	-	12,12,12	0.57	0	17,17,17	0.70	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	NAP	C	1369	-	44,52,52	1.43	7 (15%)	53,80,80	1.81	6 (11%)
2	EDO	D	1367	-	3,3,3	0.64	0	2,2,2	0.14	0
3	BGC	D	1368[A]	-	12,12,12	0.51	0	17,17,17	0.86	0
3	BGC	D	1368[B]	-	12,12,12	0.52	0	17,17,17	0.88	0
4	NAP	D	1369	-	44,52,52	1.11	5 (11%)	53,80,80	1.61	8 (15%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	EDO	A	1367	-	-	0/1/1/1	0/0/0/0
3	BGC	A	1368[A]	-	-	0/2/22/22	0/1/1/1
3	BGC	A	1368[B]	-	-	0/2/22/22	0/1/1/1
4	NAP	A	1369	-	-	0/27/67/67	0/5/5/5
2	EDO	B	1367	-	-	0/1/1/1	0/0/0/0
3	BGC	B	1368	-	-	0/2/22/22	0/1/1/1
4	NAP	B	1369	-	-	0/27/67/67	0/5/5/5
2	EDO	C	1367	-	-	0/1/1/1	0/0/0/0
3	BGC	C	1368	-	-	0/2/22/22	0/1/1/1
4	NAP	C	1369	-	-	0/27/67/67	0/5/5/5
2	EDO	D	1367	-	-	0/1/1/1	0/0/0/0
3	BGC	D	1368[A]	-	-	0/2/22/22	0/1/1/1
3	BGC	D	1368[B]	-	-	0/2/22/22	0/1/1/1
4	NAP	D	1369	-	-	0/27/67/67	0/5/5/5

The worst 5 of 22 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	C	1369	NAP	C8A-N9A	-2.47	1.33	1.36
4	D	1369	NAP	C8A-N9A	-2.13	1.34	1.36
4	B	1369	NAP	C3N-C7N	2.03	1.53	1.50
4	B	1369	NAP	P2B-O2B	2.05	1.63	1.59
4	A	1369	NAP	C6N-N1N	2.11	1.40	1.35

The worst 5 of 31 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	C	1369	NAP	N3A-C2A-N1A	-9.95	120.35	128.86
4	A	1369	NAP	N3A-C2A-N1A	-8.93	121.22	128.86
4	B	1369	NAP	N3A-C2A-N1A	-8.83	121.30	128.86

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	D	1369	NAP	N3A-C2A-N1A	-8.34	121.72	128.86
4	C	1369	NAP	C4B-O4B-C1B	-3.52	106.15	109.83

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 12 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1367	EDO	2	0
4	A	1369	NAP	1	0
2	B	1367	EDO	5	0
4	C	1369	NAP	4	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	B	2
1	D	2

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	D	255[A]:GLY	C	256[A]:ALA	N	2.08
1	B	254[A]:THR	C	255[A]:GLY	N	1.61
1	B	254[B]:THR	C	255[B]:GLY	N	1.08
1	D	255[B]:GLY	C	256[B]:ALA	N	0.63

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	360/366 (98%)	-0.12	16 (4%) 34 31	6, 10, 21, 35	17 (4%)
1	B	359/366 (98%)	-0.12	14 (3%) 39 37	5, 12, 24, 33	13 (3%)
1	C	358/366 (97%)	-0.13	18 (5%) 29 27	5, 11, 21, 31	20 (5%)
1	D	359/366 (98%)	-0.15	11 (3%) 49 47	6, 11, 22, 28	10 (2%)
All	All	1436/1464 (98%)	-0.13	59 (4%) 37 35	5, 11, 22, 35	60 (4%)

The worst 5 of 59 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	77	HIS	8.1
1	A	77	HIS	6.8
1	B	34[A]	ILE	6.8
1	B	100[A]	LEU	6.6
1	A	76	TYR	6.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	EDO	B	1367	4/4	0.75	0.20	22,25,29,32	0
2	EDO	A	1367	4/4	0.86	0.20	21,26,28,31	0
3	BGC	C	1368	12/12	0.92	0.15	9,14,20,22	0
2	EDO	C	1367	4/4	0.94	0.09	14,15,15,16	0
3	BGC	A	1368[A]	12/12	0.94	0.11	9,13,17,17	1
3	BGC	A	1368[B]	12/12	0.94	0.11	9,13,16,17	1
3	BGC	D	1368[B]	12/12	0.96	0.14	8,11,14,15	1
3	BGC	B	1368	12/12	0.96	0.10	10,13,16,18	0
3	BGC	D	1368[A]	12/12	0.96	0.14	8,11,14,14	1
4	NAP	D	1369	48/48	0.97	0.07	6,10,13,14	0
4	NAP	C	1369	48/48	0.98	0.06	4,8,11,12	0
4	NAP	A	1369	48/48	0.98	0.06	4,9,12,14	0
4	NAP	B	1369	48/48	0.98	0.07	6,9,13,14	0
2	EDO	D	1367	4/4	0.98	0.05	14,15,15,16	0
5	ZN	D	1370	1/1	1.00	0.05	7,7,7,7	1
5	ZN	C	1371	1/1	1.00	0.03	8,8,8,8	0
5	ZN	B	1371	1/1	1.00	0.02	9,9,9,9	0
5	ZN	D	1371	1/1	1.00	0.03	9,9,9,9	0
5	ZN	B	1370	1/1	1.00	0.05	6,6,6,6	1
5	ZN	A	1371	1/1	1.00	0.03	9,9,9,9	0
5	ZN	C	1370	1/1	1.00	0.03	6,6,6,6	1
5	ZN	A	1370	1/1	1.00	0.04	6,6,6,6	1

6.5 Other polymers ⓘ

There are no such residues in this entry.