



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 4, 2020 – 12:44 PM EST

PDB ID : 2CF5
Title : Crystal Structures of the Arabidopsis Cinnamyl Alcohol Dehydrogenases, At-CAD5
Authors : Youn, B.; Camacho, R.; Moinuddin, S.; Lee, C.; Davin, L.B.; Lewis, N.G.; Kang, C.
Deposited on : 2006-02-16
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.8


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	122126	8267 (2.00-2.00)
Ramachandran outliers	120053	8166 (2.00-2.00)
Sidechain outliers	120020	8165 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	357	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 2866 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CINNAMYL ALCOHOL DEHYDROGENASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	352	Total	C	N	O	S	0	0	0
			2665	1686	450	507	22			

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	Zn	0	0
			2	2		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	199	Total	O	0	0
			199	199		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: CINNAMYL ALCOHOL DEHYDROGENASE

Chain A: 



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, α , β , γ	54.22Å 54.22Å 312.99Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	10.00 – 2.00	Depositor
% Data completeness (in resolution range)	98.2 (10.00-2.00)	Depositor
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	X-PLOR	Depositor
R, R_{free}	0.195 , 0.235	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	2866	wwPDB-VP
Average B, all atoms (Å ²)	40.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.66	0/2715	1.33	13/3674 (0.4%)

There are no bond length outliers.

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	274	MET	CG-SD-CE	-7.74	87.81	100.20
1	A	119	TRP	CD1-CG-CD2	7.54	112.33	106.30
1	A	119	TRP	CE2-CD2-CG	-7.37	101.40	107.30
1	A	13	TRP	CE2-CD2-CG	-7.28	101.48	107.30
1	A	13	TRP	CD1-CG-CD2	7.24	112.09	106.30
1	A	266	LYS	CA-CB-CG	6.63	127.99	113.40
1	A	54	THR	CA-CB-CG2	6.33	121.26	112.40
1	A	75	VAL	CG1-CB-CG2	-6.11	101.12	110.90
1	A	58	LEU	CA-CB-CG	6.04	129.20	115.30
1	A	353	SER	N-CA-C	5.60	126.13	111.00
1	A	343	ARG	CA-C-N	-5.40	105.33	117.20
1	A	65	MET	CG-SD-CE	-5.15	91.95	100.20
1	A	265	LEU	CA-CB-CG	5.03	126.87	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2665	0	2666	56	0
2	A	2	0	0	0	0
3	A	199	0	0	1	0
All	All	2866	0	2666	56	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (56) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:156:GLU:HB3	1:A:347:VAL:HG11	1.67	0.76
1:A:278:ASN:H	1:A:278:ASN:HD22	1.36	0.71
1:A:274:MET:HE3	1:A:298:SER:HB3	1.76	0.65
1:A:189:LEU:HD12	1:A:216:LYS:HB3	1.80	0.63
1:A:54:THR:HG22	1:A:55:LYS:HG3	1.83	0.61
1:A:278:ASN:H	1:A:278:ASN:ND2	2.01	0.59
1:A:58:LEU:HD13	1:A:58:LEU:H	1.67	0.59
1:A:53:GLN:HE21	1:A:58:LEU:HD21	1.66	0.59
1:A:165:GLY:HA3	1:A:309:MET:HE3	1.87	0.56
1:A:21:ILE:HG23	1:A:331:ASN:ND2	2.20	0.56
1:A:43:CYS:SG	1:A:72:VAL:HG13	2.45	0.56
1:A:21:ILE:HG23	1:A:331:ASN:HD21	1.71	0.55
1:A:56:ASN:ND2	1:A:61:SER:H	2.04	0.54
1:A:153:MET:SD	1:A:158:ALA:HB2	2.48	0.54
1:A:287:LEU:HA	1:A:290:LEU:HB3	1.90	0.53
1:A:267:LEU:HD23	1:A:291:GLY:O	2.09	0.53
1:A:327:MET:O	1:A:330:VAL:HG22	2.09	0.52
1:A:103:CYS:SG	1:A:105:PRO:HD2	2.51	0.51
1:A:274:MET:CE	1:A:298:SER:HB3	2.41	0.51
1:A:309:MET:HE1	1:A:313:CYS:SG	2.51	0.51
1:A:38:ASN:HB3	1:A:77:GLU:HB3	1.93	0.50
1:A:198:LYS:HZ1	1:A:319:SER:H	1.61	0.49
1:A:66:VAL:H	1:A:132:GLN:NE2	2.10	0.49
1:A:167:THR:O	1:A:274:MET:HE1	2.13	0.49
1:A:65:MET:SD	1:A:122:ASN:ND2	2.86	0.49
1:A:220:ALA:HA	1:A:224:LEU:HD12	1.95	0.48
1:A:75:VAL:HG23	1:A:87:VAL:HA	1.95	0.48
1:A:62:ASN:H	1:A:122:ASN:HD21	1.59	0.48
1:A:324:VAL:HA	1:A:347:VAL:O	2.13	0.47
1:A:53:GLN:NE2	1:A:58:LEU:HD21	2.30	0.46
1:A:10:THR:HG21	1:A:138:ALA:HB3	1.97	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:216:LYS:HB2	1:A:229:TYR:OH	2.16	0.46
1:A:71:VAL:CG2	1:A:134:GLY:HA3	2.47	0.45
1:A:189:LEU:HD11	1:A:220:ALA:HB2	1.99	0.45
1:A:158:ALA:HA	1:A:161:LEU:HD22	1.98	0.45
1:A:249:ILE:HD13	1:A:272:ILE:HB	1.99	0.44
1:A:187:LEU:HD23	1:A:250:ASP:OD1	2.17	0.44
1:A:158:ALA:O	1:A:162:LEU:HG	2.17	0.44
1:A:93:VAL:HG21	1:A:141:VAL:HG21	2.01	0.43
1:A:56:ASN:O	1:A:58:LEU:HD22	2.18	0.43
1:A:8:ARG:CZ	1:A:8:ARG:HA	2.49	0.43
1:A:108:ARG:HB3	1:A:110:LEU:HD22	2.01	0.43
1:A:67:PRO:O	1:A:133:GLY:HA3	2.18	0.43
1:A:154:ALA:HB3	1:A:157:GLN:HB2	2.01	0.43
1:A:51:LEU:HD12	1:A:337:LEU:HD23	2.01	0.43
1:A:336:ARG:HB3	1:A:341:ASP:HB3	2.01	0.42
1:A:137:LYS:HD2	1:A:350:VAL:HG12	2.01	0.42
1:A:220:ALA:HA	1:A:224:LEU:HB2	2.02	0.42
1:A:168:VAL:HG23	1:A:192:VAL:HG12	2.02	0.42
1:A:194:HIS:O	1:A:198:LYS:HG3	2.20	0.42
1:A:309:MET:CE	1:A:313:CYS:SG	3.08	0.42
1:A:201:LYS:HA	1:A:201:LYS:HD2	1.91	0.41
1:A:11:THR:HG21	1:A:356:ASP:H	1.86	0.41
1:A:270:LYS:HG3	1:A:294:VAL:HB	2.02	0.40
1:A:304:LYS:HG3	1:A:305:GLU:N	2.36	0.40
1:A:18:PRO:HG2	3:A:2011:HOH:O	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	350/357 (98%)	309 (88%)	35 (10%)	6 (2%)	10 4

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	57	ASP
1	A	228	ASP
1	A	242	ALA
1	A	300	ILE
1	A	122	ASN
1	A	344	TYR

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	295/300 (98%)	269 (91%)	26 (9%)	11 6

All (26) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	8	ARG
1	A	28	THR
1	A	56	ASN
1	A	58	LEU
1	A	82	VAL
1	A	100	CYS
1	A	109	ASP
1	A	110	LEU
1	A	111	GLU
1	A	147	VAL
1	A	156	GLU
1	A	157	GLN
1	A	166	VAL
1	A	222	GLN
1	A	252	VAL
1	A	254	VAL
1	A	264	LEU
1	A	265	LEU
1	A	267	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	278	ASN
1	A	290	LEU
1	A	303	MET
1	A	309	MET
1	A	314	LYS
1	A	322	ILE
1	A	354	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	38	ASN
1	A	53	GLN
1	A	56	ASN
1	A	122	ASN
1	A	129	GLN
1	A	132	GLN
1	A	255	HIS
1	A	278	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.