



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2018 – 05:20 am GMT

PDB ID : 2CFA  
Title : Structure of viral flavin-dependant thymidylate synthase ThyX  
Authors : Graziani, S.; Bernauer, J.; Skouloubris, S.; Graille, M.; Zhou, C.-Z.; Marchand, C.; Decottignies, P.; van Tilbeurgh, H.; Myllykallio, H.; Liebl, U.  
Deposited on : 2006-02-17  
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.3 (157068), CSD as539be (2018)  
Xtriage (Phenix) : 1.13  
EDS : trunk30967  
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)  
Refmac : 5.8.0158  
CCP4 : 7.0 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk30967


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	111664	4477 (2.30-2.30)
Clashscore	122126	5072 (2.30-2.30)
Ramachandran outliers	120053	5022 (2.30-2.30)
Sidechain outliers	120020	5021 (2.30-2.30)
RSRZ outliers	108989	4374 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	217	<div> <div>8%</div> <div> <div></div> <div>65%</div> <div>14%</div> <div>•</div> <div>17%</div> </div> </div>
2	B	217	<div> <div>7%</div> <div> <div></div> <div>63%</div> <div>18%</div> <div>•</div> <div>18%</div> </div> </div>

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 3201 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called THYMIDYLATE SYNTHASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	180	1460	935	252	265	8	0	0	0


- Molecule 2 is a protein called THYMIDYLATE SYNTHASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	179	1454	932	250	264	8	1	0	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	77	ILE	LEU	conflict	UNP O41156

- Molecule 3 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula: C<sub>27</sub>H<sub>33</sub>N<sub>9</sub>O<sub>15</sub>P<sub>2</sub>).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
3	B	1	Total	C	N	O	P	0	0
			53	27	9	15	2		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	106	Total	O	0	0
			106	106		
4	B	75	Total	O	0	0
			75	75		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: THYMIDYLATE SYNTHASE



#### • Molecule 2: THYMIDYLATE SYNTHASE



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	69.26Å 76.99Å 93.44Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 2.30 33.65 – 2.30	Depositor EDS
% Data completeness (in resolution range)	98.6 (20.00-2.30) 98.6 (33.65-2.30)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.92 (at 2.29Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, $R_{free}$	0.213 , 0.262 0.207 , 0.256	Depositor DCC
$R_{free}$ test set	1124 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	32.2	Xtriage
Anisotropy	0.795	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 50.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	3201	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	41.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.49% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CME, FAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.48	0/1482	0.60	0/2000
2	B	0.49	0/1473	0.60	1/1983 (0.1%)
All	All	0.49	0/2955	0.60	1/3983 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	151	LYS	N-CA-CB	7.91	124.83	110.60

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1460	0	1456	22	0
2	B	1454	0	1456	34	0
3	A	53	0	31	1	0
3	B	53	0	31	2	0
4	A	106	0	0	2	0
4	B	75	0	0	6	0
All	All	3201	0	2974	55	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (55) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:98:PRO:C	2:B:208:SER:N	1.74	1.41
1:A:159:PRO:O	1:A:162:THR:HG22	1.50	1.11
2:B:76:VAL:HG11	2:B:168:MET:CE	2.01	0.91
2:B:34:ASN:HB3	2:B:36:ILE:HG22	1.50	0.91
2:B:76:VAL:CG1	2:B:168:MET:HE2	2.01	0.90
2:B:76:VAL:HG11	2:B:168:MET:HE2	1.57	0.86
2:B:66:LEU:HD11	2:B:168:MET:HE3	1.63	0.80
2:B:76:VAL:CG1	2:B:168:MET:CE	2.59	0.79
2:B:66:LEU:CD1	2:B:168:MET:HE3	2.13	0.79
1:A:67:LYS:HD3	1:A:165:THR:HB	1.68	0.75
1:A:1:MET:N	1:A:67:LYS:O	2.20	0.74
2:B:66:LEU:HD11	2:B:168:MET:CE	2.23	0.68
2:B:76:VAL:HG11	2:B:168:MET:HE1	1.76	0.67
2:B:182:ARG:HA	2:B:187:THR:HG21	1.77	0.65
1:A:182:ARG:HA	1:A:187:THR:HG21	1.84	0.60
2:B:66:LEU:CD1	2:B:168:MET:CE	2.80	0.59
2:B:180:GLU:HG3	4:B:2057:HOH:O	2.02	0.59
1:A:29:VAL:O	1:A:29:VAL:HG22	2.03	0.58
2:B:73:ALA:O	2:B:77:ILE:HG12	2.03	0.57
2:B:13:VAL:HG22	2:B:16:VAL:HB	1.85	0.57
2:B:177:HIS:HD2	4:B:2040:HOH:O	1.86	0.57
2:B:174:ASP:OD2	4:B:2055:HOH:O	2.16	0.57
2:B:84:PHE:CE2	2:B:168:MET:HG3	2.40	0.57
1:A:28:ARG:C	1:A:30:SER:H	2.08	0.57
2:B:63:THR:HB	2:B:169:SER:OG	2.05	0.56
2:B:60:ALA:HB3	2:B:172:ILE:HG12	1.87	0.55
2:B:63:THR:CG2	4:B:2008:HOH:O	2.56	0.53
2:B:66:LEU:HD12	2:B:168:MET:HE3	1.89	0.52
2:B:76:VAL:CG1	2:B:168:MET:HE1	2.37	0.51
1:A:28:ARG:O	1:A:31:ASN:OD1	2.27	0.51
1:A:9:THR:HG23	2:B:7:SER:OG	2.10	0.51
2:B:154:ALA:O	2:B:157:ILE:HG12	2.11	0.51
1:A:10:LYS:HD3	2:B:10:LYS:HD3	1.93	0.51
1:A:70:ARG:HD2	1:A:164:THR:HB	1.93	0.50
1:A:7:SER:OG	2:B:9:THR:HG23	2.11	0.50
2:B:86:GLU:HG2	2:B:166:ILE:HG22	1.93	0.50
2:B:7:SER:HB3	2:B:63:THR:HG23	1.94	0.49

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:36:ILE:HG13	2:B:39:LYS:HD2	1.96	0.48
3:A:1219:FAD:H2'	3:A:1219:FAD:H9	1.95	0.48
1:A:182:ARG:HD3	1:A:187:THR:HG21	1.96	0.48
1:A:86:GLU:HG2	1:A:166:ILE:HG22	1.95	0.47
1:A:39:LYS:HB2	4:A:2018:HOH:O	2.15	0.47
1:A:176:ILE:O	1:A:180:GLU:HB2	2.16	0.46
1:A:204:LYS:HB3	1:A:204:LYS:HE3	1.82	0.46
3:B:1217:FAD:H1'2	3:B:1217:FAD:O4'	2.19	0.43
1:A:158:LEU:HB3	1:A:162:THR:HG21	2.01	0.42
3:B:1217:FAD:H9	3:B:1217:FAD:H2'	2.00	0.42
1:A:189:ARG:HD3	1:A:189:ARG:N	2.34	0.41
1:A:20:GLU:OE1	4:A:2012:HOH:O	2.22	0.41
1:A:61:PHE:CE1	1:A:169:SER:HB3	2.55	0.41
1:A:84:PHE:O	2:B:55:SER:HB2	2.19	0.41
1:A:50:ARG:HD3	1:A:51:HIS:CE1	2.56	0.41
2:B:76:VAL:HG12	2:B:168:MET:HE2	1.92	0.41
2:B:52:LYS:HE3	4:B:2032:HOH:O	2.21	0.41
2:B:143:LYS:HB3	4:B:2052:HOH:O	2.20	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	173/217 (80%)	168 (97%)	4 (2%)	1 (1%)	27	33
2	B	169/217 (78%)	169 (100%)	0	0	100	100
All	All	342/434 (79%)	337 (98%)	4 (1%)	1 (0%)	43	53

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	29	VAL

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	154/190 (81%)	138 (90%)	16 (10%)	8	9
2	B	154/190 (81%)	139 (90%)	15 (10%)	9	10
All	All	308/380 (81%)	277 (90%)	31 (10%)	8	10

All (31) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1	MET
1	A	35	GLN
1	A	44	LEU
1	A	50	ARG
1	A	77	LEU
1	A	125	TRP
1	A	132	LYS
1	A	158	LEU
1	A	160	LEU
1	A	162	THR
1	A	165	THR
1	A	172	ILE
1	A	173	ARG
1	A	187	THR
1	A	189	ARG
1	A	194	LEU
2	B	13	VAL
2	B	36	ILE
2	B	44	LEU
2	B	45	LEU
2	B	63	THR
2	B	131	GLU
2	B	132	LYS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
2	B	152	GLU
2	B	155	ARG
2	B	158	LEU
2	B	166	ILE
2	B	187	THR
2	B	189	ARG
2	B	194	LEU
2	B	200	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	17	ASN
1	A	34	ASN
1	A	35	GLN
1	A	85	GLN
1	A	196	ASN
1	A	217	HIS
2	B	17	ASN
2	B	34	ASN
2	B	136	GLN
2	B	177	HIS
2	B	185	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
1	CME	A	48	1	9,9,10	0.86	0	6,9,11	1.49	1 (16%)
2	CME	B	48	2	9,9,10	0.70	0	6,9,11	2.43	3 (50%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CME	A	48	1	-	0/5/8/10	0/0/0/0
2	CME	B	48	2	-	0/5/8/10	0/0/0/0

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	48	CME	CZ-CE-SD	-2.03	106.33	113.37
1	A	48	CME	CE-SD-SG	2.73	116.65	103.47
2	B	48	CME	CE-SD-SG	2.78	116.88	103.47
2	B	48	CME	CB-SG-SD	4.46	112.52	103.85

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	FAD	A	1219	-	51,58,58	1.34	6 (11%)	57,89,89	2.01	6 (10%)
3	FAD	B	1217	-	51,58,58	1.38	7 (13%)	57,89,89	2.17	10 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	FAD	A	1219	-	-	0/28/50/50	0/6/6/6
3	FAD	B	1217	-	-	0/28/50/50	0/6/6/6

All (13) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	1217	FAD	C2A-N1A	2.25	1.38	1.33
3	B	1217	FAD	C5X-N5	2.29	1.38	1.35
3	A	1219	FAD	C2A-N1A	2.48	1.38	1.33
3	B	1217	FAD	C1'-N10	2.58	1.51	1.48
3	A	1219	FAD	C5X-N5	2.60	1.39	1.35
3	B	1217	FAD	C4-N3	3.01	1.38	1.33
3	A	1219	FAD	C10-N1	3.26	1.37	1.33
3	A	1219	FAD	C4-N3	3.31	1.39	1.33
3	B	1217	FAD	C4X-N5	3.64	1.38	1.33
3	B	1217	FAD	C2A-N3A	3.86	1.38	1.32
3	A	1219	FAD	C4X-N5	3.98	1.39	1.33
3	A	1219	FAD	C2A-N3A	4.18	1.38	1.32
3	B	1217	FAD	C10-N1	4.36	1.39	1.33

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	1217	FAD	N3A-C2A-N1A	-11.30	119.19	128.86
3	A	1219	FAD	N3A-C2A-N1A	-11.01	119.44	128.86
3	B	1217	FAD	O4'-C4'-C5'	-4.96	98.95	110.02
3	A	1219	FAD	C4X-C4-N3	-2.98	119.24	123.47
3	B	1217	FAD	C4X-C4-N3	-2.41	120.05	123.47
3	B	1217	FAD	C1B-N9A-C4A	-2.25	122.74	126.64
3	B	1217	FAD	C4A-C5A-N7A	-2.04	107.44	109.41

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	1217	FAD	C5X-C9A-N10	2.05	119.28	117.71
3	A	1219	FAD	C4-C4X-N5	2.33	121.32	118.70
3	B	1217	FAD	O2P-P-O5'	2.46	119.17	107.75
3	A	1219	FAD	C4X-N5-C5X	2.76	119.65	116.76
3	B	1217	FAD	C4X-N5-C5X	2.98	119.88	116.76
3	A	1219	FAD	C1'-N10-C9A	4.37	122.20	118.31
3	B	1217	FAD	C1'-N10-C9A	5.04	122.79	118.31
3	B	1217	FAD	C4-N3-C2	5.76	120.05	115.14
3	A	1219	FAD	C4-N3-C2	5.98	120.24	115.14

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1219	FAD	1	0
3	B	1217	FAD	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
2	B	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	B	150:ALA	C	151:LYS	N	2.54

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	179/217 (82%)	0.38	18 (10%) 7 10	18, 33, 84, 94	0
2	B	178/217 (82%)	0.41	15 (8%) 11 15	18, 34, 89, 92	1 (0%)
All	All	357/434 (82%)	0.39	33 (9%) 9 12	18, 33, 88, 94	1 (0%)

All (33) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	31	ASN	6.1
1	A	32	PRO	6.0
1	A	89	GLN	5.1
1	A	148	GLY	5.0
2	B	145	LEU	4.9
2	B	127	ALA	4.8
1	A	33	GLU	4.5
1	A	124	THR	4.4
2	B	126	TRP	4.3
1	A	34	ASN	4.1
1	A	35	GLN	3.9
2	B	149	ILE	3.7
2	B	148	GLY	3.6
2	B	146	GLU	3.4
1	A	126	TRP	3.2
1	A	30	SER	3.2
1	A	1	MET	3.1
1	A	139	GLU	3.1
2	B	125	TRP	3.1
1	A	146	GLU	3.0
2	B	150	ALA	2.8
1	A	39	LYS	2.5
1	A	125	TRP	2.4
2	B	128	THR	2.4

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
2	B	143	LYS	2.4
1	A	156	PHE	2.4
2	B	155	ARG	2.3
2	B	134	TYR	2.3
2	B	189	ARG	2.2
1	A	143	LYS	2.2
2	B	139	GLU	2.1
2	B	153	CYS	2.1
1	A	217	HIS	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	CME	B	48	10/11	0.94	0.15	24,25,37,40	0
1	CME	A	48	10/11	0.96	0.12	32,33,44,47	0

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	FAD	B	1217	53/53	0.95	0.14	16,25,32,32	0
3	FAD	A	1219	53/53	0.96	0.14	15,21,25,27	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.